

N° d'ordre 4221

# THÈSE

en vue de l'obtention du titre de

*Docteur de l'Université des Sciences et Technologies de Lille*

et de

*Docteur de l'Université de Vilnius*

présentée par

Vaidotas ZEMLYS

Discipline : Mathématiques Appliquées

Spécialité : Probabilités

## **Principe d'invariance pour processus de sommation multiparamétriques et applications**

Date de soutenance : 26 septembre 2008

### Composition du Jury

|                       |                      |                                   |
|-----------------------|----------------------|-----------------------------------|
| Rapporteurs :         | Davar KHOSHNEVISAN,  | Professeur, Université de l'Utah  |
|                       | Vygantas PAULASKAS,  | Professeur, Université de Vilnius |
|                       | Dalibor VOLNÝ,       | Professeur, Université de Rouen   |
| Examineurs :          | Youri DAVYDOV,       | Professeur, Université de Lille I |
|                       | Anne PHILIPPE,       | Professeur, Université de Nantes  |
|                       | Donatas SURGAILIS,   | Professeur, Université de Vilnius |
| Directeurs de Thèse : | Alfredas RAČKAUSKAS, | Professeur, Université de Vilnius |
|                       | Charles SUQUET,      | Professeur, Université de Lille I |

**Thèse préparée à l'Université des Sciences et Technologies de Lille  
Laboratoire Paul Painlevé, UMR 8524 CNRS  
et à l'Université de Vilnius**

## Résumé

La thèse a pour objet de prouver le principe d'invariance dans des espaces de Hölder pour le processus de sommation multiparamétrique et d'utiliser ce résultat en détection de rupture dans des données de panel. On caractérise d'abord la convergence en loi dans un espace de Hölder, du processus de sommation multiparamétrique dans le cas d'un champ aléatoire i.i.d. d'éléments aléatoires centrés et de carré intégrable d'un espace de Hilbert séparable, par la finitude d'un certain moment faible dont l'ordre croît avec l'exposant de Hölder, depuis deux lorsque l'exposant est nul, jusqu'à l'infini lorsque l'exposant est un demi. Ensuite on considère les tableaux triangulaires centrés à valeurs réelles. On propose une construction adaptative du processus de sommation qui coïncide avec la construction classique pour le cas d'un seul paramètre. On prouve le théorème limite central fonctionnel hölderien pour ce processus. Le processus limite est gaussien sous certaines conditions de régularité pour les variances du tableau triangulaire, le drap de Wiener n'étant qu'un cas particulier. Enfin on fournit des applications de ces résultats théoriques en construisant des statistiques de détection de rupture épidémique dans un ensemble de données multi-indexées. On construit un test de détection d'un changement d'espérance dans un rectangle épidémique, trouve sa loi limite et donne des conditions pour sa consistance. On adapte notre statistique pour la détection de rupture du coefficient dans les modèles classiques de régression pour panel.

**Mots clés :** Théorème limite central fonctionnel, drap brownien à valeurs dans un espace de Hilbert, espace de Hölder, principe d'invariance, processus de sommation, tableau triangulaire, données panellisées, rupture structurelle, test CUSUM

**Classification JEL :** C13, C23.

**Classification AMS (2000) :** 60B12, 60F17, 60G60, 62G10, 62G20, 62J05

# Invariance principle for multiparameter summation processes and applications

## Abstract

The thesis is devoted to proving invariance principle in Hölder spaces for the multi-parameter summation process and then using this result to construct the tests for detecting structural breaks in panel data. First we characterize the weak convergence in Hölder space of multi-parameter summation process in the case of i.i.d. random field of square integrable centered random elements in the separable Hilbert space by the finiteness of certain weak moment, whose order increases with the Hölder exponent, turning to two, when exponent is zero and to infinity when exponent is one half. Next we consider real valued centered triangular arrays. We propose adaptive construction of the summation process which coincides with classical construction for the one parameter case. We prove the functional central limit theorem for this process in Hölder space. The limiting process is Gaussian under certain regularity condition for variances of the triangular array, Brownian sheet being a special case. Finally we provide some application of the theoretical results by constructing statistics for detecting an epidemic change in a given data with multi-dimensional indexes. We construct a test for detecting the change of the mean in a epidemic rectangle, find its asymptotic distribution and give the conditions for the consistency. We adapt our proposed statistic for detecting the change of the coefficient in the classical panel regression models.

**Key words :** Functional central limit theorem, Hilbert space valued Brownian sheet, Hölder space, invariance principle, summation process, triangular array, panel data, structural break, CUSUM test.

**JEL classification:** C13, C23.

**AMS (2000) classification:** 60B12, 60F17, 60G60, 62G10, 62G20, 62J05

*To my wife, Alina  
and my son, Eimantas*

# Résumé substantiel

Le principe d'invariance est un résultat de la théorie des probabilités affirmant qu'un certain processus construit sur des sommes partielles de variables aléatoires converge en loi et que le processus limite ne dépend pas de la loi de ces variables. Ce résultat est une généralisation du théorème limite central (désigné ci-dessous par son acronyme anglais CLT) et est quelquefois aussi appelé théorème limite central fonctionnel (acronyme anglais FCLT), puisque la convergence en loi considérée usuellement a lieu dans un certain espace fonctionnel.

Le premier résultat de ce type fut prouvé par Prokhorov[24] et Donsker [10]. Ils ont considéré le processus de sommation basé sur des variables aléatoires i.i.d. centrées  $X_1, \dots, X_n$  :

$$\xi_n(t) = S_{[nt]} + (nt - [nt])X_{[nt]+1}, \quad t \in [0, 1], \quad (1)$$

où  $S_k = X_1 + \dots + X_k$ , pour  $k = 1, \dots, n$  et  $S_0 = 0$ . Les trajectoires de ce processus étant des lignes polygonales, le processus peut être vu comme un élément aléatoire de l'espace de fonctions continues  $C([0, 1])$ . Donsker et Prokhorov ont prouvé qu'un tel processus (normé par  $\sqrt{n}$  et  $\sigma = \sqrt{\mathbf{E} X_1^2}$ ) converge en loi quand  $n$  tend vers l'infini, vers le mouvement brownien  $W(t)$  – processus gaussien centré de fonction de covariance  $\mathbf{E} W(t)W(s) = t \wedge s$ . La condition nécessaire et suffisante pour cette convergence est que  $\sigma < \infty$ , i.e. l'existence du moment d'ordre 2. On dit qu'une suite d'éléments aléatoires d'un espace métrique séparable  $B$  converge en loi vers un élément aléatoire  $Y$ , si  $\mathbf{E} f(Y_n) \rightarrow \mathbf{E} f(Y)$ , pour toute fonctionnelle  $f$  continue bornée sur  $B$ . Il est facile de voir pourquoi ce résultat est une généralisation du CLT puisque  $\xi_n(1) = S_n$ .

Le FCLT, combiné avec le théorème de conservation de la convergence en loi par image continue, implique la convergence en loi de  $g(\sigma^{-1}n^{-1/2}\xi_n)$  vers  $g(W)$  pour toute fonctionnelle continue  $g$ . Ceci procure de multiples applications statistiques. Un exemple classique d'une telle application est un test de détection d'une rupture dans l'espérance d'un échantillon. Une des

statistiques pour un tel test est

$$Q = \max_{1 \leq k \leq n} |S_k - \frac{k}{n} S_n|.$$

La statistique  $Q$  s'écrit  $Q = g(\xi_n)$  pour la fonctionnelle  $g(x) = \sup_{0 \leq t \leq 1} |x(t)|$ . Puisque  $g$  est continue sur l'espace  $C([0, 1])$ , le FCLT nous fournit sous l'hypothèse nulle d'un échantillon i.i.d. de variance 1 et d'espérance fixée, la convergence en loi

$$n^{-1/2} Q \xrightarrow{D} \sup_{0 < t < 1} |W(t) - tW(1)|.$$

Sous l'hypothèse alternative d'existence d'un point  $k^*$  tel que  $\mathbf{E} X_i = \mu_0 + (\mu_1 - \mu_0)\mathbf{1}(i > k^*)$ , la statistique  $n^{-1/2} Q$  tend vers l'infini, donc le test est consistant. Si on considère l'alternative dite « épidémique »,  $\mathbf{E} X_i = \mu_0 + (\mu_1 - \mu_0)\mathbf{1}(k^* < i \leq l^*)$ , alors on peut utiliser la statistique

$$R = \max_{1 \leq k < l \leq n} \frac{|S_l - S_k - \frac{l-k}{n} S_n|}{[(l-k)/n]^\alpha}.$$

Mais la fonctionnelle correspondante du processus de sommation :

$$g(x) = \sup_{0 < |t-s| < 1} \frac{|x(t) - x(s)|}{|t-s|^\alpha}$$

n'est alors plus continue sur  $C([0, 1])$ . Elle est néanmoins continue sur l'espace de Banach  $(H_\alpha^o, \|\cdot\|_\alpha)$ ,  $0 < \alpha < 1$ , des fonctions  $x : [0, 1] \rightarrow \mathbb{R}$  telles que

$$\|x\|_\alpha := |x(0)| + w_\alpha(x, 1) < \infty,$$

où

$$w_\alpha(x, \delta) := \sup_{0 < |t-s| \leq \delta} \frac{|x(t) - x(s)|}{|t-s|^\alpha} \xrightarrow{\delta \rightarrow 0} 0.$$

Pour obtenir la loi limite de la statistique  $R$ , nous avons alors besoin d'un FCLT dans  $H_\alpha^o$ . Notons que l'espace  $H_\alpha^o$  a une topologie plus fine que  $C([0, 1])$ , donc son espace de fonctionnelles continues est plus « riche ». Notons aussi qu'en raison du module de continuité du mouvement brownien, la restriction  $0 < \alpha < 1/2$  s'impose. L'intérêt d'une statistique hölderienne comme  $R$  est de permettre la détection de « courtes » épidémies, typiquement  $k^* - l^*$  de l'ordre de  $n^\gamma$  pour  $0 < \gamma < 1/2$ , le choix de  $\gamma$  étant lié à l'intégrabilité des  $X_i$ , alors que sans le dénominateur dans  $R$ , on reste limité à la détection d'épidémies de longueur au moins  $n^{1/2}$ . Dans ce travail, nous nous intéressons à l'extension de cet avantage aux processus de sommation multiparamétriques. Ces processus

apparaissent dans l'étude des sommes partielles de variables aléatoires multi-indexées. Outre la curiosité mathématique, l'intérêt de l'étude de ce type de variables est qu'il est parfois plus commode d'attacher plusieurs indices à une observation. Par exemple, si on mesure une certaine propriété pour un échantillon d'individus au cours du temps, il est naturel d'attribuer deux indices aux observations, le numéro de l'individu et l'instant d'observation. On appelle ceci données longitudinales ou données de panel.

Nous donnons maintenant une définition du processus de sommation multiparamétrique plus générale que celle utilisée dans ce travail, de façon à pouvoir situer nos résultats dans leur contexte historique. Notons  $|A|$  la mesure de Lebesgue du sous-ensemble borélien  $A$  de  $\mathbb{R}^d$ . Pour une collection  $\mathcal{A}$  de parties boréliennes de  $[0, 1]^d$ , le processus de sommation  $\{\xi_n(A); A \in \mathcal{A}\}$  basé sur le champ aléatoire  $\{X_j, \mathbf{j} \in \mathbb{N}^d\}$ , de variables aléatoires réelles i.i.d. centrées est défini par

$$\xi_n(A) = \sum_{1 \leq \mathbf{j} \leq \mathbf{n}} |R_{\mathbf{n}, \mathbf{j}}|^{-1} |R_{\mathbf{n}, \mathbf{j}} \cap A| X_j, \quad (2)$$

où  $\mathbf{j} = (j_1, \dots, j_d)$ ,  $\mathbf{n} = (n_1, \dots, n_d)$ ,  $R_{\mathbf{n}, \mathbf{j}}$  est le « rectangle »

$$R_{\mathbf{n}, \mathbf{j}} := \left[ \frac{j_1 - 1}{n_1}, \frac{j_1}{n_1} \right) \times \dots \times \left[ \frac{j_d - 1}{n_d}, \frac{j_d}{n_d} \right)$$

et la condition d'indexation «  $\mathbf{1} \leq \mathbf{j} \leq \mathbf{n}$  » s'entend composante par composante :  $1 \leq j_1 \leq n_1, \dots, 1 \leq j_d \leq n_d$ . Notons que pour  $\mathcal{A} = \{[0, t], t \in [0, 1]\}$ , le processus ainsi défini coïncide avec celui défini par (1).

En munissant la collection  $\mathcal{A}$  d'une certaine pseudo-métrique  $\delta$ , on peut définir l'espace  $C(\mathcal{A})$  des fonctions réelles continues sur  $\mathcal{A}$ , muni de la norme

$$\|f\|_{\mathcal{A}} := \sup_{A \in \mathcal{A}} |f(A)|.$$

Les semi-métriques usuelles sont  $\delta(A, B) = \sqrt{|A \Delta B|}$ , ou  $\delta(A, B) = \sqrt{m(A \Delta B)}$ , pour  $A, B \in \mathcal{A}$ , où  $m$  est une mesure de probabilité sur la tribu borélienne de  $[0, 1]^d$ . Les trajectoires du processus  $\xi_n(A)$  vivent alors dans l'espace  $C(\mathcal{A})$ . Lorsque  $\mathcal{A}$  est totalement borné relativement à  $\delta$ ,  $C(\mathcal{A})$  est un espace de Banach séparable. Le processus  $\{\xi_n(A), A \in \mathcal{A}\}$  est alors un élément aléatoire d'un espace métrique séparable et il est possible de rechercher un FCLT pour un tel processus. La limite en loi d'un tel processus de sommation est la généralisation du mouvement brownien appelée *drap brownien* ou *drap de Wiener*. C'est un processus gaussien centré  $W$ , indexé par  $\mathcal{A}$ , à trajectoires dans  $C(\mathcal{A})$  et vérifiant

$$\mathbf{E} W(A)W(B) = |A \cap B|, \quad A, B \in \mathcal{A}.$$

L'existence d'un tel processus est prouvée au prix de quelques restrictions sur la collection  $\mathcal{A}$ , qui s'expriment usuellement par une certaine condition sur son entropie métrique.

Considérons la famille de rectangles

$$\mathcal{Q}_d := \left\{ [0, t_1] \times \cdots \times [0, t_d]; \mathbf{t} = (t_1, \dots, t_d) \in [0, 1]^d \right\}. \quad (3)$$

Le premier FCLT dans  $C(\mathcal{A})$  pour  $\{\xi_n(A); A \in \mathcal{A}\}$  fut établi par Kuelbs [19] dans le cas particulier de la famille  $\mathcal{A} = \mathcal{Q}_d$  sous certaines hypothèses restrictives de moments et par Wichura [42] sous l'hypothèse de variance finie.

En 1983, Pyke [25] a établi un FCLT dans  $C(\mathcal{A})$ , pourvu que  $\mathcal{A}$  satisfasse la condition d'entropie avec crochets. Cependant son résultat supposait des conditions de moment dépendant de la taille de la collection  $\mathcal{A}$ . Bass [3] et simultanément Alexander et Pyke [1], ont étendu ce résultat aux champs aléatoires i.i.d. de variance finie. Des développements ultérieurs concernent l'affaiblissement des conditions d'entropie sur la collection  $\mathcal{A}$ , Ziegler [43], et le remplacement de l'hypothèse i.i.d. sur le champ aléatoire  $\{X_n, \mathbf{n} \in \mathbb{N}^d\}$  par des hypothèses de dépendance, cf. Dedecker [8], El Machkouri et Ouchti [12] entre autres.

Le FCLT dans les espaces de Hölder n'a pas été aussi intensivement étudié. En 1962, Lamperti [20] obtenait le premier théorème limite central fonctionnel dans les espaces de Banach séparables  $H_\alpha^o$ ,  $0 < \alpha < 1/2$ . En supposant que  $\mathbf{E} |X_1|^q < \infty$  pour un  $q > 2$ , il prouvait la convergence en loi de  $n^{-1/2}\xi_n$  vers  $W$  dans l'espace de Hölder  $H_\alpha^o$  pour tout  $\alpha < 1/2 - 1/q$ . Račkauskas et Suquet dans [31], voir aussi [28], ont obtenu une condition nécessaire et suffisante pour le théorème limite central fonctionnel de Lamperti, à savoir que pour  $0 < \alpha < 1/2$ ,  $n^{-1/2}\xi_n$  converge en loi dans  $H_\alpha^o$  vers  $W$  si et seulement si

$$\lim_{t \rightarrow \infty} t^{p(\alpha)} P(|X_1| > t) = 0,$$

où

$$p(\alpha) := \frac{1}{\frac{1}{2} - \alpha}. \quad (4)$$

L'espace de Hölder général  $H_\rho(\mathcal{A})$  pour la collection  $\mathcal{A}$  et la fonction de poids  $\rho$  est défini comme le sous-espace de  $C(\mathcal{A})$  constitué des fonctions pour lesquelles

$$\sup_{0 < \delta(A, B) < 1} \frac{|x(A) - x(B)|}{\rho(\delta(A, B))} < \infty.$$

Pour l'existence de  $W$  dans un tel espace, voir Dudley [11] et Erickson [13]. Dans le cas où  $\rho(h) = h^\alpha$ , Erickson [13] prouve que  $\alpha$  ne peut excéder  $1/2$

et décroît quand l'entropie de  $\mathcal{A}$  croît. Les résultats les plus généraux pour le FCLT dans les espaces  $H_\rho(\mathcal{A})$  sont fournis par Erickson [13] qui montre que si  $\mathbf{E}|X_j|^q < \infty$  pour un  $q > 2$ , le FCLT est vérifié dans  $H_\rho(\mathcal{A})$  pour un  $\rho$  dépendant de  $q$  et des propriétés de  $\mathcal{A}$ . Pour  $d = 1$  et la classe  $\mathcal{A}$  des intervalles  $[0, t]$ ,  $0 \leq t \leq 1$ , les résultats d'Erickson coïncident avec ceux de Lamperti [20], tandis que son cas  $d > 1$  requiert des moments d'ordre  $q > dp(\alpha)$  avec le même  $p(\alpha)$  que dans (4).

La classe de sous-ensembles  $\mathcal{Q}_d$  est isomorphe au cube unité  $[0, 1]^d$  par la bijection  $[0, \mathbf{t}] \leftrightarrow \mathbf{t}$ . Il est alors naturel de traiter le processus  $\xi_n(A)$  comme une fonction à plusieurs variables ou « multiparamétrique »  $\xi_n(\mathbf{t})$ . Račkauskas et Suquet ont étudié intensivement les espaces hölderiens de fonctions multiparamétriques et mis au point tous les outils pour prouver le FCLT dans de tels espaces. Au vu de la généralisation du principe d'invariance de Lamperti, la question se pose naturellement d'un résultat analogue dans le cas multiparamétrique. Cette question est complètement résolue dans la présente contribution.

Le premier chapitre de la thèse collecte tous les outils et résultats utiles pour prouver le théorème limite central fonctionnel dans des espaces de Hölder. Le processus de sommation  $\{\xi_n, \mathbf{n} \in \mathbb{N}^d\}$  est en réalité un filet (*net*), alors que tous les critères usuels concernent des suites. Donc dans ce chapitre, nous adaptons tous les résultats existants au cas des filets au lieu des suites.  $\mathbb{H}$  désignant un espace de Hilbert séparable, nous commençons par définir l'espace de Hölder  $H_\alpha^o(\mathbb{H})$  des fonctions multiparamétriques à valeurs hilbertiennes comme l'espace vectoriel des fonctions  $x : [0, 1]^d \rightarrow \mathbb{H}$  telles que

$$\|x\|_\alpha := \|x(0)\| + \omega_\alpha(x, 1) < \infty,$$

avec

$$\omega_\alpha(x, \delta) := \sup_{0 < |\mathbf{t} - \mathbf{s}| \leq \delta} \frac{\|x(\mathbf{t}) - x(\mathbf{s})\|}{|\mathbf{t} - \mathbf{s}|^\alpha} \xrightarrow{\delta \rightarrow 0} 0.$$

Muni de la norme  $\|\cdot\|_\alpha$ ,  $H_\alpha^o(\mathbb{H})$  est un espace de Banach séparable, voir [27] ou [33]. Nous rappelons alors tous les faits utiles concernant cet espace, d'après Račkauskas et Suquet [27]. Ceci nous permet de prouver le critère d'équitension suivant pour les filets d'éléments aléatoires de  $H_\alpha^o(\mathbb{H})$ .

**Théorème 1** *Soient  $\{\zeta_n, \mathbf{n} \in \mathbb{N}^d\}$  et  $\zeta$  des éléments aléatoires de l'espace  $H_\alpha^o(\mathbb{H})$ . Supposons satisfaites les conditions suivantes.*

- i) *Pour tout point dyadique  $\mathbf{t} \in [0, 1]^d$ , le filet d'éléments aléatoires de  $\mathbb{H}$   $\zeta_n(\mathbf{t})$  est asymptotiquement équitendu dans  $\mathbb{H}$ .*
- ii) *Pour tout  $\varepsilon > 0$ ,*

$$\lim_{J \rightarrow \infty} \limsup_{m(\mathbf{n}) \rightarrow \infty} P(\sup_{j \geq J} 2^{\alpha j} \max_{v \in V_j} \|\lambda_{j,v}(\zeta_n)\| > \varepsilon) = 0.$$

Alors le filet  $\zeta_{\mathbf{n}}$  est asymptotiquement équitendu dans l'espace  $H_{\alpha}^o(\mathbb{H})$ .

Dans cet énoncé,  $m(\mathbf{n})$  désigne le minimum des composantes du vecteur  $\mathbf{n}$  et la quantité  $\lambda_{j,v}(\zeta_{\mathbf{n}})$  provient d'une décomposition de Schauder particulière de l'espace  $H_{\alpha}^o(\mathbb{H})$  et est définie comme une différence seconde de  $\zeta_{\mathbf{n}}$  en certains points dyadiques.

Le chapitre se poursuit par la définition du processus limite pour le FCLT dans l'espace  $H_{\alpha}^o(\mathbb{H})$ , à savoir un drap de Wiener à valeurs dans  $\mathbb{H}$  avec un opérateur de covariance  $\Gamma$ . Il est défini comme un processus gaussien centré à valeurs dans  $\mathbb{H}$ , indexé par  $[0, 1]^d$  et vérifiant

$$\mathbf{E} \langle W(\mathbf{t}), x \rangle \langle W(\mathbf{s}), y \rangle = (t_1 \wedge s_1) \dots (t_d \wedge s_d) \langle \Gamma x, y \rangle$$

pour tous  $\mathbf{t}, \mathbf{s} \in [0, 1]^d$  et tous  $x, y \in \mathbb{H}$ . Comme ce processus n'est pas d'un emploi usuel dans la littérature, nous donnons une preuve de son existence.

Ensuite, on rappelle les inégalité de Rosenthal et de Doob qui sont utiles pour prouver l'équitension. Pour des éléments aléatoires  $(Y_i)_{i \in I}$  indépendants d'un espace de Hilbert, l'inégalité de Rosenthal s'écrit

$$\mathbf{E} \left\| \sum_{i \in I} Y_i \right\|^q \leq C'_q \left( \mathbf{E} \left\| \sum_{i \in I} G(Y_i) \right\|^q + \sum_{i \in I} \mathbf{E} \|Y_i\|^q \right),$$

et peut donc s'appliquer pour des sommes finies indexées par un ensemble quelconque. La généralisation de l'inégalité de Doob pour des sommes de variables aléatoires réelles multiparamétriques est donnée par Khoshnevisan. Nous montrons qu'elle s'étend aussi au cas d'éléments aléatoires hilbertiens :

$$\mathbf{E} \max_{0 \leq j \leq \mathbf{n}} \|S_j\|^p \leq \left( \frac{p}{p-1} \right)^{dp} \mathbf{E} \|S_{\mathbf{n}}\|^p,$$

pour tous  $\mathbf{n} \in \mathbb{N}^d$  et  $p > 1$ .

Le second chapitre donne les propriétés détaillées du processus de sommation (2). Il se trouve que ce processus admet l'utile représentation suivante.

**Proposition 2** Pour  $\mathbf{t} \in [0, 1]^d$ , notons  $\mathbf{s} = \{\mathbf{nt}\}$  et représentons les sommets  $V(\mathbf{u})$  du rectangle  $R_{\mathbf{n}, [\mathbf{nt}] + 1}$  par

$$V(\mathbf{u}) := \frac{[\mathbf{nt}]}{\mathbf{n}} + \frac{\mathbf{u}}{\mathbf{n}}, \quad \mathbf{u} \in \{0, 1\}^d.$$

On peut alors écrire  $\mathbf{t}$  comme barycentre de ces  $2^d$  sommets munis des masses  $w(\mathbf{u}) \geq 0$  dépendant de  $\mathbf{t}$ , i.e.,

$$\mathbf{t} = \sum_{\mathbf{u} \in \{0, 1\}^d} w(\mathbf{u}) V(\mathbf{u}), \quad \text{avec} \quad \sum_{\mathbf{u} \in \{0, 1\}^d} w(\mathbf{u}) = 1,$$

où

$$w(\mathbf{u}) = \prod_{l=1}^d s_l^{u_l} (1 - s_l)^{1-u_l}.$$

À partir de cette représentation, définissons pour une collection  $\{X_{\mathbf{i}}, \mathbf{1} \leq \mathbf{i} \leq \mathbf{n}\}$  d'éléments aléatoires d'un espace de Banach, le champ aléatoire  $\xi_n^*$  par

$$\xi_n^*(\mathbf{t}) = \sum_{\mathbf{u} \in \{0,1\}^d} w(\mathbf{u}) S_{[\mathbf{n}\mathbf{t}]+\mathbf{u}}, \quad \mathbf{t} \in [0, 1]^d.$$

Alors  $\xi_n^*$  coïncide avec le processus de sommation défini par (2).

Cette proposition permet d'obtenir la majoration suivante de l'expression qui apparaît dans le critère d'équitension du théorème 1, ii) :

$$\sup_{j \geq J} 2^{\alpha j} \max_{v \in V_j} \|\lambda_{j,v}(\xi_n)\| \leq 3^d \sum_{m=1}^d \left( \max_{J \leq j \leq \log n_m} 2^{\alpha j} \max_{0 \leq k < 2^j} \psi_n^{(m)}(t_{k+1}, t_k) + n_m^\alpha Z_n^{(m)} \right)$$

où

$$\psi_n^{(m)}(t', t) := \max_{\mathbf{1}-m \leq \mathbf{k}-m \leq \mathbf{n}-m} \left\| \sum_{k_m=[n_m t]+1}^{[n_m t']} \Delta_{k_m}^{(m)} S_{\mathbf{k}} \right\|, \quad Z_n^{(m)} := \max_{1 \leq \mathbf{k} \leq \mathbf{n}} \|\Delta_{k_1}^{(m)} S_{\mathbf{k}}\|.$$

Les opérateurs de différence  $\Delta$  utilisés dans ces formules sont définis par

$$\Delta_k^{(i)} S_j := S_{(j_1, \dots, j_{i-1}, k, j_{i+1}, \dots, j_d)} - S_{(j_1, \dots, j_{i-1}, k-1, j_{i+1}, \dots, j_d)}.$$

L'obtention de ce majorant n'utilise aucune hypothèse sur la loi ou la dépendance des variables aléatoires concernées, il peut donc être réutilisé pour prouver un FCLT dans d'autres situations que le cas i.i.d.

Le principe d'invariance dans l'espace  $H_\alpha^o(\mathbb{H})$  est établi dans le chapitre 3.

**Théorème 3** Pour  $0 < \alpha < 1/2$ , posons  $p = p(\alpha) := 1/(1/2 - \alpha)$ . Pour  $d \geq 2$ , soit  $\{X_{\mathbf{i}}; \mathbf{i} \in \mathbb{N}^d, \mathbf{i} \geq \mathbf{1}\}$  une collection i.i.d. d'éléments aléatoires centrés et de carré intégrable de l'espace de Hilbert séparable  $\mathbb{H}$  et soit  $\xi_n$  le processus de sommation défini par

$$\xi_n(\mathbf{t}) = \sum_{\mathbf{i} \leq \mathbf{n}} \pi(\mathbf{n}) \left| \left[ \frac{\mathbf{i} - \mathbf{1}}{\mathbf{n}}, \frac{\mathbf{i}}{\mathbf{n}} \right) \cap [0, \mathbf{t}] \right| X_{\mathbf{i}}.$$

Notons  $W$  un drap brownien à valeurs dans  $\mathbb{H}$  avec le même opérateur de covariance que  $X_{\mathbf{1}}$ . Alors la convergence en loi

$$\pi(\mathbf{n})^{-1/2} \xi_n \xrightarrow[\mathfrak{m}(\mathbf{n}) \rightarrow \infty]{H_\alpha^o(\mathbb{H})} W$$

a lieu si et seulement si

$$\pi(\mathbf{n})P(\|X_1\| > n_m^{1/p} \pi(\mathbf{n}_{-m})^{1/2}) \xrightarrow{m(\mathbf{n}) \rightarrow \infty} 0, \quad (5)$$

pour  $m = 1, \dots, d$ , où l'on a noté  $\pi(\mathbf{n})$  le produit des composantes du multiindice  $\mathbf{n}$  et  $\pi(\mathbf{n}_{-m})$  celui de toutes les composantes sauf  $n_m$ .

Bien que la condition (5) paraisse assez technique, nous montrons qu'elle équivaut à la finitude du moment faible d'ordre  $p$  de  $X_1$ , i.e.

$$\sup_{t>0} t^p P(\|X_1\| > t) < \infty.$$

Ce nous donne immédiatement l'amélioration de résultats d'Erickson, où la condition pour FTCL est l'existence  $\mathbf{E}|X_1|^q < \infty$  pour  $q > dp$ .

En outre, l'utilité de (5) est de montrer quel type de condition de moment est requise si on considère une convergence plus faible que  $m(\mathbf{n}) \rightarrow \infty$ . Regardons par exemple la convergence restreinte aux  $\mathbf{n}$  de la forme  $\mathbf{n} = (n, \dots, n)$ , ce qui est le cadre usuel pour prouver le FCLT pour des processus de sommation généraux. Alors la condition de moment se réduit à

$$\lim_{t \rightarrow \infty} t^{\frac{2d}{d-2\alpha}} P(\|X_1\| > t) = 0,$$

ce qui est vérifié pour tout  $d > 1$  et tout  $0 < \alpha < 1/2$  dès que  $\mathbf{E}\|X_1\|^4 < \infty$ . Ceci contraste avec le résultat correspondant pour la convergence hölderienne de la ligne polygonale de Donsker-Prokhorov où nécessairement  $\mathbf{E}|X_1|^q < \infty$  pour tout  $q < p(\alpha)$ .

Les chapitres 2 et 3 explorent aussi la possibilité d'affaiblir l'hypothèse i.i.d. Quelques résultats sont prouvés dans le cadre des tableaux triangulaires. Pour des processus de sommation généraux (2), le cas des tableaux triangulaires a été étudié par Goldie et Greenwood [14], [15]. Ils ont utilisé la construction classique du processus de sommation, ce qui fait que leur résultat ne coïncide pas quand  $d = 1$ , avec celui de Prokhorov [24] pour le processus adaptatif  $\Xi_n$  indexé par  $[0, 1]$ , défini comme la ligne polygonale de sommets  $(b_n(k), S_n(k))$ , où  $b_n(k) = \mathbf{E} X_{n,1}^2 + \dots + \mathbf{E} X_{n,k}^2$ , où l'on suppose que  $b_n(k_n) = 1$ , et que les  $X_{n,k}$  sont des variables aléatoires indépendantes non identiquement distribuées.

Bickel et Wichura [5] ont tenté d'introduire une construction adaptative pour des processus de sommation généraux. Néanmoins, ils posent des conditions restrictives sur les variances des variables aléatoires du tableau triangulaire. Pour des variables aléatoires indépendantes centrées  $\{X_{n,ij}, 1 \leq$

$i \leq I_n, 1 \leq j \leq J_n\}$  dont les variances sont de la forme  $\mathbf{E} X_{n,ij} = a_{n,i} b_{n,j}$  avec  $\sum a_{n,i} = 1 = \sum b_{n,j}$ , ils définissent le processus de sommation comme

$$\zeta_n(t_1, t_2) = \sum_{i \leq A_n(t_1)} \sum_{j \leq B_n(t_2)} X_{n,ij},$$

où

$$A_n(t_1) = \max\{k : \sum_{i \leq k} a_{n,i} < t_1\}, \quad B_n(t_2) = \max\{l : \sum_{j \leq l} b_{n,j} < t_2\}.$$

Il est facile de voir que cette construction est la généralisation 2-dimensionnelle de la version processus de saut de la construction de Prokhorov. Bickel et Wichura ont prouvé la convergence du processus  $\zeta_n$  vers un drap de Wiener  $W$  dans l'espace  $D([0, 1]^2)$  des fonctions càdlàg à deux variables, en supposant la négligibilité infinitésimale pour  $a_{n,i}$  et  $b_{n,j}$  et la condition de Lindeberg pour les  $X_{n,ij}$ .

La généralisation de la construction de Prokhorov pour le processus de sommation multiparamétrique est proposée dans le chapitre 2. On y montre que la construction de Bickel et Wichura—cas particulier de celle que nous proposons—n'est pas suffisamment générale pour travailler avec tout tableau triangulaire.

La construction de Prokhorov ne place pas les sommes partielles en des points régulièrement espacés, mais en des points qui sont les variances des sommes partielles. Pour le processus de sommation multiparamétrique, les constructions standard placent les sommes partielles sur les points d'une grille rectangulaire uniforme. Malheureusement, il n'est pas possible de construire une grille rectangulaire sur le cube unité  $[0, 1]^d$  de telle sorte que ses points soient les variances des sommes partielles multiparamétriques. Nous proposons alors d'utiliser une grille où les coordonnées des points sur chaque axe sont les sommes des variances le long des autres axes. Pour clarifier cette idée, introduisons quelques notations. Pour le tableau triangulaire

$$(X_{n,\mathbf{k}}, \mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n), \quad \mathbf{n} \in \mathbb{N}^d,$$

posons

$$b_n(\mathbf{k}) := \sum_{j \leq \mathbf{k}} \sigma_{n,\mathbf{k}}^2,$$

où  $\sigma_{n,\mathbf{k}}^2 = \mathbf{E} X_{n,\mathbf{k}}^2$  (nous imposons au tableau d'être centré et que la somme des variances soit 1, i.e.  $b_n(\mathbf{k}_n) = 1$ ). Pour  $i = 1, \dots, d$  posons

$$b_i(k) := b_n(k_n^1, \dots, k_n^{i-1}, k, k_n^{i+1}, \dots, k_n^d),$$

et définissons le rectangle de grille comme

$$Q_{\mathbf{n}, \mathbf{k}} := \left[ b_1(k_1 - 1), b_1(k_1) \right) \times \cdots \times \left[ b_d(k_d - 1), b_d(k_d) \right).$$

Alors le processus de sommation que nous proposons est défini par

$$\Xi_{\mathbf{n}}(\mathbf{t}) = \sum_{1 \leq j \leq n} |Q_{\mathbf{n}, j}|^{-1} |Q_{\mathbf{n}, j} \cap [0, \mathbf{t}]| X_{\mathbf{n}, j}.$$

Comme pour le processus de sommation  $\xi_{\mathbf{n}}(\mathbf{t})$  nous pouvons donner une représentation de  $\Xi_{\mathbf{n}}(\mathbf{t})$  par interpolation affine des valeurs aux sommets du rectangle contenant  $\mathbf{t}$  et de l'utiliser pour obtenir une majoration de normes de Hölder séquentielles :

$$\sup_{j \geq J} 2^{\alpha j} \max_{v \in V_j} \|\lambda_{j,v}(\Xi_{\mathbf{n}})\| \leq 3^d \sum_{m=1}^d \left( \max_{j \geq J} 2^{j\alpha} \max_{r \in D_j} [\psi_{\mathbf{n}}^{(m)}(r, r^-) + \psi_{\mathbf{n}}^{(m)}(r, r^+)] + Z_{\mathbf{n}}^{(m)} \right)$$

où

$$\begin{aligned} \psi_{\mathbf{n}}(r, r^-)^{(m)} &:= \max_{\mathbf{k}_{-m} \leq \mathbf{k}_{\mathbf{n}, -m}} \left| \sum_{k_m = u_m(r^-) + 2}^{u_m(r)} \Delta_{k_m}^{(m)} S_{\mathbf{n}}(\mathbf{k}) \right|, \\ \psi_{\mathbf{n}}(r, r^+)^{(m)} &:= \max_{\mathbf{k}_{-m} \leq \mathbf{k}_{\mathbf{n}, -m}} \left| \sum_{k_m = u_m(r) + 2}^{u_m(r^+)} \Delta_{k_m}^{(m)} S_{\mathbf{n}}(\mathbf{k}) \right|, \\ Z_{\mathbf{n}}^{(m)} &:= \max_{1 \leq \mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} \frac{|\Delta_{k_m}^{(m)} S_{\mathbf{n}}(\mathbf{k})|}{(\Delta b_m(k_m))^\alpha}. \end{aligned}$$

Dans le cas  $d = 1$  le processus de sommation proposé coïncide avec la construction de Prokhorov. Les conditions pour un FCLT hölderien dans ce cas ont été données par Račkauskas et Suquet [30]. Ils combinent une certaine condition de moment avec la négligibilité infinitésimale du tableau triangulaire. Leur condition de moment étant — dans ce contexte — plus forte que celle de Lindeberg, la convergence des lois de dimension finie en découle immédiatement. Ce n'est plus le cas pour le processus  $\Xi_{\mathbf{n}}(\mathbf{t})$  quand  $d \geq 2$ . L'équitension est obtenue via les théorèmes suivants.

**Théorème 4** *Pour  $0 < \alpha < 1/2$ , posons  $p(\alpha) := 1/(1/2 - \alpha)$ . Si*

$$\max_{1 \leq l \leq d} \max_{1 \leq k_l \leq k_{\mathbf{n}}^l} \Delta b_l(k_l) \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty$$

*et pour un certain  $q > p(\alpha)$*

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} \sigma_{\mathbf{n}, \mathbf{k}}^{-2q\alpha} \mathbf{E} |X_{\mathbf{n}, \mathbf{k}}|^q = 0,$$

*alors le filet  $\{\Xi_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}\}$  est asymptotiquement équitendu dans l'espace  $H_\alpha^o([0, 1]^d)$ .*

Introduisons pour tout  $\tau > 0$  les variables aléatoires tronquées :

$$X_{n,k,\tau} := X_{n,k} \mathbf{1}\{|X_{n,k}| \leq \tau \sigma_{n,k}^{2\alpha}\}.$$

**Théorème 5** *Supposons que*

$$\max_{1 \leq l \leq d} \max_{1 \leq k_l \leq k_n^l} \Delta b_l(k_l) \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty \quad (6)$$

*et que les conditions suivantes soient vérifiées.*

*i. Pour tout  $\varepsilon > 0$ ,*

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq k \leq k_n} P(|X_{n,k}| \geq \varepsilon \sigma_{n,k}^{2\alpha}) = 0.$$

*ii. Pour tout  $\varepsilon > 0$ ,*

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq k \leq k_n} \mathbf{E} X_{n,k}^2 \mathbf{1}\{|X_{n,k}| \geq \varepsilon\} = 0.$$

*iii. Pour un certain  $q > 1/(1/2 - \alpha)$ ,*

$$\lim_{\tau \rightarrow 0} \lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq k \leq k_n} \sigma_{n,k}^{-2q\alpha} \mathbf{E} |X_{n,k,\tau}|^q = 0.$$

*Alors le filet  $\{\Xi_n, \mathbf{n} \in \mathbb{N}\}$  est asymptotiquement équitendu dans l'espace  $H_\alpha^o([0, 1]^d)$ .*

Ces théorèmes constituent une généralisation directe des conditions données par Račkauskas et Suquet.

Même si les conditions d'équitension ci-dessus assurent que le tableau triangulaire vérifie le CLT, elles ne nous disent rien sur la convergence des lois fini-dimensionnelles du processus  $\Xi_n(\mathbf{t})$ . Nous montrons que les limites des lois fini-dimensionnelles, si elles existent, sont gaussiennes, mais ce ne sont pas nécessairement les lois fini-dimensionnelles du drap de Wiener. Plus précisément, nous montrons que si on a la convergence

$$\mu_n(\mathbf{t}) := \sum_{1 \leq k \leq k_n} \mathbf{1}\{\mathbf{B}(\mathbf{k}) \in [0, \mathbf{t}]\} \sigma_{n,k}^2 \rightarrow \mu(\mathbf{t}), \text{ quand } m(\mathbf{n}) \rightarrow \infty$$

pour une certaine fonction  $\mu(\mathbf{t})$ , alors le processus limite  $G(\mathbf{t})$  est gaussien de covariance  $\mathbf{E} G(\mathbf{t})G(\mathbf{s}) = \mu(\mathbf{s} \wedge \mathbf{t})$ . Nous donnons des exemples montrant à quel type de comportement asymptotique on peut s'attendre.

**Exemple 1**  $X_{n,k} = a_{n,k}Y_k$ , où les  $\{Y_k, \mathbf{k} \leq \mathbf{k}_n\}$  sont i.i.d. gaussiennes standard, et

$$a_{n,k}^2 = \begin{cases} \frac{1}{10n^2}, & \text{pour } \mathbf{k} \leq (n, n) \\ \frac{3}{10n^2}, & \text{sinon.} \end{cases} \quad (7)$$

Le tableau triangulaire ainsi défini vérifie la condition (6), mais un calcul élémentaire montre que pour ce tableau

$$\begin{aligned} \mu_n(\mathbf{t}) \rightarrow \nu(\mathbf{t}) := & \frac{1}{10} \left( \frac{5}{2}t_1 \wedge 1 \right) \left( \frac{5}{2}t_2 \wedge 1 \right) + \frac{(5t_1 - 2) \vee 0}{10} \left( \frac{5}{2}t_2 \vee 1 \right) \\ & + \frac{(5t_2 - 2) \vee 0}{10} \left( \frac{5}{2}t_1 \vee 1 \right) + \frac{((5t_1 - 2) \vee 0)((5t_2 - 2) \vee 0)}{30}. \end{aligned}$$

Dans l'exemple suivant, la situation est encore pire puisque  $\mu_n(\mathbf{t})$  ne converge pour aucun  $\mathbf{t}$ .

**Exemple 2**  $X_{n,k} = b_{n,k}Y_k$  où les  $\{Y_k, \mathbf{k} \leq \mathbf{k}_n\}$  sont i.i.d. gaussiennes standard, et

$$b_{n,k}^2 = \begin{cases} \pi(\mathbf{k}_n)^{-1}, & \text{pour } \mathbf{n} = (2l-1, 2l-1), l \in \mathbb{N} \\ a_{n,k}^2, & \text{pour } \mathbf{n} = (2l, 2l), l \in \mathbb{N} \end{cases}$$

où les  $a_{n,k}$  sont définis comme dans (7).

Dans le quatrième chapitre, on applique les résultats théoriques ci-dessus à la construction de statistiques de détection de ruptures épidémiques dans un ensemble de données multi-indexées. On commence par la détection d'un changement d'espérance dans l'échantillon à double indice  $\{X_{ij}, 1 \leq i \leq n, 1 \leq j \leq m\}$ . Nous testons l'hypothèse nulle

$(H_0)$  : Les  $X_{ij}$  ont tous même espérance  $\mu_0$ ,  
contre l'hypothèse

$(H_A)$  : Il existe des entiers  $1 < a^* \leq b^* < n$ ,  $1 < c^* \leq d^* < m$  et une constante  $\mu_1 \neq \mu_0$  tels que

$$\mathbf{E} X_{ij} = \mu_0 + \mu_1 \mathbf{1} \left( (i, j) \in [a^*, b^*] \times [c^*, d^*] \cap \mathbb{N}^2 \right).$$

Nous appelons cette alternative *rectangle épidémique*. La statistique que nous proposons pour détecter ce type de rupture est la classique statistique du rapport de vraisemblance cf. Csörgő et Horváth [7], pondérée par une puissance du diamètre du rectangle épidémique :

$$DUI(n, m, \alpha) = \max_{\substack{1 \leq a < b \leq n \\ 1 \leq c < d \leq m}} \frac{|\Delta_{b-a}^{(1)} \Delta_{d-c}^{(2)} S_{bd} - (s_b - s_a)(t_d - t_c) S_{nm}|}{\max\{s_b - s_a, t_d - t_c\}^\alpha},$$

où  $s_i = i/n$  et  $t_j = j/m$  pour  $1 \leq i \leq n, 1 \leq j \leq m$ . Nous montrons que cette statistique est une fonctionnelle du processus de sommation  $\xi_n$ , continue sur l'espace de Hölder  $H_\alpha^o([0, 1]^2)$ . Ainsi, par image continue, le principe d'invariance hölderien du chapitre 3 nous permet d'établir la convergence :

$$\sigma^{-1}(nm)^{-1/2}DUI(n, m, \alpha) \xrightarrow{D} DUI(\alpha),$$

où

$$DUI(\alpha) = \sup_{\mathbf{0} \leq \mathbf{s} < \mathbf{t} \leq \mathbf{1}} \frac{|\Delta_{[\mathbf{s}, \mathbf{t}]}W - (t_1 - s_1)(t_2 - s_2)W(\mathbf{1})|}{|\mathbf{t} - \mathbf{s}|^\alpha},$$

en supposant que

$$\sup_{t>0} t^{1/(1/2-\alpha)} P(|X_{11}| > t) < \infty,$$

pour un certain  $\alpha \in ]0, 1/2[$  et en notant

$$\Delta_{[\mathbf{s}, \mathbf{t}]}x := x(\mathbf{t}) - x(s_1, t_2) - x(t_1, s_2) + x(\mathbf{s}),$$

pour toute fonction à deux variables  $x$ . La consistance du test est donnée par le théorème suivant.

**Théorème 6** *Sous  $(H_A)$ , supposons que les  $X_{ij}$  sont indépendantes et que  $\sigma_0^2 = \sup_n \text{var}(X_n)$  est fini. Supposons en outre que*

$$\lim_{m(n) \rightarrow \infty} (nm)^{1/2} \frac{h_{nm}}{d_{nm}^\alpha} |\mu_1 - \mu_0| \rightarrow \infty, \quad (8)$$

où en posant  $k^* = b^* - a^*$  et  $l^* = d^* - c^*$ ,

$$h_{nm} = \frac{k^* l^*}{nm} \left( 1 - \frac{k^* l^*}{nm} \right) \quad \text{et} \quad d_{nm} = \max \left\{ \frac{k^*}{n}, \frac{l^*}{m} \right\}.$$

Alors

$$(nm)^{-1/2}DUI(n, m, \alpha) \rightarrow \infty.$$

Ceci est une généralisation directe du résultat analogue de Račkauskas et Suquet [33] dans le cas  $d = 1$ . Cependant dans notre situation, la division par une puissance du diamètre du rectangle épidémique n'apporte pas un aussi grand avantage que la division par une puissance de la longueur de l'intervalle d'épidémie dans le cas  $d = 1$ . Pour le voir, supposons que  $k^* = n^\gamma$ ,  $l^* = m^\delta$  et que  $\mu_1 - \mu_0$  ne dépend pas de  $(n, m)$ . Alors la condition (8) devient

$$\frac{n^{\gamma-1/2} m^{\delta-1/2}}{[n^{\gamma-1} \vee m^{\delta-1}]^\alpha} \rightarrow \infty.$$

Si  $n^{\gamma-1} > m^{\delta-1}$ , alors

$$n^{\gamma(1-\alpha)+\alpha-1/2} m^{\delta-1/2} \rightarrow \infty,$$

ce qui permet la détection d'épidémies très courtes pour  $k^*$ , mais les plus courtes longueurs d'épidémie pour  $l^*$  ne peuvent être atteintes. Remarquons que pour  $\delta > 1/2$  la condition  $n^{\gamma-1} > m^{\delta-1}$  est vérifiée si  $m > n^2$ . Ainsi pour détecter des épidémies très courtes selon un indice, on a besoin de beaucoup plus de données épidémiques pour l'autre indice. Autrement dit, on n'arrive pas par cette méthode à détecter un rectangle épidémique dont les deux dimensions sont toutes deux très courtes. Il n'y a pas ce type de restriction pour le cas  $d = 1$  puisqu'alors  $h_n = l^*/n(1 - l^*/n)$  et  $d_n = l^*/n$ . Ainsi pour  $d = 1$  et une épidémie de longueur  $l^* = n^\gamma$ , la statistique de test diverge pour  $\gamma > (1/2 - \alpha)/(1 - \alpha)$ . Dans le cas  $d = 2$ , on arrive à un résultat similaire si on teste contre l'alternative

$(H'_A)$  : Il existe des entiers  $1 < c^* \leq d^* < m$  et une constante  $\mu_1 \neq \mu_0$  tels que

$$\mathbf{E} X_{ij} = \mu_0 + \mu_1 \mathbf{1} \left( (i, j) \in [1, n] \times [c^*, d^*] \cap \mathbb{N}^2 \right).$$

Pour tester contre cette alternative, nous proposons la statistique modifiée

$$DUI_3(n, m, \alpha) = \max_{1 \leq c < d \leq m} \frac{|S_{n,d} - S_{n,c} - (t_d - t_c)S_{n,m}|}{(t_d - t_c)^\alpha}.$$

Avec la même hypothèse de moment faible que pour  $DUI$ , on montre que sous l'hypothèse nulle, la loi limite de  $DUI_3(n, m, \alpha)$  est celle de

$$DUI_3(\alpha) = \sup_{0 < s < t < 1} \frac{|W(1, t) - W(1, s) - (t - s)W(1, 1)|}{|t - s|^\alpha}.$$

Les conditions pour la consistance sont alors données par le corollaire suivant.

**Corollaire 7** *La famille des  $X_{ij}$  étant indépendante et telle que  $\sup_n \text{var}(X_n) < \infty$ , sous l'hypothèse alternative  $H'_A$  nous avons*

$$(nm)^{-1/2} DUI_3(n, m, \alpha) \rightarrow \infty,$$

*pourvu que*

$$\lim_{m(n) \rightarrow \infty} (nm)^{1/2} \left( \frac{l^*}{m} \right)^{1-\alpha} \left( 1 - \frac{l^*}{m} \right) |\mu_1 - \mu_0| \rightarrow \infty.$$

Nous tournons ensuite notre attention vers les modèles de régression pour panels. On dispose d'un échantillon de données de panel  $\{(y_{ij}, \mathbf{x}'_{ij}), i = 1, \dots, n; j = 1, \dots, m\}$  où  $\mathbf{x}'_{ij} = (x_{ij1}, \dots, x_{ijK})$ . On considère les modèles de régression *pooled* (moindres carrés ordinaires) ou à effets fixes décrits par Baltagi [2] :

$$y_{ij} = \mathbf{x}'_{ij}\boldsymbol{\beta} + u_{ij}, \quad (9)$$

$$y_{ij} = \mathbf{x}'_{ij}\boldsymbol{\beta} + \mu_i + u_{ij}. \quad (10)$$

On propose des statistiques pour détecter un changement du coefficient de régression  $\boldsymbol{\beta}$ . L'approche utilisée est similaire à celle de Proberger et Krämer [23]. On prouve d'abord un FCLT pour les résidus de la régression, puis on utilise une statistique pour tester la rupture.

Les résidus de régression pour le modèle (9) sont définis comme

$$\hat{u}_{ij} = y_{ij} - \mathbf{x}'_{ij}\hat{\boldsymbol{\beta}} = u_{ij} - \mathbf{x}'_{ij}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}),$$

où

$$\hat{\boldsymbol{\beta}} = \left( \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}'_{ij} \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} y_{ij}.$$

est l'estimateur par moindres carrés du coefficient  $\boldsymbol{\beta}$ . On suppose que les perturbations  $u_{ij}$  vérifient un FCLT.

**Supposition F** On suppose en outre que le processus de sommation basé sur ces variables, défini par

$$\xi_{n,m}(t, s) = nm \sum_{i=1}^n \sum_{j=1}^m \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, s] \right| u_{ij},$$

vérifie le FCLT suivant :

$$\frac{1}{\sigma\sqrt{nm}} \xi_{n,m} \xrightarrow{D} W, \text{ quand } n \wedge m \rightarrow \infty,$$

dans l'espace  $H_\alpha^\alpha([0, 1]^2)$  pour un  $\alpha \in ]0, 1/2[$ .

On fait sur les  $\mathbf{x}_{ij}$  les hypothèses suivantes.

**Supposition A**  $1 \leq (i, j) \leq (n, m)$ . On suppose que

$$\lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}'_{ij} = R,$$

pour une matrice  $(K \times K)$  non singulière  $R$ . On suppose de plus que le modèle est reparamétrisé de sorte que

$$R = \begin{bmatrix} 1 & 0 \\ 0 & R^* \end{bmatrix},$$

ce qui à son tour, implique que

$$\mathbf{c} \equiv \lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} = [1, 0, \dots, 0]'$$

On prouve alors le théorème suivant.

**Théorème 8** *Sous les conditions F et A, pour le modèle de régression*

$$y_{ij} = \mathbf{x}_{ij}'\boldsymbol{\beta} + u_{ij},$$

définissons le processus de sommation

$$\widehat{W}^{(n,m)}(t, s) = nm \sum_{i=1}^n \sum_{j=1}^m \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, s] \right| \widehat{u}_{ij}.$$

Alors

$$\frac{1}{\sigma\sqrt{nm}} \widehat{W}^{(n,m)}(t, s) \xrightarrow{D} W(t, s) - tsW(1, 1), \text{ as } n \wedge m \rightarrow \infty,$$

dans l'espace  $H_\alpha^\alpha([0, 1]^2)$ , pour le  $\alpha \in ]0, 1/2[$  indiqué dans les conditions F.

Pour le modèle de régression à effets fixes (10), les résidus sont définis comme

$$\widehat{u}^{FE} = \widetilde{y}_{ij} - \widetilde{\mathbf{x}}_{ij}'\widehat{\boldsymbol{\beta}}^{FE},$$

où  $\widetilde{y}_{ij} = y_{ij} - \bar{y}_i$  avec  $\widetilde{\mathbf{x}}_{ij}$  défini de manière analogue et  $\widehat{\boldsymbol{\beta}}^{FE}$ , l'estimateur classique dans le cas des effets fixes du coefficient de régression  $\boldsymbol{\beta}$ , est donné par

$$\widehat{\boldsymbol{\beta}}^{FE} = \left( \sum_{i=1}^n \sum_{j=1}^m \widetilde{\mathbf{x}}_{ij} \widetilde{\mathbf{x}}_{ij}' \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \widetilde{\mathbf{x}}_{ij} \widetilde{y}_{ij}.$$

Introduisons les hypothèses suivantes.

**Supposition B** *On suppose que*

$$\lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \tilde{\mathbf{x}}'_{ij} = \tilde{R}$$

*pour une matrice  $(K \times K)$  non singulière  $R$ , et que*

$$\lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} = \mathbf{c}$$

*pour un  $\mathbf{c} \in \mathbb{R}^K$ .*

On prouve alors le théorème suivant.

**Théorème 9** *pour le modèle de régression de panel*

$$y_{ij} = \mathbf{x}'_{ij} \boldsymbol{\beta} + \mu_i + u_{ij},$$

*le processus de sommation*

$$\widehat{W}_{nm}^{FE}(t, s) = nm \sum_{i=1}^n \sum_{j=1}^m \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, s] \right| \widehat{u}_{ij}^{FE}.$$

*Alors*

$$\widehat{W}_{nm}^{FE}(t, s) \xrightarrow{D} W(t, s) - sW(t, 1), \text{ quand } n \wedge m \rightarrow \infty,$$

*dans l'espace  $H_\alpha^o([0, 1]^2)$ , pour le  $\alpha \in ]0, 1/2[$  de la condition F.*

Pour les deux modèles, on étudie des alternatives locales de la forme

$$\boldsymbol{\beta}_{ij} = \boldsymbol{\beta} + \frac{1}{\sqrt{nm}} \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right)$$

avec  $\mathbf{g} : [0, 1]^2 \rightarrow \mathbb{R}^K$  continue. Sous les conditions F et A (resp. B), on obtient un FCLT dans  $H_\alpha^o([0, 1]^2)$  pour les résidus associés aux alternatives locales. Le processus limite pour le modèle (9) est alors

$$W(t, s) - tsW(1, 1) + \int_0^t \int_0^s \mathbf{c}' \mathbf{g}(u, v) du dv - ts \mathbf{c}' \int_0^1 \int_0^1 \mathbf{g}(u, v) du dv$$

et pour le modèle (10)

$$W(t, s) - sW(t, 1) + \int_0^t \int_0^s \mathbf{c}' \mathbf{g}(u, v) du dv - s \int_0^t \int_0^1 \mathbf{c}' \mathbf{g}(u, v) du dv.$$

Pour tester une rupture dans le coefficient de régression, nous proposons d'utiliser la statistique  $DUI(n, m, \alpha)$ , où les sommes sont celles des résidus de régression. Nous testons l'hypothèse nulle

$(H_0)$  : *Le coefficient  $\beta$  ne change pas,*  
contre l'alternative

$(H_A)$  : *Il existe des entiers  $1 < a^* \leq b^* < n$ ,  $1 < c^* \leq d^* < m$  et des constantes  $\beta_1 \neq \beta_0$  tels que le vrai coefficient  $\beta$  admette la représentation*

$$\beta = \beta_0 + \beta_1 \mathbf{1} \left( (i, j) \in [a^*, b^*] \times [c^*, d^*] \cap \mathbb{N}^2 \right).$$

On donne la loi asymptotique de la statistique de test sous l'hypothèse nulle pour les deux modèles de régression. Pour la régression ordinaire par moindres carrés, celle ci demeure la même :

$$\sigma^{-1}(nm)^{-1/2} \widehat{DUI}(n, m, \alpha) \xrightarrow{D} DUI(\alpha), \text{ quand } n \wedge m \rightarrow \infty,$$

Pour la régression à effets fixes, elle se change en celle de

$$DUI^{FE}(\alpha) = \sup_{\mathbf{0} \leq \mathbf{s} < \mathbf{t} \leq \mathbf{1}} \frac{|\Delta_{[\mathbf{s}, \mathbf{t}]} W - (t_2 - s_2)[W(t_1, 1) - W(s_1, 1)]|}{|\mathbf{t} - \mathbf{s}|^\alpha}.$$

On donne aussi des conditions pour la consistance des deux tests et les lois limites des statistiques sous les alternatives locales.

Le chapitre se termine par une discussion des résultats. On y note que dans les applications, l'hypothèse i.i.d. est parfois trop restrictive, mais que nos résultats sont prouvés d'une manière « portable » en ce sens que si l'on dispose de nouveaux principes d'invariance pour une classe plus large de variables aléatoires à double indice (par exemple vérifiant certaines conditions de dépendance faible), les statistiques de test et leurs lois limites restent les mêmes, aussi longtemps que le processus limite dans ces nouveaux principes d'invariance est le drapeau de Wiener.

# Disertacijos santrauka

## Invariantiškumo principas daugiamačio indekso sumavimo procesams ir taikymai

Invariantiškumo principas tai tikimybių teorijos rezultatas teigiantis, kad tam tikras atsitiktinių dydžių dalinių sumų procesas silpnai konverguoja ir kad ribinis procesas nepriklauso nuo sumuojamų atsitiktinių dydžių pasiskirstymo. Šis rezultatas yra centrinės ribinės teoremos (CRT) apibendrinimas, kartais dar taip pat vadinamas funkcinė centrine ribine teorema (FCRT).

Pirmus tokio tipo rezultatus gavo Prokhorov [24] ir Donsker [10]. Jie nagrinėjo sumavimo procesą su nuliniu vidurkiu nepriklausomais ir vienodai pasiskirsčiusiais n.v.p. atsitiktiniais dydžiais  $X_1, \dots, X_n$ :

$$\xi_n(t) = S_{[nt]} + (nt - [nt])X_{[nt]+1}, t \in [0, 1], \quad (11)$$

čia  $S_k = X_1 + \dots + X_k$ ,  $k = 1, \dots, n$  ir  $S_0 = 0$ . Šio proceso trajektorijos yra laužtės, taigi procesas gali būti nagrinėjamas kaip atsitiktinis elementas su reikšmėmis tolydžių funkcijų erdvėje  $C([0, 1])$ . Donsker ir Prokhorov įrodė, kad toks procesas (padalintas iš  $\sqrt{n}$  ir  $\sigma = \sqrt{\mathbf{E} X_1^2}$ ) silpnai konverguoja į atsitiktinį Brauno judesį  $W(t)$  – Gauso procesą su nuliniu vidurkiu ir kovariacijos funkcija  $\mathbf{E} W(t)W(s) = t \wedge s$ . Būtina ir pakankama sąlyga konvergavimui yra baigtinis antras momentas, t.y.  $\sigma < \infty$ . (Laikoma, kad atsitiktinių elementų seka  $Y_n$  iš separabilios metrinės erdvės  $B$  silpnai konverguoja į atsitiktinį elementą  $Y$ , jeigu kiekvienam tolydžiam aprėžtam funkcionalui  $f$ ,  $\mathbf{E} f(Y_n) \rightarrow \mathbf{E} f(Y)$ ). Kadangi  $\xi_n(1) = S_n$  iš karto matome, kodėl šis rezultatas yra CRT apibendrinimas.

Iš FCRT bei tolydaus atvaizdžio teoremos išplaukia, kad kiekvienam tolydžiam funkcionalui  $g$ ,  $g(\sigma^{-1}n^{-1/2}\xi_n)$  silpnai konverguoja į  $g(W)$ . Šis rezultatas yra plačiai taikomas statistikoje. Klasikinis tokio pritaikymo pavyzdys yra statistinių testų skirtų nustatyti imties vidurkio struktūrinį pasikeitimą

konstravimas. Vieno iš tokių testų statistika yra

$$Q = \max_{1 \leq k \leq n} |S_k - \frac{k}{n} S_n|.$$

Galima įsitikinti, kad  $Q = g(\xi_n)$ , funkcionalui  $g(x) = \sup_{0 \leq t \leq 1} |x(t)|$ . Šis funkcionalas yra tolydus erdvėje  $C([0, 1])$ , taigi iš FCRT ir tolydaus atvaizdžio teoremos, prie nulinės hipotezės, kad imties  $\{X_1, \dots, X_n\}$  elementų vidurkis nesikeičia, iš karto išplaukia, kad

$$n^{-1/2}Q \xrightarrow{D} \sup_{0 < t < 1} |W(t) - tW(1)|.$$

Prie alternatyvios hipotezės, tarus, kad egzistuoja  $k^*$ , toks, kad  $\mathbf{E} X_i = \mu_0 + (\mu_1 - \mu_0)\mathbf{1}(i > k^*)$ , galima įrodyti, kad statistika  $n^{-1/2}Q$  diverguoja. Taigi testas yra suderintas. Nagrinėjant taip vadinamą epideminę alternatyvą  $\mathbf{E} X_i = \mu_0 + (\mu_1 - \mu_0)\mathbf{1}(k^* < i \leq l^*)$ , paprastai naudojama statistika

$$R = \max_{1 \leq k \leq n} \frac{|S_l - S_k - \frac{l-k}{n} S_n|}{[(l-k)/n]^\alpha}.$$

Funkcionalas atitinkantis šią statistiką

$$g(x) = \sup_{0 < |t-s| < 1} \frac{|x(t) - x(s)|}{|t-s|^\alpha}$$

jau nėra tolydus erdvėje  $C([0, 1])$ . Jis yra tolydus Banacho erdvėje  $(H_\alpha^o, \|\cdot\|_\alpha)$ ,  $0 < \alpha < 1$ , kuriai priklauso funkcijos  $x : [0, 1] \rightarrow \mathbb{R}$  tenkinančios

$$\|x\|_\alpha := |x(0)| + w_\alpha(x, 1) < \infty,$$

ir

$$w_\alpha(x, \delta) := \sup_{0 < |t-s| \leq \delta} \frac{|x(t) - x(s)|}{|t-s|^\alpha} \xrightarrow{\delta \rightarrow 0} 0.$$

Taigi norint rasti ribinį statistikos  $R$  pasiskirstymą, reikia turėti FCRT erdvėje  $H_\alpha^o$ . Reikėtų pastebėti, kad erdvės  $H_\alpha^o$  topologija yra stipresnė nei erdvės  $C([0, 1])$ , taigi jos funkcionalų erdvė yra turtingesnė. Taip pat reikėtų pastebėti, kad dėl Brauno judesio tolydumo modulio, atsiranda apribojimas  $0 < \alpha < 1/2$ . Statistika  $R$  yra naudinga tuo, kad leidžia aptikti trumpas epidemijas, kai  $k^* - l^*$  yra  $n^\gamma$  eilės, su  $0 < \gamma < 1/2$ . Rodiklio  $\gamma$  pasirinkimas priklauso nuo  $X_i$  integruojamumo. Be vardiklio statistika  $R$  leistų aptikti epidemijas ne trumpesnes kaip  $n^{1/2}$ . Vienas iš disertacinio darbo tikslų buvo pabandyti perkelti šį pranašumą daugiamačio indekso sumavimo procesams.

Daugiamačio indekso sumavimo procesai sutinkami studijuojant atsitiktinių dydžių su daugiamačiu indeksu dalines sumas. Be matematinio smalsumo tokių dydžių tyrimas turi ir praktinę vertę, nes kartais keletas indeksų vienam stebėjimui priskiriami natūraliai. Pavyzdžiui jeigu mes matuojame keleto individų kokią nors savybę tam tikrais laiko tarpais, yra natūralu kiekvienam stebėjimui priskirti du indeksus, vieną pažymėti individui, kitą pažymėti laikui. Tokio tipo duomenys yra vadinami paneliniais bei jų tyrimui pastaruoju metu skiriama daug dėmesio.

Norėdami pademonstruoti mūsų gautų rezultatų kontekstą pateiksime bendresnį nei darbe naudojamą daugiamačio indekso sumavimo proceso apibrėžimą. Tegų  $|A|$  žymi Borelio aibės  $A \subset \mathbb{R}^d$  Lebego matą. Sumavimo procesas  $\{\xi_n(A); A \in \mathcal{A}\}$  aibių klasei  $\mathcal{A}$  nepriklausomai vienodai pasiskirsčiusių realių atsitiktinių dydžių su nuliniu vidurkiu atveju yra apibrėžiamas taip:

$$\xi_n(A) = \sum_{1 \leq j \leq n} |R_{n,j}|^{-1} |R_{n,j} \cap A| X_j, \quad (12)$$

čia  $\mathbf{j} = (j_1, \dots, j_d)$ ,  $\mathbf{n} = (n_1, \dots, n_d)$ ,  $R_{n,j}$  yra stačiakampis

$$R_{n,j} := \left[ \frac{j_1 - 1}{n_1}, \frac{j_1}{n_1} \right) \times \dots \times \left[ \frac{j_d - 1}{n_d}, \frac{j_d}{n_d} \right)$$

ir užrašas “ $1 \leq \mathbf{j} \leq \mathbf{n}$ ” suprantamas pakoordinačiui:  $1 \leq j_1 \leq n_1, \dots, 1 \leq j_d \leq n_d$ . Pastebėkime, kad aibių klasei  $\mathcal{A} = \{[0, t], t \in [0, 1]\}$  taip apibrėžtas procesas sutampa su procesu apibrėžtu (11).

Aibių klasėje  $\mathcal{A}$  apibrėžus semimetriką  $\delta$ , galima nagrinėti realių tolydžių funkcijų erdvę  $C(\mathcal{A})$  su norma

$$\|f\|_{\mathcal{A}} := \sup_{A \in \mathcal{A}} |f(A)|.$$

Įprastinės semimetrikos yra  $\delta(A, B) = \sqrt{|A \Delta B|}$ , arba  $\delta(A, B) = \sqrt{m(A \Delta B)}$ , čia  $A, B \in \mathcal{A}$ , ir  $m$  yra tikimybinis matas apibrėžtas vienetinio  $d$ -mačio kubo  $[0, 1]^d$  Borelio poaibiams. Tokiu atveju, proceso  $\xi_n(A)$  trajektorijos priklauso erdvei  $C(\mathcal{A})$ . Jei  $\mathcal{A}$  yra pilnai aprėžta semimetrikos  $\delta$  atžvilgiu, tai  $C(\mathcal{A})$  yra separabili Banacho erdvė. Kadangi tokiu atveju procesas  $\{\xi_n(A), A \in \mathcal{A}\}$  yra atsitiktinis elementas su reikšmėmis separabilioje metrinėje erdvėje, tai šiam procesui galime tirti FCRT. Ribinis procesas tada yra Brauno judesio apibendrinimas, vadinamas Vynerio arba Brauno paklode. Tai nulinio vidurkio Gauso procesas  $W$  indeksuotas  $\mathcal{A}$  su trajektorijomis erdvėje  $C(\mathcal{A})$ , bei

$$\mathbf{E} W(A)W(B) = |A \cap B|, \quad A, B \in \mathcal{A}.$$

Tokie procesai egzistuoja, jei aibių klasei  $\mathcal{A}$  įvedami apribojimai, dažniausiai išreiškiami entropijos sąlyga.

Pirmoji FCRT procesui  $\{\xi_n(A); A \in \mathcal{A}\}$  erdvėje  $C(\mathcal{A})$  buvo įrodyta aibių klasei  $\mathcal{A} = \mathcal{Q}_d$ , kai

$$\mathcal{Q}_d := \left\{ [0, t_1] \times \cdots \times [0, t_d]; \mathbf{t} = (t_1, \dots, t_d) \in [0, 1]^d \right\}. \quad (13)$$

Ją įrodė Kuelbs [19], prie tam tikrų momentinių sąlygų bei Wichura [42] baigtinės dispersijos atveju.

Platesnėms aibių klasėms  $\mathcal{A}$  tenkinančioms tam tikrą entropijos sąlygą, Pyke [25] įrodė FCRT erdvėje  $C(\mathcal{A})$ . Jo rezultate sąlygos sumuojamiems atsitiktiniams dydžiams priklausė nuo  $\mathcal{A}$  savybių. Bass [3] ir tuo pačiu metu Alexander and Pyke [1] pagerino Pyke rezultatą n.v.p. atsitiktiniams laukams turintiems baigtinę dispersiją. Jie įrodė kad FCRT galioja erdvėje  $C(\mathcal{A})$ , kai sumuojami atsitiktiniai dydžiai yra n.v.p. ir turi baigtinę dispersiją ir kai aibių klasė  $\mathcal{A}$  tenkina tam tikrą entropijos sąlygą.

Vėlesni autoriai nagrinėjo galimybes susilpninti entropijos sąlygas aibių klasei  $\mathcal{A}$ , Ziegler [43], bei n.v.p. reikalavimą atsitiktiniam laukui  $\{X_n, n \in \mathbb{N}^d\}$ , Dedeker [8], El Machkouri and Ouchti [12] ir kiti.

FCRT Hiolderio erdvėse buvo skirtas mažesnis dėmesys. 1962 metais Lamperti [20] nagrinėjo FCRT erdvėje  $H_\alpha^o$ ,  $0 < \alpha < 1/2$ . Laikydamas, kad  $\mathbf{E} |X_1|^q < \infty$ , kai  $q > 2$ , jis parodė, kad procesas  $n^{-1/2}\xi_n$  silpnai konverguoja į  $W$  Hiolderio erdvėje  $H_\alpha^o$  bet kokiam  $\alpha < 1/2 - 1/q$ . Račkauskas ir Suquet [31] (taip pat [28]) gavo būtinas ir pakankamas sąlygas Lamperti FCRT. Jie parodė, kad  $n^{-1/2}\xi_n$  konverguoja į  $W$  erdvėje  $H_\alpha^o$ ,  $0 < \alpha < 1/2$ , tada ir tik tada, kai

$$\lim_{t \rightarrow \infty} t^{p(\alpha)} P(|X_1| > t) = 0, \quad (14)$$

čia

$$p(\alpha) := \frac{1}{\frac{1}{2} - \alpha}. \quad (15)$$

Apibendrinta Hiolderio erdvė  $H_\rho(\mathcal{A})$  aibių klasei  $\mathcal{A}$  ir svorio funkcijai  $\rho$  yra apibrėžiama, kaip erdvės  $C(\mathcal{A})$  poaibis, kuriam priklauso funkcijos tenkinančios sąlygą

$$\sup_{0 < \delta(A, B) < 1} \frac{|x(A) - x(B)|}{\rho(\delta(A, B))} < \infty.$$

Sąlygas  $W$  egzistavimui tokiose erdvėse pateikia Dudley [11] ir Erickson [13]. Kai  $\rho(h) = h^\alpha$ , Erickson [13] parodo, kad parametras  $\alpha$  negali viršyti  $1/2$  bei priklauso nuo aibių klasės  $\mathcal{A}$  entropijos. Erickson [13] parodo, kad bendriausia FCRT erdvėje  $H_\rho(\mathcal{A})$  galioja, jei  $\mathbf{E} |X_j|^q < \infty$  tam tikram  $q > 2$  bei  $\rho$ ,

kuris priklauso nuo  $q$  bei  $\mathcal{A}$  savybių. Vienmačiu atveju  $d = 1$ , aibių klasei  $\mathcal{A}$  sutampant su intervalų  $[0, t]$ ,  $0 \leq t \leq 1$  aibe, Erickson rezultatas sutampa su Lamperti. Daugiamačiu atveju  $d > 1$  Erickson parodo, kad atsitiktiniai dydžiai privalo turėti ne mažesnę kaip  $q > dp(\alpha)$  momentą, su  $p(\alpha)$  apibrėžtu (15).

Aibių klasė  $\mathcal{Q}_d$  yra izomorfiška vienetiniam kubui  $[0, 1]^d$  dėl bijekcijos  $[0, t] \leftrightarrow \mathbf{t}$ . Taigi natūralu nagrinėti procesą  $\xi_{\mathbf{n}}(A)$  kaip keleto kintamųjų funkciją  $\xi_{\mathbf{n}}(\mathbf{t})$ . Žinant tai, kad Račkauskas ir Suquet yra išsamiai išstudijavę keleto kintamųjų funkcijų Hiolderio erdves ir pateikę visus FCRT įrodyti reikalingus rezultatus, bei tai, kad Lamperti invariantiškumo principą vienmačiu atveju galima įrodyti būtinas ir pakankamas sąlygas, kyla klausimas, ar įmanoma gauti būtinas ir pakankamas sąlygas daugiamačio indekso atveju. Į šią klausimą disertacijoje pilnai atsakoma.

Pirmame disertacijos skyriuje yra surinkti visi reikalingi rezultatai FCRT įrodinėti Hiolderio erdvėse. Kadangi sumavimo procesas  $\{\xi_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}^d\}$  yra apibendrinta seka, o visi įprastiniai rezultatai yra sekoms, skyriuje jie yra adaptuojami. Skyriuje Hiolderio erdvė  $H_{\alpha}^o(\mathbb{H})$ , kuriai priklauso daugiamačio argumento funkcijos su reikšmėmis Hilberto erdvėje  $H$  apibrėžiama kaip vektorinė funkcijų  $x : [0, 1]^d \rightarrow \mathbb{H}$  tenkinančių

$$\|x\|_{\alpha} := \|x(0)\| + \omega_{\alpha}(x, 1) < \infty,$$

ir

$$\omega_{\alpha}(x, \delta) := \sup_{0 < |\mathbf{t} - \mathbf{s}| \leq \delta} \frac{\|x(\mathbf{t}) - x(\mathbf{s})\|}{|\mathbf{t} - \mathbf{s}|^{\alpha}} \xrightarrow{\delta \rightarrow 0} 0$$

erdvė. Su norma  $\|\cdot\|_{\alpha}$ ,  $H_{\alpha}^o(\mathbb{H})$  yra separabili Banacho erdvė, žr. [27] arba [33]. Toliau skyriuje yra pateikiami naudingi faktai apie šią erdvę iš Račkausko ir Suquet [27]. Remiantis šiais faktais yra įrodomas tirštumo kriterijus apibendrintoms erdvės  $H_{\alpha}^o(\mathbb{H})$  sekoms.

**Teorema 1** *Tegu  $\{\zeta_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}^d\}$  ir  $\zeta$  yra atsitiktiniai elementai su reikšmėmis erdvėje  $H_{\alpha}^o(\mathbb{H})$ . Jei*

i) *Kiekvienam diadiniam  $\mathbf{t} \in [0, 1]^d$ , apibendrinta atsitiktinių elementų su reikšmėmis  $\mathbb{H}$  seka  $\zeta_{\mathbf{n}}(\mathbf{t})$  yra asimptotiškai tiršta  $\mathbb{H}$ .*

ii) *Kiekvienam  $\varepsilon > 0$*

$$\lim_{J \rightarrow \infty} \limsup_{m(\mathbf{n}) \rightarrow \infty} P\left(\sup_{j \geq J} 2^{\alpha j} \max_{v \in V_j} \|\lambda_{j,v}(\zeta_{\mathbf{n}})\| > \varepsilon\right) = 0,$$

*tai apibendrinta seka  $\zeta_{\mathbf{n}}$  yra asimptotiškai tiršta erdvėje  $H_{\alpha}^o(\mathbb{H})$ .*

Naudojantis šia išraiška apibrėžus atsitiktinį lauką  $\xi_n^*$ :

$$\xi_n^*(\mathbf{t}) = \sum_{\mathbf{u} \in \{0,1\}^d} w(\mathbf{u}) S_{[\mathbf{n}\mathbf{t}]+\mathbf{u}}, \quad \mathbf{t} \in [0, 1]^d,$$

jis sutampa su sumavimo procesu apibrėžtu (12), kai  $\{X_i, \mathbf{1} \leq \mathbf{i} \leq \mathbf{n}\}$  yra aibė atsitiktinių dydžių su reikšmėmis Banacho erdvėje.

Šio teiginio pagalba mes galime įvertinti išraišką panaudotą tirštumo kriterijuje 1:

$$\sup_{j \geq J} 2^{\alpha j} \max_{\mathbf{v} \in V_j} \|\lambda_{j,\mathbf{v}}(\xi_n)\| \leq 3^d \sum_{m=1}^d \left( \max_{J \leq j \leq \log n_m} 2^{\alpha j} \max_{0 \leq k < 2^j} \psi_n^{(m)}(t_{k+1}, t_k) + n_m^\alpha Z_n^{(m)} \right)$$

čia

$$\psi_n^{(m)}(t', t) := \max_{\mathbf{1}-m \leq \mathbf{k}-m \leq \mathbf{n}-m} \left\| \sum_{k_m=[n_m t]+1}^{[n_m t']} \Delta_{k_m}^{(m)} S_{\mathbf{k}} \right\|,$$

$$Z_n^{(m)} := \max_{1 \leq \mathbf{k} \leq \mathbf{n}} \|\Delta_{k_1}^{(m)} S_{\mathbf{k}}\|.$$

Šiose išraiškose naudojami skirtuminiai operatoriai  $\Delta$  yra apibrėžiami taip:

$$\Delta_k^{(i)} S_j := S_{(j_1, \dots, j_{i-1}, k, j_{i+1}, \dots, j_d)} - S_{(j_1, \dots, j_{i-1}, k-1, j_{i+1}, \dots, j_d)}.$$

Gautas įvertis nereikalauja jokių prielaidų atsitiktinių dydžių pasiskirstymams, taigi jis gali būti naudojamas įrodinėjant FCRT ne tik n.v.p. a. d. atveju.

Trečiame skyriuje yra įrodomas invariantiškumo principas erdvėje  $H_\alpha^o(\mathbb{H})$ .

**Teorema 3** Tegu  $0 < \alpha < 1/2$  ir  $p = p(\alpha) := 1/(1/2 - \alpha)$ . Tarkime, kad kiekvienam  $d \geq 2$ ,  $\{X_i; \mathbf{i} \in \mathbb{N}^d, \mathbf{i} \geq \mathbf{1}\}$  yra apibendrinta n.v.p. atsitiktinių dydžių su reikšmėmis separabilioje Hilberto erdvėje  $\mathbb{H}$  seka, kurių vidurkis yra nulis ir kuriems egzistuoja antras momentas. Tegu  $\xi_n$  yra sumavimo procesas apibrėžtas

$$\xi_n(\mathbf{t}) = \sum_{\mathbf{i} \leq \mathbf{n}} \pi(\mathbf{n}) \left| \left[ \frac{\mathbf{i} - \mathbf{1}}{\mathbf{n}}, \frac{\mathbf{i}}{\mathbf{n}} \right) \cap [0, \mathbf{t}] \right| X_i,$$

o  $W$  yra Vynerio paklodė su reikšmėmis  $\mathbb{H}$  ir tokiu pačiu kovariacijos operatoriumi kaip  $X_1$ . Tada

$$\pi(\mathbf{n})^{-1/2} \xi_n \xrightarrow[\mathfrak{m}(\mathbf{n}) \rightarrow \infty]{H_\alpha^o(\mathbb{H})} W$$

tada ir tik tada, kai

$$\pi(\mathbf{n})P(\|X_1\| > n_m^{1/p}\pi(\mathbf{n}_{-m})^{1/2}) \xrightarrow{m(\mathbf{n}) \rightarrow \infty} 0. \quad (18)$$

kiekvienam  $m = 1, \dots, d$ . Čia  $\pi(\mathbf{n})$  žymi indekso  $\mathbf{n}$  koordinatų sandaugą, o  $\pi(\mathbf{n}_{-m})$  sandaugą visų koordinatų išskyrus  $n_m$ .

Nors sąlyga (18) iš pirmo žvilgsnio atrodo pakankamai sudėtinga, yra įrodoma, kad ji yra ekvivalenti atsitiktinio dydžio  $X_1$  silpno  $p$ -momento egzistavimui t.y.:

$$\sup_{t>0} t^p P(\|X_1\| > t) < \infty.$$

Iš čia iš karto išplaukia, kad darbe pavyko pagerinti Erickson rezultata, nes jo darbe FCRT reikalaujama, kad  $\mathbf{E}|X_1|^q < \infty$ , kai  $q > dp$ .

Taip pat sąlyga (18) yra naudinga tuo, kad ji parodo, kokios momentinės sąlygos turi būti tenkinamos, jeigu nagrinėjamas silpnėsnis konvergavimas, nei  $m(\mathbf{n}) \rightarrow \infty$ . Imkime atskirą atvejį, kai  $\mathbf{n} = (n, \dots, n)$ . Pastebėsime, kad literatūroje tai dažnai paplitusi prielaida. Tada būtina ir pakankama sąlyga FCRT tampa

$$\lim_{t \rightarrow \infty} t^{\frac{2d}{d-2\alpha}} P(\|X_1\| > t) = 0.$$

Ji yra tenkinama visiems  $d > 1$  ir  $0 < \alpha < 1/2$ , jeigu  $\mathbf{E}\|X_1\|^4 < \infty$ . Ši sąlyga parodo, kad perėjus prie daugiamatės indeksų situacija pasikeičia, nes tam kad FCRT galėtų įprastam Donsker-Prokhorov laužčių procesui yra būtina, kad  $\mathbf{E}|X_1|^q < \infty$  bet kokiam  $q < p(\alpha)$ .

Antrame ir trečiame skyriuose tiriant galimybę susilpninti n.v.p. sąlygą buvo nagrinėjamas serijų schemas atvejais. Daugiamatės indeksų sumavimo procesams serijų schemas atvejais taip pat buvo tiriamas Goldie ir Greenwood [14], [15]. Jie naudojo klasikinę sumavimo proceso konstrukciją (12), taigi jų rezultatas atveju  $d = 1$  nesutapo su Prokhorov [24] rezultatu *adaptyviam* laužčių procesui  $\Xi_n$  su viršūnėmis  $(b_n(k), S_n(k))$ , čia  $b_n(k) = \mathbf{E}X_{n,1}^2 + \dots + \mathbf{E}X_{n,k}^2$ , variant, kad  $b_n(k_n) = 1$ , ir kad  $X_{n,k}$  – nepriklausomi, bet nevienodai pasiskirstę atsitiktiniai dydžiai.

Pirmieji adaptyvią konstrukciją daugiamatės indeksų sumavimo procesams pasiūlė Bickel ir Wichura [5]. Tačiau jų darbe yra reikalavimas, kad atsitiktinių dydžių iš serijų schemas dispersijos tenkintų tam tikras sąlygas. Nulinio vidurkio nepriklausomiems realiems atsitiktiniams dydžiams  $\{X_{n,ij}, 1 \leq i \leq I_n, 1 \leq j \leq J_n\}$  su dispersijomis  $\mathbf{E}X_{n,ij} = a_{n,i}b_{n,j}$ , čia  $\sum a_{n,i} = 1 = \sum b_{n,j}$ , jų pasiūlyta konstrukcija buvo

$$\zeta_n(t_1, t_2) = \sum_{i \leq A_n(t_1)} \sum_{j \leq B_n(t_1)} X_{n,ij},$$

Čia  $m(\mathbf{n})$  žymi vektoriaus  $\mathbf{n}$  koordinačių minimumą, o išraiška  $\lambda_{j,v}(\zeta_{\mathbf{n}})$  yra paimta iš tam tikros erdvės  $H_{\alpha}^{\circ}(\mathbb{H})$  Šauderio dekompozicijos bei apibrėžiama kaip antros eilės  $\zeta_{\mathbf{n}}$  skirtumas tam tikruose diadiniuose taškuose.

Toliau skyriuje apibrėžiamas FCRT erdvėje  $H_{\alpha}^{\circ}(\mathbb{H})$  ribinis procesas. Tai Vynerio paklodė su reikšmėmis erdvėje  $\mathbb{H}$  ir kovariacijos operatoriumi  $\Gamma$ . Šis procesas apibrėžiamas kaip Gauso procesas su reikšmėmis erdvėje  $\mathbb{H}$  ir argumentų aibe  $[0, 1]^d$ , tenkinantis sąlygą

$$\mathbf{E} \langle W(\mathbf{t}), x \rangle \langle W(\mathbf{s}), y \rangle = (t_1 \wedge s_1) \dots (t_d \wedge s_d) \langle \Gamma x, y \rangle$$

čia  $\mathbf{t}, \mathbf{s} \in [0, 1]^d$  ir  $x, y \in \mathbb{H}$ . Kadangi toks procesas yra retai sutinkamas literatūroje yra pateikiamas jo egzistavimo įrodymas.

Skyriaus gale yra primenamos Rosenthal ir Doob nelygybės, kadangi darbe jos plačiai naudojamos tirštumui įrodyti. Atsitiktiniams elementams  $(Y_i)_{i \in I}$  su reikšmėmis Hilberto erdvėje, Rosenthal nelygybė yra pateikiama pavidalu

$$\mathbf{E} \left\| \sum_{i \in I} Y_i \right\|^q \leq C_q' \left( \mathbf{E} \left\| \sum_{i \in I} G(Y_i) \right\|^q + \sum_{i \in I} \mathbf{E} \|Y_i\|^q \right),$$

čia  $I$  yra indeksų aibė, kuriai nekeliame jokie reikalavimai. Daugiamatnio indekso realioms dalinėms sumoms Doob nelygybės apibendrinimą yra pateikęs Khoshnevisan[18]. Skyriuje parodoma, kad nelygybė galioja ir daugiamatnio indekso dalinėms sumoms su reikšmėmis Hilberto erdvėse:

$$\mathbf{E} \max_{0 \leq j \leq \mathbf{n}} \|S_j\|^p \leq \left( \frac{p}{p-1} \right)^{dp} \mathbf{E} \|S_{\mathbf{n}}\|^p,$$

visiems  $\mathbf{n} \in \mathbb{N}^d$  ir  $p > 1$ .

Antrame skyriuje yra detalios išnagrinėtos sumavimo proceso (12) savybės. Pasirodo, kad šis procesas gali būti išreikštas kitokia, patogesne forma.

**Teiginys 2** Tegu  $\mathbf{t} \in [0, 1]^d$ . Pažymėkime  $\mathbf{s} = \{\mathbf{n}\mathbf{t}\}$  ir stačiakampio  $R_{\mathbf{n}, [\mathbf{n}\mathbf{t}] + 1}$  viršūnes  $V(\mathbf{u})$ :

$$V(\mathbf{u}) := \frac{[\mathbf{n}\mathbf{t}]}{\mathbf{n}} + \frac{\mathbf{u}}{\mathbf{n}}, \quad \mathbf{u} \in \{0, 1\}^d. \quad (16)$$

Tada yra įmanoma išreikšti  $\mathbf{t}$  šitų viršūnių svorine suma, su svoriais priklausančiais nuo  $\mathbf{t}$ :

$$\mathbf{t} = \sum_{\mathbf{u} \in \{0, 1\}^d} w(\mathbf{u}) V(\mathbf{u}), \quad \text{where} \quad \sum_{\mathbf{u} \in \{0, 1\}^d} w(\mathbf{u}) = 1, \quad (17)$$

čia

$$w(\mathbf{u}) = \prod_{l=1}^d s_l^{u_l} (1 - s_l)^{1-u_l}.$$

čia

$$A_n(t_1) = \max\{k : \sum_{i \leq k} a_{n,i} < t_1\},$$

$$B_n(t_2) = \max\{l : \sum_{j \leq l} b_{n,j} < t_2\}.$$

Lengva pastebėti, kad ši konstrukcija yra Prokhorov konstrukcijos apibendrinimas dvimačio indekso atveju. Bickel ir Wichura įrodė, kad procesas  $\zeta_n$  konverguoja į Vynerio paklodę dvimačio argumento càdlàg funkcijų erdvėje  $D([0, 1]^2)$ , jei  $a_{n,i}$  ir  $b_{n,j}$  tenkina begalinio mažumo, o atsitiktiniai dydžiai  $\{X_{n,ij}\}$  Lindebergo sąlygas.

Antrame skyriuje yra pasiūlomas naujas Prokhorov konstrukcijos apibendrinimas daugiamatnio indekso sumavimo procesams. Disertaciniame darbe įrodoma, kad nors Bickel ir Wichura yra atskiras pasiūlytos konstrukcijos atvejis, ši konstrukcija vis tiek netinkama bet kokiai serijų schemai.

Prokhorov konstrukcijoje dalinės sumos yra išdėliojamos ne ant tolygiai išsidėsčiusių taškų, bet ant taškų kurie yra dalinių sumų dispersijos. Daugiamatnio indekso sumavimo proceso įprastinė konstrukcija dalines sumas išdėlioja ant stačiakampės gardelės. Deja bendru atveju nėra įmanoma sukonstruoti tokios stačiakampės gardelės, kurios taškai būtų dalinių sumų dispersijos. Darbe yra pasiūloma naudoti gardelę, kurios koordinatės ant kiekvienos ašies yra dispersijų sumos pagal visas likusias ašis. Aiškumo dėlei serijų schemai

$$(X_{n,k}, \mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n), \mathbf{n} \in \mathbb{N}^d,$$

pažymėkime

$$b_n(\mathbf{k}) := \sum_{j \leq \mathbf{k}} \sigma_{n,\mathbf{k}}^2.$$

čia  $\sigma_{n,\mathbf{k}}^2 = \mathbf{E} X_{n,\mathbf{k}}^2$  (mes tariame, kad serijų schema yra centruota ir kad visų dispersijų suma yra vienetasis, t.y.  $b_n(\mathbf{k}_n) = 1$ ). Kiekvienam  $i = 1, \dots, d$  tegu

$$b_i(k) := b_n(k_n^1, \dots, k_n^{i-1}, k, k_n^{i+1}, \dots, k_n^d).$$

Apibrėžkime gardelės stačiakampį

$$Q_{n,\mathbf{k}} := \left[ b_1(k_1 - 1), b_1(k_1) \right) \times \dots \times \left[ b_d(k_d - 1), b_d(k_d) \right).$$

Darbe pasiūlytas *adaptyvus* daugiamatnio indekso sumavimo procesas tada yra apibrėžiamas taip:

$$\Xi_n(\mathbf{t}) = \sum_{1 \leq j \leq n} |Q_{n,j}|^{-1} |Q_{n,j} \cap [0, \mathbf{t}]| X_{n,j}.$$

Kaip ir sumavimo proceso  $\xi_n(\mathbf{t})$  atveju buvo gauta alternatyvi procesos  $\Xi_n(\mathbf{t})$  išraiška kurios pagalba buvo gautas įvertis

$$\sup_{j \geq J} 2^{\alpha j} \max_{v \in V_j} \|\lambda_{j,v}(\Xi_n)\| \leq 3^d \sum_{m=1}^d \left( \max_{j \geq J} 2^{j\alpha} \max_{r \in D_j} [\psi_n^{(m)}(r, r^-) + \psi_n^{(m)}(r, r^+)] + Z_n^{(m)} \right),$$

čia

$$\begin{aligned} \psi_n(r, r^-)^{(m)} &:= \max_{\mathbf{k}_{-m} \leq \mathbf{k}_n, -m} \left| \sum_{k_m=u_m(r^-)+2}^{u_m(r)} \Delta_{k_m}^{(m)} S_n(\mathbf{k}) \right| \\ \psi_n(r, r^+)^{(m)} &:= \max_{\mathbf{k}_{-m} \leq \mathbf{k}_n, -m} \left| \sum_{k_m=u_m(r)+2}^{u_m(r^+)} \Delta_{k_m}^{(m)} S_n(\mathbf{k}) \right| \\ Z_n^{(m)} &:= \max_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \frac{|\Delta_{k_m}^{(m)} S_n(\mathbf{k})|}{(\Delta b_m(k_m))^\alpha}. \end{aligned}$$

Kai  $d = 1$  pasiūlytas sumavimo procesas sutampa su Prokhorov konstrukcija. Šiuo atveju sąlygas FCRT Hiolderio erdvėse gavo Račkauskas ir Suquet[30]. Jie reikalavo tam tikros momentinės sąlygos bei kad serijų schema tenkintų begalinio mažumo sąlygą. Jų gauta momentinė sąlyga yra griežtesnė nei Lindebergo sąlyga, taigi iš jos iš karto išplaukia baigtiniamaičių pasiskirstymų konvergavimas. To nėra procesui  $\Xi_n(\mathbf{t})$ . Jo tirštumą charakterizuoja sekančios teoremos:

**Teorema 4** Tegu  $0 < \alpha < 1/2$  ir  $p(\alpha) := 1/(1/2 - \alpha)$ . Jei

$$\max_{1 \leq l \leq d} \max_{1 \leq k_l \leq k_n^l} \Delta b_l(k_l) \rightarrow 0, \text{ kai } m(\mathbf{n}) \rightarrow \infty.$$

ir tam tikram  $q > p(\alpha)$

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \sigma_{n,\mathbf{k}}^{-2q\alpha} \mathbf{E} |X_{n,\mathbf{k}}|^q = 0,$$

tai apibendrinta seka  $\{\Xi_n, \mathbf{n} \in \mathbb{N}\}$  yra asimptotiškai tiršta erdvėje  $H_\alpha^o([0, 1]^d)$ .

Kiekvienam  $\tau > 0$  apibrėžkime nupjautus atsitiktinius dydžius:

$$X_{n,\mathbf{k},\tau} := X_{n,\mathbf{k}} \mathbf{1}\{|X_{n,\mathbf{k}}| \leq \tau \sigma_{n,\mathbf{k}}^{2\alpha}\}$$

**Teorema 5** *Tegu*

$$\max_{1 \leq l \leq d} \max_{1 \leq k_l \leq k_n^l} \Delta b_l(k_l) \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty. \quad (19)$$

ir yra tenkinamos sąlygos:

i. Kiekvienam  $\varepsilon > 0$ ,

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} P(|X_{\mathbf{n}, \mathbf{k}}| \geq \varepsilon \sigma_{\mathbf{n}, \mathbf{k}}^{2\alpha}) = 0.$$

ii. Kiekvienam  $\varepsilon > 0$ ,

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \mathbf{E} X_{\mathbf{n}, \mathbf{k}}^2 \mathbf{1}\{|X_{\mathbf{n}, \mathbf{k}}| \geq \varepsilon\} = 0.$$

iii. Kuriam nors  $q > 1/(1/2 - \alpha)$ ,

$$\lim_{\tau \rightarrow 0} \lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \sigma_{\mathbf{n}, \mathbf{k}}^{-2q\alpha} \mathbf{E} |X_{\mathbf{n}, \mathbf{k}, \tau}|^q = 0.$$

Tada apibendrinta seka  $\{\Xi_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}\}$  yra asimptotiškai tiršta erdvėje  $H_\alpha^o([0, 1]^d)$ .

Šios dvi teoremos yra tiesioginis Račkausko ir Suquet įrodytų sąlygų FCRT erdvėje  $H_\alpha^o([0, 1]^d)$  apibendrinimas.

Nors sąlygos tirštumui užtikrina, kad serijų schemai galioja CRT, reikia papildomų sąlygų, kad baigtiniam procesui  $\Xi_{\mathbf{n}}(\mathbf{t})$  skirstiniai konverguotų. Trečiame skyriuje yra įrodoma, kad baigtiniam skirstinių ribos yra normaliosios, jei jos egzistuoja, bet tai nebūtinai Vynerio paklodės baigtiniam skirstiniai. Yra įrodoma, kad jei egzistuoja riba

$$\mu_{\mathbf{n}}(\mathbf{t}) := \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \mathbf{1}\{\mathbf{B}(\mathbf{k}) \in [0, \mathbf{t}]\} \sigma_{\mathbf{n}, \mathbf{k}}^2 \rightarrow \mu(\mathbf{t}), \text{ kai } m(\mathbf{n}) \rightarrow \infty,$$

kokiai nors funkcijai  $\mu(\mathbf{t})$ , tai ribinis procesas yra Gauso procesas  $G(\mathbf{t})$  su kovariacijos funkcija  $\mathbf{E} G(\mathbf{t}) G(\mathbf{s}) = \mu(\mathbf{s} \wedge \mathbf{t})$ . Toliau pateikiami pavyzdžiai kaip nuo serijų schemas dispersijų gali priklausyti ribinis pasiskirstymas.

**Pavyzdys 1** Tegu  $\mathbf{n} = (n, n)$ ,  $\mathbf{k}_n = (2n, 2n)$  ir  $X_{\mathbf{n}, \mathbf{k}} = a_{\mathbf{n}, \mathbf{k}} Y_{\mathbf{k}}$ , čia  $\{Y_{\mathbf{k}}, \mathbf{k} \leq \mathbf{k}_n\}$  n.v.p. atsitiktiniai dydžiai su standartiniu normaliuoju pasiskirstymu, o

$$a_{\mathbf{n}, \mathbf{k}}^2 = \begin{cases} \frac{1}{10n^2}, & \text{kai } \mathbf{k} \leq (n, n) \\ \frac{3}{10n^2}, & \text{kitais atvejais.} \end{cases} \quad (20)$$

Taip apibrėžta serijų schema tenkina (19), bet paprastų algebrinių veiksmų dėka galima įsitikinti, kad tokiai serijų schemai

$$\begin{aligned}\mu_n(\mathbf{t}) \rightarrow \nu(\mathbf{t}) := & \frac{1}{10} \left( \frac{5}{2} t_1 \wedge 1 \right) \left( \frac{5}{2} t_2 \wedge 1 \right) \\ & + \frac{(5t_1 - 2) \vee 0}{10} \left( \frac{5}{2} t_2 \vee 1 \right) \\ & + \frac{(5t_2 - 2) \vee 0}{10} \left( \frac{5}{2} t_1 \vee 1 \right) \\ & + \frac{((5t_1 - 2) \vee 0)((5t_2 - 2) \vee 0)}{30}.\end{aligned}$$

Taigi baigtiniamačiai pasiskirstymai nekonverguoja į Vynerio paklodės baigtiniamačius pasiskirstymus. Kitame pavyzdyje pateikiamas serijų schemas pavyzdys, kai  $\mu_n(\mathbf{t})$  diverguoja kiekvienam  $\mathbf{t}$ .

**Pavyzdys 2** Tegu  $\mathbf{n} = (n, n)$ ,  $\mathbf{k}_n = (n, n)$  ir  $X_{\mathbf{n}, \mathbf{k}} = b_{\mathbf{n}, \mathbf{k}} Y_{\mathbf{k}}$ , čia  $\{Y_{\mathbf{k}}, \mathbf{k} \leq \mathbf{k}_n\}$  n.v.p. atsitiktiniai dydžiai su standartiniu normaliuoju pasiskirstymu, o

$$b_{\mathbf{n}, \mathbf{k}}^2 = \begin{cases} \pi(\mathbf{k}_n)^{-1}, & \text{for } \mathbf{n} = (2l - 1, 2l - 1), l \in \mathbb{N} \\ a_{\mathbf{n}, \mathbf{k}}^2, & \text{for } \mathbf{n} = (2l, 2l), l \in \mathbb{N} \end{cases}$$

čia  $a_{\mathbf{n}, \mathbf{k}}$  apibrėžti kaip (20).

Ketvirtame skyriuje pateikiami teorinių rezultatų taikymai, konstruojant statistikas epideminių pasikeitimų aptikimui duomenyse su daugiamačiu indeksu. Iš pradžių yra nagrinėjamas vidurkio pasikeitimo dvimačio indekso imtyje  $\{X_{ij}, 1 \leq i \leq n, 1 \leq j \leq m\}$ . Laikoma, kad nulinė hipotezė yra

$(H_0)$ : Visų  $X_{ij}$  vidurkis yra vienodas ir lygus  $\mu_0$ ,  
o alternatyvioji hipotezė

$(H_A)$ : Egzistuoja tokie natūriniai skaičiai  $1 < a^* \leq b^* < n$ ,  $1 < c^* \leq d^* < m$  ir konstanta  $\mu_1 \neq \mu_0$  tokia, kad

$$\mathbf{E} X_{ij} = \mu_0 + \mu_1 \mathbf{1} \left( (i, j) \in [a^*, b^*] \times [c^*, d^*] \cap \mathbb{N}^2 \right).$$

Disertacijoje tokia alternatyva pavadinama *epideminiu stačiakampiu*. Tokiam pasikeitimui aptikti pasiūloma statistika, kuri yra klasikinė didžiausio tikėtinumo santykio statistika pasiūlyta Csörgő and Horváth [7] su tam tikru svoriu - epideminio stačiakampio diametru:

$$DUI(n, m, \alpha) = \max_{\substack{1 \leq a < b \leq n \\ 1 \leq c < d \leq m}} \frac{|\Delta_{b-a}^{(1)} \Delta_{d-c}^{(2)} S_{bd} - (s_b - s_a)(t_d - t_c) S_{nm}|}{\max\{s_b - s_a, t_d - t_c\}^\alpha},$$

čia  $s_i = i/n$  ir  $t_j = j/m$  kiekvienam  $1 \leq i \leq n, 1 \leq j \leq m$ . Yra parodoma, kad ši statistika yra sumavimo proceso  $\xi_n$  funkcionalas, kuris yra tolydus Hiolderio erdvėje  $H_\alpha^2([0, 1]^d)$ . Taigi pasinaudojus tolydaus atvaizdžio teorema bei invariantiškumo principu iš trečio skyriaus yra įrodomas pasiūlytos statistikos silpnas konvergavimas prie nulinės hipotezės:

$$\sigma^{-1}(nm)^{-1/2}DUI(n, m, \alpha) \xrightarrow{D} DUI(\alpha),$$

čia

$$DUI(\alpha) = \sup_{\mathbf{0} \leq \mathbf{s} < \mathbf{t} \leq \mathbf{1}} \frac{|\Delta_{[\mathbf{s}, \mathbf{t}]}W - (t_1 - s_1)(t_2 - s_2)W(\mathbf{1})|}{|\mathbf{t} - \mathbf{s}|^\alpha}$$

kai

$$\sup_{t>0} t^{1/(1/2-\alpha)} P(|X_{11}| > t) < \infty,$$

ir  $0 < \alpha < 1/2$ , laikant kad

$$\Delta_{[\mathbf{s}, \mathbf{t}]}x = x(\mathbf{t}) - x(s_1, t_2) - x(t_1, s_2) + x(\mathbf{s}).$$

bet kuriai dviejų kintamųjų funkcijai  $x$ . Taip pat yra įrodomas ir sukonstruoto testo suderinamumas:

**Teorema 6** *Prie alternatyvos ( $H_A$ ) tarkime, kad  $X_{ij}$  yra nepriklausomi ir kad  $\sigma_0^2 = \sup_n \text{var}(X_n) < \infty$ . Jei*

$$\lim_{m(n) \rightarrow \infty} (nm)^{1/2} \frac{h_{nm}}{d_{nm}^\alpha} |\mu_1 - \mu_0| \rightarrow \infty, \quad (21)$$

čia

$$h_{nm} = \frac{k^* l^*}{nm} \left( 1 - \frac{k^* l^*}{nm} \right) \text{ and } d_{nm} = \max \left\{ \frac{k^*}{n}, \frac{l^*}{m} \right\},$$

tai

$$(nm)^{-1/2}DUI(n, m, \alpha) \rightarrow \infty$$

Gautas rezultatas yra tiesioginis Račkausko ir Suquet [33] rezultato apibendrinimas. Jie nagrinėjo įprastinės (vienmačio indekso) imties atvejį bei testą skirtą aptikti vidurkio pasikeitimą prie taip vadinamos epideminės alternatyvos. Disertacijoje gautą rezultatą pritaikius vienmačio indekso imčiai,

gautumėme tą pačią testo statistiką, jos ribinį pasiskirstymą bei sąlygas suderinamumui. Bet gautame dvimačiame apibendrinime dalinimas iš epideminio stačiakampio diametro nebėra toks pat naudingas kaip dalinimas iš epideminio intervalo ilgio vienmačiu atveju. Tarkime, kad  $k^* = n^\gamma$ ,  $l^* = m^\delta$  ir kad  $\mu_1 - \mu_0$  nepriklauso nuo  $(n, m)$ . Tada sąlyga (21) atrodo taip:

$$\frac{n^{\gamma-1/2}m^{\delta-1/2}}{[n^{\gamma-1} \vee m^{\delta-1}]^\alpha} \rightarrow \infty.$$

Jei  $n^{\gamma-1} > m^{\delta-1}$  tai

$$n^{\gamma(1-\alpha)+\alpha-1/2}m^{\delta-1/2} \rightarrow \infty.$$

Matome, kad  $\gamma$  galima priklausomai nuo  $\alpha$  parinkti kiek norimai mažą ir suderinamumo sąlyga bus tenkinama, bet  $\delta$  negalime imti mažesnio nei  $1/2$ . Taigi gali būti aptiktos labai trumpos  $k^*$  epidemijos, bet geresnė  $l^*$  epidemijos ilgio eilė negali būti pasiekta. Sąlyga  $n^{\gamma-1} > m^{\delta-1}$ , kai  $\delta > 1/2$  yra patenkinama jei  $m > n^2$ , vadinasi norint aptikti labai trumpos epidemijas vienam indeksui, reikia daugiau duomenų kitam indeksui. Tokio apribojimo nėra vienmačiu atveju, nes tada  $h_n = l^*/n(1 - l^*/n)$  and  $d_n = l^*/n$ . Taigi epidemijai  $l^* = n^\gamma$  statistika diverguoja, kai  $\gamma > (1/2 - \alpha)/(1 - \alpha)$ .

Gautas apribojimą dvimačiu atveju galima panaikinti nagrinėjant pakeitus alternatyvą:

$(H'_A)$  : Egzistuoja natūriniai skaičiai ,  $1 < c^* \leq d^* < m$  ir konstanta  $\mu_1 \neq \mu_0$  tokia, kad

$$\mathbf{E} X_{ij} = \mu_0 + \mu_1 \mathbf{1} \left( (i, j) \in [1, n] \times [c^*, d^*] \cap \mathbb{N}^2 \right).$$

Testuoti tokiai alternatyvai disertacijoje pasiūloma modifikuota statistika

$$DUI_3(n, m, \alpha) = \max_{1 \leq c < d \leq m} \frac{|S_{n,d} - S_{n,c} - (t_d - t_c)S_{n,m}|}{(t_d - t_c)^\alpha}.$$

Prie panašių prielaidų yra parodoma, kad šios statistikos ribinis pasiskirstymas yra

$$DUI_3(\alpha) = \sup_{0 < s < t < 1} \frac{|W(1, t) - W(1, s) - (t - s)W(1, 1)|}{|t - s|^\alpha},$$

o suderinamumo sąlygos yra:

**Išvada 7** Nepriklausomų atsitiktinių dydžių šeimai  $X_{ij}$  su  $\sigma_0^2 = \sup_n \text{var}(X_n) < \infty$  prie alternatyvos  $H'_A$

$$(nm)^{-1/2} DUI_3(n, m, \alpha) \rightarrow \infty,$$

jei

$$\lim_{m(n) \rightarrow \infty} (nm)^{1/2} \left( \frac{l^*}{m} \right)^{1-\alpha} \left( 1 - \frac{l^*}{m} \right) |\mu_1 - \mu_0| \rightarrow \infty.$$

Toliau disertacijoje sprendžiami pasikeitimo uždaviniai panelinių duomenų regresijos modeliams. Nagrinėjami klasikiniai panelinių duomenų regresijos modeliai aprašyti Baltagi [2]:

$$y_{ij} = \mathbf{x}'_{ij} \boldsymbol{\beta} + u_{ij}, \quad (22)$$

$$y_{ij} = \mathbf{x}'_{ij} \boldsymbol{\beta} + \mu_i + u_{ij}, \quad (23)$$

Panelinių duomenų literatūroje modelis (22) paprastai vadinamas įprastinės regresijos modeliu, o modelis (23) - fiksuotų efektų regresijos modeliu.

Disertacijoje pasiūlomos statistikos skirtos nustatyti regresijos koeficiento  $\beta$  pasikeitimui. Statistikos išvedamos naudojantis idėja pasiūlyta Proberger and Krämer [23]. Iš pradžių įrodoma FCRT regresijos liekanoms, tada naudojamos statistikos vidurkio pasikeitimui, nes regresijos koeficiento pasikeitimas, keičia regresijos liekanų vidurkį.

Modelio (22) liekanos yra apibrėžiamos taip:

$$\hat{u}_{ij} = y_{ij} - \mathbf{x}'_{ij} \hat{\boldsymbol{\beta}} = u_{ij} - \mathbf{x}'_{ij} (\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}),$$

čia

$$\hat{\boldsymbol{\beta}} = \left( \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}'_{ij} \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} y_{ij}$$

yra koeficiento  $\boldsymbol{\beta}$  mažiausių kvadratų įvertis. Jeigu regresijos paklaidoms  $u_{ij}$  galioja FCRT ir  $\mathbf{x}_{ij}$  tenkina prielaidą:

**Prielaida A** Tegu  $x_{ij1} = 1$  visiems  $1 \leq (i, j) \leq (n, m)$  ir

$$\lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}'_{ij} = R,$$

tam tikrai neišsigimusiai  $(K \times K)$  matricai  $R$ . Taip pat tarkime, kad modelis yra pertvarkytas taip, kad

$$R = \begin{bmatrix} 1 & 0 \\ 0 & R^* \end{bmatrix}.$$

Tai savo ruožtu lemia, kad

$$\mathbf{c} \equiv \lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} = [1, 0, \dots, 0]'$$

tai teisinga tokia teorema

**Teorema 8** *Panelinių duomenų regresijos modeliui*

$$y_{ij} = \mathbf{x}_{ij}'\boldsymbol{\beta} + u_{ij},$$

*apibrėžkite sumavimo procesą*

$$\widehat{W}^{(n,m)}(t, s) = nm \sum_{i=1}^n \sum_{j=1}^m \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, s] \right| \widehat{u}_{ij}.$$

Tada

$$\frac{1}{\sigma\sqrt{nm}} \widehat{W}^{(n,m)}(t, s) \xrightarrow{D} W(t, s) - tsW(1, 1), \text{ as } n \wedge m \rightarrow \infty,$$

erdvėje  $H_\alpha^\alpha([0, 1]^2)$ , čia  $0 < \alpha < 1/2$ .

Fiksuotų efektų regresijos modeliui (23) regresijos liekanos yra apibrėžiamos taip:

$$\widehat{u}^{FE} = \widetilde{y}_{ij} - \widetilde{\mathbf{x}}_{ij}'\widehat{\boldsymbol{\beta}}^{FE},$$

čia  $\widetilde{y}_{ij} = y_{ij} - \bar{y}_i$  ir  $\widetilde{\mathbf{x}}_{ij}$  apibrėžiamas analogiškai, o  $\widehat{\boldsymbol{\beta}}^{FE}$  – klasikinis fiksuotų efektų koeficiento  $\boldsymbol{\beta}$  įvertis:

$$\widehat{\boldsymbol{\beta}}^{FE} = \left( \sum_{i=1}^n \sum_{j=1}^m \widetilde{\mathbf{x}}_{ij} \widetilde{\mathbf{x}}_{ij}' \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \widetilde{\mathbf{x}}_{ij} \widetilde{y}_{ij}.$$

Jeigu regresijos paklaidos  $u_{ij}$  tenkina FCRT bei daroma prielaida

**Prielaida B** *Tegu*

$$\lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \widetilde{\mathbf{x}}_{ij} \widetilde{\mathbf{x}}_{ij}' = \widetilde{R}$$

*tam tikrai neišsigimusiai  $(K \times K)$  matricai  $R$ , ir*

$$\lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} = \mathbf{c}$$

*tam tikram  $\mathbf{c} \in \mathbb{R}^K$ .*

tai įrodoma teorema

**Teorema 9** *Panelinių duomenų regresijos modeliui*

$$y_{ij} = \mathbf{x}'_{ij}\boldsymbol{\beta} + \mu_i + u_{ij},$$

*apibrėžkime sumavimo procesą*

$$\widehat{W}_{nm}^{FE}(t, s) = nm \sum_{i=1}^n \sum_{j=1}^m \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, s] \right| \widehat{u}_{ij}^{FE}.$$

*Tada*

$$\widehat{W}_{nm}^{FE}(t, s) \xrightarrow{D} W(t, s) - sW(t, 1), \text{ as } n \wedge m \rightarrow \infty,$$

*erdvėje  $H_\alpha^0([0, 1]^2)$ , čia  $0 < \alpha < 1/2$ .*

Abiems panelinių duomenų regresijos modeliams darbe išnagrinėtos lokalsios alternatyvos

$$\boldsymbol{\beta}_{ij} = \boldsymbol{\beta} + \frac{1}{\sqrt{nm}} \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right)$$

čia  $\mathbf{g}$  - vektorinė funkcija tolydi  $[0, 1]^2$ . Prie tų pačių prielaidų lokalių alternatyvų modeliams yra įrodoma FCRT erdvėje  $H_\alpha^0([0, 1]^d)$ . Ribinis procesas modeliui (22) tada yra

$$\begin{aligned} W(t, s) - tsW(1, 1) + \int_0^t \int_0^s \mathbf{c}' \mathbf{g}(u, v) du dv \\ - ts \mathbf{c}' \int_0^1 \int_0^1 \mathbf{g}(u, v) du dv, \end{aligned}$$

o modeliui (23)

$$\begin{aligned} W(t, s) - sW(t, 1) + \int_0^t \int_0^s \mathbf{c}' \mathbf{g}(u, v) du dv \\ - s \int_0^t \int_0^1 \mathbf{c}' \mathbf{g}(u, v) du dv. \end{aligned}$$

Regresijos koeficiento pasikeitimo testavimui darbe siūloma naudoti statistiką  $DUI(n, m, \alpha)$ , atsitiktinės dvimačio indekso imties elementų sumas pakeičiant regresijos liekanų sumomis. Laikoma, kad testuojama nulinė hipotezė

$(H_0)$  : Koeficientas  $\boldsymbol{\beta}$  nesikeičia,

prie alternatyvos

$(H_A)$  : Egzistuoja tokie natūriniai skaičiai  $1 < a^* \leq b^* < n, 1 < c^* \leq d^* < m$  ir konstantos  $\boldsymbol{\beta}_1 \neq \boldsymbol{\beta}_0$  tokios, kad tikrasis koeficientas  $\boldsymbol{\beta}$  yra išreiškiamas kaip

$$\boldsymbol{\beta} = \boldsymbol{\beta}_0 + \boldsymbol{\beta}_1 \mathbf{1} \left( (i, j) \in [a^*, b^*] \times [c^*, d^*] \cap \mathbb{N}^2 \right).$$

Prie nulinės hipotezės disertacijoje gaunamos ribinis statistikos pasiskirstymas abiemis panelinių duomenų regresijos modeliams. Įprastinei regresijai ribinis pasiskirstymas išlieka toks pat kaip ir atsitiktinės imties atveju:

$$\sigma^{-1}(nm)^{-1/2}\widehat{DUI}(n, m, \alpha) \xrightarrow{D} DUI(\alpha), \text{ as } n \wedge m \rightarrow \infty,$$

Fiksuotų efektų regresijai jis pasikeičia:

$$DUI^{FE}(\alpha) = \sup_{\mathbf{0} \leq \mathbf{s} < \mathbf{t} \leq \mathbf{1}} \frac{|\Delta_{[\mathbf{s}, \mathbf{t}]}W - (t_2 - s_2)[W(t_1, 1) - W(s_1, 1)]|}{|\mathbf{t} - \mathbf{s}|^\alpha}.$$

Darbe nurodomos sąlygos abiejų testų suderinamumui bei statistikų ribiniai pasiskirstymai prie lokalių alternatyvų.

Skyriaus pabaigoje yra pateikiamas gautų praktinių rezultatų aptarimas. Yra pastebima, kad praktikoje n.v.p. prielaida regresijos paklaidoms kartais yra pernelyg griežta, bet gautus rezultatus nesunku adaptuoti prielaidas pakeitus. Tam tereikia turėti invariantiškumo principą prie naujų prielaidų, statistikos ir jų ribiniai pasiskirstymai išliks tokie patys, jeigu tik ribinis procesas invariantiškumo principu yra Vynerio paklodė.

# Contents

|                                                         |           |
|---------------------------------------------------------|-----------|
| <b>Résumé substantiel</b>                               | <b>5</b>  |
| <b>Disertacijos santrauka</b>                           | <b>23</b> |
| <b>Notations</b>                                        | <b>43</b> |
| <b>Introduction</b>                                     | <b>45</b> |
| <b>1 Weak convergence in Hölder spaces</b>              | <b>52</b> |
| 1.1 General results . . . . .                           | 52        |
| 1.1.1 Basic definitions . . . . .                       | 52        |
| 1.1.2 Nets and asymptotic tightness . . . . .           | 53        |
| 1.1.3 Schauder decomposition . . . . .                  | 54        |
| 1.2 Hölder space and its properties . . . . .           | 55        |
| 1.2.1 Definition . . . . .                              | 55        |
| 1.2.2 Pyramidal functions . . . . .                     | 55        |
| 1.2.3 Tightness criteria . . . . .                      | 58        |
| 1.3 Results in probability . . . . .                    | 59        |
| 1.3.1 Gaussian processes . . . . .                      | 59        |
| 1.3.2 A variant of continuous mapping theorem . . . . . | 62        |
| 1.3.3 Rosenthal inequality . . . . .                    | 63        |
| 1.3.4 Doob inequality . . . . .                         | 64        |
| <b>2 Summation processes</b>                            | <b>67</b> |
| 2.1 Uniform variance case . . . . .                     | 67        |
| 2.1.1 Differences of partial sums . . . . .             | 67        |
| 2.1.2 Definitions and representations . . . . .         | 68        |
| 2.1.3 Estimate of sequential Hölder norm . . . . .      | 74        |
| 2.2 Unequal variance . . . . .                          | 80        |
| 2.2.1 Definitions and representations . . . . .         | 80        |
| 2.2.2 Estimate of sequential norm . . . . .             | 84        |

|          |                                                                  |            |
|----------|------------------------------------------------------------------|------------|
| <b>3</b> | <b>Functional central limit theorems</b>                         | <b>90</b>  |
| 3.1      | Invariance principle . . . . .                                   | 90         |
| 3.1.1    | Finite dimensional distributions . . . . .                       | 90         |
| 3.1.2    | Necessity . . . . .                                              | 94         |
| 3.1.3    | Tightness . . . . .                                              | 96         |
| 3.1.4    | Corollaries . . . . .                                            | 101        |
| 3.1.5    | Truncated variables . . . . .                                    | 104        |
| 3.2      | Triangular array . . . . .                                       | 107        |
| 3.2.1    | Finite dimensional distributions . . . . .                       | 107        |
| 3.2.2    | Tightness . . . . .                                              | 113        |
| 3.2.3    | Corollaries . . . . .                                            | 121        |
| <b>4</b> | <b>Applications</b>                                              | <b>126</b> |
| 4.1      | Tests for epidemic alternatives . . . . .                        | 128        |
| 4.1.1    | Epidemic rectangles . . . . .                                    | 128        |
| 4.1.2    | Some special cases . . . . .                                     | 134        |
| 4.2      | Functional central limit theorems for panel data regressions . . | 137        |
| 4.2.1    | Models and the assumptions . . . . .                             | 137        |
| 4.2.2    | Local alternatives . . . . .                                     | 142        |
| 4.3      | Change point statistics for panel regressions . . . . .          | 146        |
| 4.3.1    | Tests and their behaviour under null hypothesis . . . .          | 146        |
| 4.3.2    | Consistency of the epidemic alternatives . . . . .               | 148        |
| 4.3.3    | Practical considerations and discussion . . . . .                | 151        |
|          | <b>Bibliography</b>                                              | <b>155</b> |

# Notations

$\mathbf{t}$  denotes a real vector  $(t_1, \dots, t_d)$ .

$\mathbb{R}^d$  denotes the set of real vectors  $\mathbf{t}$ .

$\mathbb{N}$  denotes the set of natural numbers,  $\mathbb{N} = \{1, 2, \dots\}$ .

$\mathbb{Z}$  denotes the set of integers,  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$ .

$\mathbb{H}$  denotes Hilbert space.

$\mathbf{0}$  denotes element  $(0, \dots, 0)$  from the space  $\mathbb{R}^d$ .

$\mathbf{1}$  denotes element  $(1, \dots, 1)$  from the space  $\mathbb{R}^d$ .

$\mathbf{t}_{k:l}$  denotes the “subvector”  $(t_k, t_{k+1}, \dots, t_l)$ .

$\mathbf{t}_{-k}$  denotes the “subvector”  $(t_1, \dots, t_{k-1}, t_{k+1}, \dots, t_d)$ .

$\mathbf{t}_K$  denotes the subvector  $(t_{k_1}, \dots, t_{k_l})$  with  $K = \{k_1, \dots, k_l\}$  and  $1 \leq k_1 < k_2 < \dots < k_l \leq d$ .

$\mathbf{s} \leq \mathbf{t}$  means  $s_k \leq t_k$  for all  $k = 1, \dots, d$ .

$|\mathbf{t}|$  denotes  $\max(|t_1|, \dots, |t_d|)$  for  $\mathbf{t} \in \mathbb{R}^d$ .

$|A|$  denotes Lebesgue measure for the set  $A \subset \mathbb{R}^d$ .

$\text{card } A$  denotes the cardinality of the set  $A \subset \mathbf{R}^d$ .

$\mathbf{s}\mathbf{t}$  denotes  $(s_1 t_1, \dots, s_d t_d)$ ,  $\mathbf{s}/\mathbf{t}$  denotes  $(s_1/t_1, \dots, s_d/t_d)$ .

$\boldsymbol{\pi}(\mathbf{n})$  denotes  $n_1 \cdots n_d$  for  $\mathbf{n} \in \mathbb{N}^d$ .

$\text{m}(\mathbf{t})$  denotes  $\min(t_1, \dots, t_d)$ .

$[x]$  denotes integer part of real number  $x$ .

$\{x\}$  denotes the fractional part of real number  $x$ , ( $\{x\} = x - [x]$ ).

$[\mathbf{t}]$  denotes  $([t_1], \dots, [t_d])$ , respectively  $\{\mathbf{t}\}$  denotes  $(\{t_1\}, \dots, \{t_d\})$ .

$x \wedge y$  denotes  $\min(x, y)$  for real numbers  $x$  and  $y$ .

$x \vee y$  denotes  $\max(x, y)$  for real numbers  $x$  and  $y$ .

$\mathbf{t} \wedge \mathbf{s}$  denotes  $(t_1 \wedge s_1, \dots, t_d \wedge s_d)$ , respectively  $\mathbf{t} \vee \mathbf{s}$  denotes  $(t_1 \vee s_1, \dots, t_d \vee s_d)$ .

$\|\cdot\|$  denotes the norm of Hilbert space  $\mathbb{H}$ .

$\langle \cdot, \cdot \rangle$  denotes the scalar product of Hilbert space  $\mathbb{H}$ .

$C([0, 1]^d)$  denotes the set of continuous functions  $x : [0, 1]^d \rightarrow \mathbb{R}$ .

$H_\alpha^o([0, 1]^d)$  denotes the set of continuous functions  $x : [0, 1]^d \rightarrow \mathbb{R}$  satisfying

$$\lim_{\delta \rightarrow 0} \sup_{|\mathbf{t} - \mathbf{s}| < \delta} |x(\mathbf{t}) - x(\mathbf{s})| / |\mathbf{t} - \mathbf{s}|^\alpha = 0.$$

$H_\alpha^o(\mathbb{H})$  denotes the set of continuous functions  $x : [0, 1]^d \rightarrow \mathbb{H}$  satisfying

$$\lim_{\delta \rightarrow 0} \sup_{|\mathbf{t} - \mathbf{s}| < \delta} \|x(\mathbf{t}) - x(\mathbf{s})\| / |\mathbf{t} - \mathbf{s}|^\alpha = 0.$$

$X_\alpha = Y_\alpha + o_P(1)$  iff  $\|X_\alpha - Y_\alpha\| \rightarrow 0$  in probability.

$\xrightarrow{D}$  denotes weak convergence in the space specified afterwards. If no space is specified it is assumed that weak convergence takes place in  $\mathbb{R}$ .

For the net  $\{X_\alpha\}$  of Banach space valued random variables

$\mathbf{1}(\cdot)$  denotes the indicator function.

$\Delta_s^{(m)}$  denotes the difference operator acting on  $m$ -th coordinate,  $\Delta_s^{(1)}x(\mathbf{t}) = x(\mathbf{t}) - x((t_1 - s, t_2, \dots, t_d))$ .

$\xi_n$  denotes polygonal summation process.

$\xi_n$  denotes continuous multi-parameter summation process.

# Introduction

Convergence of stochastic processes to some Brownian motion or related process is an important topic in probability theory and mathematical statistics. The first functional central limit theorem by Donsker and Prokhorov states the  $C[0, 1]$ -weak convergence of  $n^{-1/2}\xi_n$  to the standard Brownian motion  $W$ . Here  $\xi_n$  denotes the random polygonal line process indexed by  $[0, 1]$ :

$$\xi_n(t) = S_{[nt]} + (nt - [nt])X_{[nt]+1}, t \in [0, 1],$$

where  $S_0, S_k := X_1 + \dots + X_k, k \geq 1$ , are the partial sums of a sequence  $(X_i)_{i \geq 1}$  of i.i.d. random variables such that  $\mathbf{E} X_1 = 0$  and  $\mathbf{E} X_1^2 = 1$ . (We say that sequence of random elements  $Y_n$  with values in separable metric space  $B$  converges weakly to random element  $Y$ , if  $\mathbf{E} f(Y_n) \rightarrow \mathbf{E} f(Y)$ , for every continuous bounded functional  $f$ ).

This theorem implies via continuous mapping the convergence in distribution of  $f(n^{-1/2}\xi_n)$  to  $f(W)$  for any continuous functional  $f : C[0, 1] \rightarrow \mathbb{R}$ . Clearly this provides many statistical applications. On the other hand, considering that the paths of  $\xi_n$  are piecewise linear and that  $W$  has roughly speaking, an  $\alpha$ -Hölder regularity for any exponent  $\alpha < 1/2$ , it is tempting to look for a stronger topological framework for the weak convergence of  $n^{-1/2}\xi_n$  to  $W$ . In addition to the satisfaction of mathematical curiosity, the practical interest of such an investigation is to obtain a richer set of continuous functionals of the paths. For instance, Hölder norms of  $\xi_n$  are closely related to some test statistics to detect short “epidemic” changes in the distribution of the  $X_i$ ’s, see [32, 34].

In 1962, Lamperti [20] obtained the first functional central limit theorem in the separable Banach spaces  $H_\alpha^o, 0 < \alpha < 1/2$ , of functions  $x : [0, 1] \rightarrow \mathbb{R}$  such that

$$\|x\|_\alpha := |x(0)| + w_\alpha(x, 1) < \infty,$$

with

$$w_\alpha(x, \delta) := \sup_{0 < |t-s| \leq \delta} \frac{|x(t) - x(s)|}{|t - s|^\alpha} \xrightarrow{\delta \rightarrow 0} 0.$$

More precise definitions are given in the section 1.2.

Assuming that  $\mathbf{E} |X_1|^q < \infty$  for some  $q > 2$ , he proved the weak convergence of  $n^{-1/2}\xi_n$  to  $W$  in the Hölder space  $H_\alpha^o$  for any  $\alpha < 1/2 - 1/q$ . Račkauskas and Suquet in [31] (see also [28]) obtained a necessary and sufficient condition for the Lamperti's functional central limit theorem. Namely for  $0 < \alpha < 1/2$ ,  $n^{-1/2}\xi_n$  converges weakly in  $H_\alpha^o$  to  $W$  if and only if

$$\lim_{t \rightarrow \infty} t^{p(\alpha)} P(|X_1| > t) = 0, \quad (24)$$

where

$$p(\alpha) := \frac{1}{\frac{1}{2} - \alpha}. \quad (25)$$

Further extensions of Donsker-Prokhorov's functional central limit theorem concern summation processes. Let  $|A|$  denote the Lebesgue measure of the Borel subset  $A$  of  $\mathbb{R}^d$ . For a collection  $\mathcal{A}$  of Borel subsets of  $[0, 1]^d$ , summation process  $\{\xi_n(A); A \in \mathcal{A}\}$  based on a random field  $\{X_j, \mathbf{j} \in \mathbb{N}^d\}$ , of independent identically distributed real random variables with zero mean is defined by

$$\xi_n(A) = \sum_{1 \leq \mathbf{j} \leq \mathbf{n}} |R_{\mathbf{n}, \mathbf{j}}|^{-1} |R_{\mathbf{n}, \mathbf{j}} \cap A| X_j,$$

where  $\mathbf{j} = (j_1, \dots, j_d)$ ,  $\mathbf{n} = (n_1, \dots, n_d)$ ,  $R_{\mathbf{n}, \mathbf{j}}$  is the “rectangle”

$$R_{\mathbf{n}, \mathbf{j}} := \left[ \frac{j_1 - 1}{n_1}, \frac{j_1}{n_1} \right) \times \dots \times \left[ \frac{j_d - 1}{n_d}, \frac{j_d}{n_d} \right)$$

and the indexation condition “ $\mathbf{1} \leq \mathbf{j} \leq \mathbf{n}$ ” is understood componentwise :  $1 \leq j_1 \leq n_1, \dots, 1 \leq j_d \leq n_d$ . Of special interest are the partial sum processes based on the collection of sets  $\mathcal{A} = \mathcal{Q}_d$  where

$$\mathcal{Q}_d := \left\{ [0, t_1] \times \dots \times [0, t_d]; \mathbf{t} = (t_1, \dots, t_d) \in [0, 1]^d \right\}, \quad (26)$$

Note that when  $d = 1$  the partial sum process  $\xi_n$  based on  $\mathcal{Q}_d$  is the random polygonal line of Donsker-Prokhorov's theorem.

By equipping the collection  $\mathcal{A}$  with some pseudo-metric  $\delta$ , one define the space  $C(\mathcal{A})$  of real continuous functions on  $\mathcal{A}$ , endowed with the norm

$$\|f\|_{\mathcal{A}} := \sup_{A \in \mathcal{A}} |f(A)|.$$

The usual semimetrics are  $\delta(A, B) = \sqrt{|A \Delta B|}$ , or  $\delta(A, B) = \sqrt{m(A \Delta B)}$ , for  $A, B \in \mathcal{A}$ , where  $m$  is a probability measure on the  $\sigma$ -algebra of Borel

subsets of  $[0, 1]^d$ . When  $\mathcal{A}$  is totally bounded with respect to  $\delta$ ,  $C(\mathcal{A})$  is a separable Banach space.

A Brownian sheet process indexed by  $\mathcal{A}$  is a mean zero Gaussian process  $W$  with sample paths in  $C(\mathcal{A})$  and

$$\mathbf{E} W(A)W(B) = |A \cap B|, \quad A, B \in \mathcal{A}. \quad (27)$$

Existence of such process is proved by placing restrictions on collection  $\mathcal{A}$  which are usually expressed by some condition on its metric entropy. Dudley [11] and Erickson [13] give conditions for  $W$  to exist in a general Hölder space  $H_\rho(\mathcal{A})$ . It is defined as the subspace of the space  $C(\mathcal{A})$  of the functions satisfying

$$\sup_{0 < \delta(A, B) < 1} \frac{|x(A) - x(B)|}{\rho(\delta(A, B))} < \infty,$$

with the weight function  $\rho$ . For  $\rho(h) = h^\alpha$ , Erickson [13] proves that for process  $W$ , the Hölder exponent  $\alpha$  cannot exceed  $1/2$  and it decreases as the entropy of  $\mathcal{A}$  increases. The functional central limit theorem (FCLT) in  $C(\mathcal{A})$  or in  $H_\rho(\mathcal{A})$  means the weak convergence of the summation process  $\{\xi_n(A); A \in \mathcal{A}\}$ , suitably normalized, to a Brownian sheet process indexed by  $\mathcal{A}$ .

The first FCLT for  $\{\xi_n(A); A \in \mathcal{Q}_d\}$  in  $C(\mathcal{Q}_d)$  was established by Kuelbs [19] under some moment restrictions and by Wichura [42] under finite variance condition. In 1983, Pyke [25] derived a FCLT for summation process in  $C(\mathcal{A})$ , provided that the collection  $\mathcal{A}$  satisfies the bracketing entropy condition. However, his result required moment conditions which depend on the size of the collection  $\mathcal{A}$ . Bass [3] and simultaneously Alexander and Pyke [1] extended Pyke's result to i.i.d. random fields with finite variance. Further developments were concerned with relaxing entropy conditions on the collection  $\mathcal{A}$ , Ziegler [43], and with relaxing i.i.d. condition on the random field  $\{X_n, \mathbf{n} \in \mathbb{N}^d\}$ , Dedecker [8], El Machkouri and Ouchti [12] to name a few.

The FCLT for summation process in  $H_\rho(\mathcal{A})$  is not so extensively studied. Most general results are provided by Erickson [13] who shows that if  $\mathbf{E}|X_j|^q < \infty$  for some  $q > 2$  then the FCLT holds in  $H_\rho(\mathcal{A})$  for some  $\rho$  which depends on  $q$  and properties of  $\mathcal{A}$ . For  $d = 1$  and the class  $\mathcal{A}$  of intervals  $[0, t]$ ,  $0 \leq t \leq 1$ , Erickson's results coincide with Lamperti's ones [20], whereas his case  $d > 1$  requires moments of order  $q > dp(\alpha)$  with the same  $p(\alpha)$  as in (25). In Račkauskas and Zemlys [36], the result by Erickson was improved in the case  $d = 2$ . In thesis this result was extended for  $d > 2$  and Hilbert space valued random variables. Before stating it in full we need some

definitions.

With  $\mathbb{H}$  as the real separable Hilbert space, define the Hölder space  $\mathbb{H}_\alpha^o([0, 1]^d)$  of Hilbert-valued multi-parameter functions as the vector space of functions  $x : [0, 1]^d \rightarrow \mathbb{H}$  such that

$$\|x\|_\alpha := \|x(0)\| + w_\alpha(x, 1) < \infty,$$

with

$$w_\alpha(x, \delta) := \sup_{0 < |\mathbf{t} - \mathbf{s}| \leq \delta} \frac{\|x(\mathbf{t}) - x(\mathbf{s})\|}{|\mathbf{t} - \mathbf{s}|^\alpha} \xrightarrow{\delta \rightarrow 0} 0.$$

Note that for  $\mathbb{H} = \mathbb{R}$ , the space  $\mathbb{H}_\alpha^o([0, 1]^d)$  is a subset of  $H_\rho(\mathcal{Q}_d)$  with  $\rho(h) = h^\alpha$  and  $\mathcal{Q}_d$  defined by (3).

Define  $\mathbb{H}$ -valued Brownian sheet  $W$  with the covariance operator  $\Gamma$ , as a  $\mathbb{H}$ -valued zero mean Gaussian process indexed by  $[0, 1]^d$  and satisfying

$$\mathbf{E} \langle W(\mathbf{t}), x \rangle \langle W(\mathbf{s}), y \rangle = (t_1 \wedge s_1) \cdots (t_d \wedge s_d) \langle \Gamma x, y \rangle \quad (28)$$

for  $\mathbf{t}, \mathbf{s} \in [0, 1]^d$  and  $x, y \in \mathbb{H}$ . For  $\mathbb{H} = \mathbb{R}$ , the space of covariance operators is isomorphic to  $\mathbb{R}_+$  and (28) collapses to

$$\mathbf{E} W(\mathbf{t}) W(\mathbf{s}) = \sigma^2(t_1 \wedge s_1) \cdots (t_d \wedge s_d).$$

which is the same as (27) for  $A, B \in \mathcal{Q}_d$  and  $\sigma^2 = 1$ .

The following theorem holds.

**Theorem 1** *For  $0 < \alpha < 1/2$ , set  $p = p(\alpha) := 1/(1/2 - \alpha)$ . For  $d \geq 2$ , let  $\{X_{\mathbf{j}}; \mathbf{j} \in \mathbb{N}^d, \mathbf{j} \geq \mathbf{1}\}$  be an i.i.d. collection of square integrable centered random elements in the separable Hilbert space  $\mathbb{H}$  and  $\xi_{\mathbf{n}}$  be the summation process defined by*

$$\xi_{\mathbf{n}}(\mathbf{t}) = \sum_{1 \leq j \leq n} |R_{\mathbf{n}, j}|^{-1} |R_{\mathbf{n}, j} \cap [0, \mathbf{t}]| X_{\mathbf{j}}.$$

*Let  $W$  be a  $\mathbb{H}$ -valued Brownian sheet with the same covariance operator as  $X_{\mathbf{1}}$ . Then the convergence*

$$(n_1 \cdots n_d)^{-1/2} \xi_{\mathbf{n}} \xrightarrow[m(\mathbf{n}) \rightarrow \infty]{\mathbb{H}_\alpha^o([0, 1]^d)} W$$

*holds if and only if*

$$\sup_{t > 0} t^{p(\alpha)} P(\|X_{\mathbf{1}}\| > t) < \infty. \quad (29)$$

As we see, condition (29) does not depend on the dimension  $d$  provided  $d > 1$  and is weaker than necessary and sufficient condition (24) in the extension by Račkauskas and Suquet of Lamperti's functional central limit theorem. Moreover, we show that summation process considered along the diagonal, namely the sequence  $n^{-d/2}\xi_n = n^{-d/2}\xi_{n,\dots,n}$ ,  $n \in \mathbb{N}$ , converges in  $\mathbb{H}_\alpha^o([0, 1]^d)$  if and only if

$$\lim_{t \rightarrow \infty} t^{2d/(d-2\alpha)} P(\|X_1\| > t) = 0. \quad (30)$$

As dimension  $d$  increases, this condition weakens. For example, (30) is satisfied for any  $d > 1$  provided  $\mathbf{E} \|X_1\|^4 < \infty$ . This again shows up a difference between the cases  $d = 1$  and  $d > 1$  for functional central limit theorems in Hölder spaces.

The result in theorem 1 was obtained together with Račkauskas and Suquet [35]. Its proof and the prerequisites take up a sizeable part of the thesis. Necessary results from Hölder spaces and probability theory are given in the chapter 1. The properties of the summation process  $\xi_n$  are given in the section 2.1 and the result is proved in the section 3.1.

After i.i.d. case we considered the case of triangular array, when random variables are independent but not identically distributed. For general summation processes, the case of non-identically distributed variables was investigated by Goldie and Greenwood [14], [15]. They used classical construction of summation process, so their result does not coincide with classical Prokhorov [24] result for adaptive polygonal line process  $\Xi_n$  indexed by  $[0, 1]$  with vertices  $(b_n(k), S_n(k))$ , where  $b_n(k) = \mathbf{E} X_{n,1}^2 + \dots + \mathbf{E} X_{n,k}^2$ , with assumption that  $b_n(k_n) = 1$ , and  $X_{n,k}$  – independent non-identically distributed random variables.

The attempt to introduce adaptive construction for general summation processes was made by Bickel and Wichura [5]. However they put some restrictions on variance of random variables in triangular array. For zero mean independent random variables  $\{X_{n,ij}, 1 \leq i \leq I_n, 1 \leq j \leq J_n\}$  with variances  $\mathbf{E} X_{n,ij}^2 = a_{n,i}b_{n,j}$  satisfying  $\sum a_{n,i} = 1 = \sum b_{n,j}$ , they defined summation process as

$$\zeta_n(t_1, t_2) = \sum_{i \leq A_n(t_1)} \sum_{j \leq B_n(t_2)} X_{n,ij},$$

where

$$A_n(t_1) = \max\{k : \sum_{i \leq k} a_{n,i} < t_1\}, \quad B_n(t_2) = \max\{l : \sum_{j \leq l} b_{n,j} < t_2\}.$$

It is easy to see that this construction is two-dimensional time generalization

of jump version of Prokhorov construction. Bickel and Wichura proved that the process  $\zeta_n$  converges in the space  $D([0, 1]^2)$  to a Brownian sheet, if  $a_{n,i}$  and  $b_{n,j}$  are infinitesimally small and the random variables  $\{X_{n,ij}\}$  satisfy Lindeberg condition.

In this contribution we introduced new adaptive construction of summation process which reduces to classical construction for triangular arrays in one dimensional case. Sufficient conditions for the weak convergence in Hölder spaces are given. For the case  $d = 1$  they coincide with conditions given by Račkauskas and Suquet. The limiting process in general case is not a standard Brownian sheet. It is a mean zero Gaussian process with covariance depending on the limit of  $\mathbf{E} \Xi_n(\mathbf{t})^2$ . Examples of possible limiting processes are given. In case of special variance structure of triangular array as in Bickel and Wichura it is shown that the limiting process is a standard Brownian sheet.

Finally we provide the application of the theoretical results by constructing statistics for detecting the epidemic change in a given data with multi-dimensional indexes. Such data naturally arise if for example we measure some property of sample of individuals through time. It is natural then to assign two indexes to observation, the number of the individual and the time period when it was observed. This is so called longitudinal or panel data. First we consider the detection of the change of the mean in the double indexed sample  $\{X_{ij}, 1 \leq i \leq n, 1 \leq j \leq m\}$ . We test the null hypothesis of no change in mean against the alternative hypothesis of the change in a *epidemic rectangle*, i.e. the mean is different for indexes in the rectangle  $D^* = [a^*, b^*] \times [c^*, d^*] \cap \mathbb{N}^2$ . Our proposed statistic for detecting such change is the classical likelihood ratio statistic of Csörgő and Horváth [7], weighted with the power of diameter of the epidemic rectangle. We show that this statistic is the functional of summation process  $\xi_n$ , with the functional continuous in the Hölder space. Thus using continuous mapping theorem and our theoretical result we find the asymptotic distribution of our statistic. We give the conditions for the consistency of the test and show that division by diameter, improves the ability to detect shorter epidemics, but that the result is not optimal compared to the one-dimensional case considered by Račkauskas and Suquet [33].

Next we turn our attention to panel regression models. We consider classical pooled or ordinary least squares and fixed effects regressions described by Baltagi [2]. We prove functional central limit theorem (FCLT) for the regression residuals under condition that regression disturbances satisfy FCLT and classical conditions on the regressors. This result generalizes the result of Ploberger and Krämer [23] for the time-series regression. Using the FCLT for regression residuals we adapt our statistic for detecting the change of the

mean, to detect the change of the regression coefficient in both regression models. We find asymptotic distributions and give conditions for consistency of the statistics. We also investigate the behaviour of these statistics under local alternatives and derive results similar to those of Ploberger and Krämer.

# Chapter 1

## Weak convergence in Hölder spaces

### 1.1 General results

#### 1.1.1 Basic definitions

Let us introduce some notation. Vectors  $\mathbf{t} = (t_1, \dots, t_d)$  of  $\mathbb{R}^d$ ,  $d \geq 2$ , are typeset in italic bold. In particular,

$$\mathbf{0} := (0, \dots, 0), \quad \mathbf{1} := (1, \dots, 1).$$

For  $1 \leq k < l \leq d$ ,  $\mathbf{t}_{k:l}$  denotes the “subvector”

$$\mathbf{t}_{k:l} := (t_k, t_{k+1}, \dots, t_l),$$

$\mathbf{t}_{-k}$  denotes the “subvector”

$$\mathbf{t}_{-k} = (t_1, \dots, t_{k-1}, t_{k+1}, \dots, t_d).$$

and  $\mathbf{t}_K$  denotes the “subvector”

$$\mathbf{t}_K = (t_{k_1}, \dots, t_{k_m}),$$

with  $K = \{k_1, \dots, k_m\} \subset \{1, \dots, d\}$  and  $1 \leq k_1 < k_2 < \dots < k_m \leq d$ . The set  $\mathbb{R}^d$  is equipped with the partial order

$$\mathbf{s} \leq \mathbf{t} \quad \text{if and only if} \quad s_k \leq t_k, \quad \text{for all } k = 1, \dots, d. \quad (1.1)$$

As a vector space,  $\mathbb{R}^d$  is endowed with the norm

$$|\mathbf{t}| = \max(|t_1|, \dots, |t_d|), \quad \mathbf{t} = (t_1, \dots, t_d) \in \mathbb{R}^d.$$

Together with the usual addition of vectors and multiplication by a scalar, we use also the componentwise multiplication and division of vectors  $\mathbf{s} = (s_1, \dots, s_d)$ ,  $\mathbf{t} = (t_1, \dots, t_d)$  in  $\mathbb{R}^d$  defined whenever it makes sense, by

$$\mathbf{s}\mathbf{t} := (s_1 t_1, \dots, s_d t_d), \quad \mathbf{s}/\mathbf{t} := (s_1/t_1, \dots, s_d/t_d).$$

Partial order as well as all these operations are also intended componentwise when one of the two involved vectors is replaced by a scalar. So for  $c \in \mathbb{R}$  and  $\mathbf{t} \in \mathbb{R}^d$ ,  $c \leq \mathbf{t}$  means  $c \leq t_k$  for  $k = 1, \dots, d$ ,  $\mathbf{t} + c := (t_1 + c, \dots, t_d + c)$ ,  $c/\mathbf{t} := (c/t_1, \dots, c/t_d)$ .

For  $\mathbf{n} = (n_1, \dots, n_d) \in \mathbb{N}^d$ , we write

$$\boldsymbol{\pi}(\mathbf{n}) := n_1 \cdots n_d,$$

and for  $\mathbf{t} = (t_1, \dots, t_d) \in \mathbb{R}^d$ ,

$$m(\mathbf{t}) := \min(t_1, \dots, t_d).$$

For any real number  $x$ , denote by  $[x]$  and  $\{x\}$  its integer part and fractional part defined respectively by

$$[x] \leq x < [x] + 1, \quad [x] \in \mathbb{Z} \quad \text{and} \quad \{x\} := x - [x].$$

When applied to vectors  $\mathbf{t}$  of  $\mathbb{R}^d$ , these operations are defined componentwise:

$$[\mathbf{t}] := ([t_1], \dots, [t_d]), \quad \{\mathbf{t}\} := (\{t_1\}, \dots, \{t_d\}).$$

The context should dispel any notational confusion between the fractional part of  $x$  (or  $\mathbf{t}$ ) and the set having  $x$  (or  $\mathbf{t}$ ) as unique element.

We denote by  $\mathbb{H}$  a separable real Hilbert space with norm  $\|\cdot\|$  and inner product  $\langle \cdot, \cdot \rangle$ .

### 1.1.2 Nets and asymptotic tightness

Let  $A$  be a set with a partial order and let  $A$  be a directed set. For a general topological space  $X$ , a map from  $A$  to  $X$  is called a net and denoted by  $\{x_\alpha, \alpha \in A\} \subset X$ . We say that this net has a limit  $x$  if and only if for every neighborhood  $U$  of  $x$  there exists  $\alpha_0 \in A$  such that  $x_\alpha \in U$  for each  $\alpha_0 \leq \alpha$ . When the space  $X$  is Hausdorff, any net in  $X$  has *at most* one limit. All the

spaces we are dealing with are Banach, therefore Hausdorff, so it is always implicit that if the limit of the net exists, it is unique.

We are mainly interested in the nets  $\{x_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}^d\}$ , where  $\mathbb{N}^d$  is a directed set with partial order  $\mathbf{s} \leq \mathbf{t}$  defined in (1.1). Note that if we have  $\mathbf{n}_0 \leq \mathbf{n}$ , then  $m(\mathbf{n}) \geq m(\mathbf{n}_0)$  and if  $m(\mathbf{n}) \geq N$ , then  $(N, \dots, N) \leq \mathbf{n}$ . Thus if the net  $\{x_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}^d\}$  has the limit  $x$  it makes sense to write

$$\lim_{m(\mathbf{n}) \rightarrow \infty} x_{\mathbf{n}} = x.$$

We will use this notation throughout the thesis.

Let  $B$  be some separable Banach space and  $(Y_{\alpha})_{\alpha \in A}$  be a net of random elements in  $B$ . We write

$$Y_{\alpha} \xrightarrow{B} Y,$$

for weak convergence in the space  $B$  to the random element  $Y$ , i.e.  $\mathbf{E} f(Y_{\alpha}) \rightarrow \mathbf{E} f(Y)$  for any continuous and bounded  $f : B \rightarrow \mathbb{R}$ .

For proving weak convergence of the nets we use some variant of Prokhorov's theorem (see e.g. van der Vaart and Wellner [41] p.21 theorem 1.3.9) which asserts that the net  $\{Y_{\alpha}\}$  has a weakly convergent subnet if it is asymptotically tight, i. e. for each  $\varepsilon > 0$  there exists a compact set  $K_{\varepsilon} \in B$  such that

$$\liminf_{\alpha} P(Y_{\alpha} \in K_{\varepsilon}) > 1 - \varepsilon. \quad (1.2)$$

Thus weak convergence of the net  $Y_{\alpha}$  can be proved by classical approach, by checking the property of asymptotical tightness and proving the convergence of the finite-dimensional distributions.

### 1.1.3 Schauder decomposition

To check the property of asymptotical tightness we need some way of characterizing compact subsets of the paths space. Suquet [39] gives us a criteria exploiting the notion of Schauder decomposition.

**Definition 1** *An infinite sequence  $(\mathcal{B}_j, j \in \mathbb{N})$  of closed linear subspaces of a Banach space  $B$  such that  $\mathcal{B}_j \neq 0$  ( $j \in \mathbb{N}$ ) is called a Schauder decomposition of  $B$  if for every  $x \in B$  there exists a unique sequence  $(y_n, n \in \mathbb{N})$  with  $y_j \in \mathcal{B}_j$  ( $j \in \mathbb{N}$ ) such that:*

$$x = \sum_{j=0}^{\infty} y_j$$

and if the coordinate projections defined by  $v_n(x) = y_n$ , are continuous on  $B$ .

Let us denote  $\mathcal{Z}_j = \bigoplus_{i \leq j} \mathcal{B}_i$  and  $E_j = \sum_{i \leq j} v_i$  the continuous projections of  $B$  onto  $\mathcal{Z}_j$ . Operation  $\bigoplus$  here means the direct sum of vector subspaces, i.e. if  $U = V \bigoplus W$  then for each  $u \in U$  there exists a unique decomposition  $u = v + w$ , with  $v \in V$  and  $w \in W$ .

Relatively compact subsets (whose closure are compacts) of separable Banach spaces with Schauder decomposition are then characterized by the following theorem.

**Theorem 2 (Suquet, [39])** *Let  $B$  be a separable Banach space having a Schauder decomposition  $(\mathcal{B}_j, j \in \mathbb{N})$ . A subset  $K$  is relatively compact in  $B$  if and only if:*

- i) for each  $j \in \mathbb{N}$ ,  $E_j K$  is relatively compact in  $\mathcal{Z}_j := \bigoplus_{i \leq j} \mathcal{B}_i$ ;
- ii)  $\sup_{x \in K} \|x - E_j x\| \rightarrow 0$  as  $j \rightarrow \infty$ .

## 1.2 Hölder space and its properties

The functional framework for our study of convergence of random fields is a certain class of Hölder spaces whose definition and some useful properties are gathered in this section.

### 1.2.1 Definition

For  $0 < \alpha < 1$ , define the Hölder space  $\mathbb{H}_\alpha^o([0, 1]^d)$  as the vector space of functions  $x : [0, 1]^d \rightarrow \mathbb{H}$  such that

$$\|x\|_\alpha := \|x(0)\| + w_\alpha(x, 1) < \infty,$$

with

$$w_\alpha(x, \delta) := \sup_{0 < |\mathbf{t} - \mathbf{s}| \leq \delta} \frac{\|x(\mathbf{t}) - x(\mathbf{s})\|}{|\mathbf{t} - \mathbf{s}|^\alpha} \xrightarrow{\delta \rightarrow 0} 0.$$

Endowed with the norm  $\|\cdot\|_\alpha$ ,  $\mathbb{H}_\alpha^o([0, 1]^d)$  is a separable Banach space. In the special case  $\mathbb{H} = \mathbb{R}$ , we write  $H_\alpha^o([0, 1]^d)$  instead of  $\mathbb{H}_\alpha^o([0, 1]^d)$ . For other Hilbert spaces  $H$  we write  $H_\alpha^o([0, 1]^d, H)$ .

### 1.2.2 Pyramidal functions

From works of Račkauskas and Suquet [27, 33] we know that the space  $\mathbb{H}_\alpha^o([0, 1]^d)$  has a Schauder decomposition. The linear subspaces featuring

in the decomposition contain linear combinations of certain pyramidal functions. We give now the precise definitions.

For convenience we write  $T = [0, 1]^d$  in this section. If  $A$  is a convex subset of  $T$ , the function  $f : T \rightarrow \mathbb{H}$  is said to be affine on  $A$  if it preserves the barycenter, i.e. for any finite sequence  $\mathbf{u}_1, \dots, \mathbf{u}_m$  in  $A$  and non negative scalars  $r_1, \dots, r_m$  such that  $\sum_{i=1}^m r_i = 1$ ,  $f(\sum_{i=1}^m r_i \mathbf{u}_i) = \sum_{i=1}^m r_i f(\mathbf{u}_i)$ .

Let us define the *standard triangulation* of the unit cube  $T = [0, 1]^d$ . Write  $\Pi_d$  for the set of permutations of the indexes  $1, \dots, d$ . For any  $\pi = (i_1, \dots, i_d) \in \Pi_d$ , let  $\Delta_\pi(T)$  be the convex hull of the  $d + 1$  points

$$\mathbf{0}, \mathbf{e}_{i_1}, (\mathbf{e}_{i_1} + \mathbf{e}_{i_2}), \dots, \sum_{k=1}^d \mathbf{e}_{i_k},$$

where the  $\mathbf{e}_i$ 's are the vectors of the canonical basis of  $\mathbb{R}^d$ . So, each simplex  $\Delta_\pi(T)$  corresponds to one path from  $\mathbf{0}$  to  $(1, \dots, 1)$  via vertices of  $T$  and such that along each segment of the path, only one coordinate increases while the others remain constants. Thus  $T$  is divided into  $d!$  simplexes with disjoint interiors. The standard triangulation of  $T$  is the family  $T_0$  of simplexes  $\{\Delta_\pi(T), \pi \in \Pi_d\}$ .

Next, we divide  $T$  into  $2^{jd}$  dyadic cubes with edge  $2^{-j}$ . By dyadic translations and change of scale, each of them is equipped with a triangulation similar to  $T_0$ . And  $T_j$  is the set of the  $2^{jd}d!$  simplexes so constructed. For  $j \geq 1$  the set  $W_j := \text{vert}(T_j)$  of vertices of the simplexes in  $T_j$  is

$$W_j = \{k2^{-j}; 0 \leq k \leq 2^j\}^d.$$

In what follows we put  $V_0 := W_0$  and  $V_j := W_j \setminus W_{j-1}$ . So  $V_j$  is the set of new vertices born with the triangulation  $T_j$ . More explicitly,  $V_j$  is the set of dyadic points  $\mathbf{v} = (k_1 2^{-j}, \dots, k_d 2^{-j})$  in  $W_j$  with at least one  $k_i$  odd.

The  $T_j$ -pyramidal function  $\Lambda_{j,\mathbf{v}}$  with peak vertex  $\mathbf{v} \in V_j$  is the *real valued* function defined on  $T$  by three conditions

- i.  $\Lambda_{j,\mathbf{v}}(\mathbf{v}) = 1$ ;
- ii.  $\Lambda_{j,\mathbf{v}}(\mathbf{w}) = 0$  if  $\mathbf{w} \in \text{vert}(T_j)$  and  $\mathbf{w} \neq \mathbf{v}$ ;
- iii.  $\Lambda_{j,\mathbf{v}}$  is affine on each simplex  $\Delta$  in  $T_j$ .

It follows clearly due to affinity from above definition that the support of  $\Lambda_{j,\mathbf{v}}$  is the union of all simplexes in  $T_j$  containing the peak vertex  $\mathbf{v}$ . By [37](Prop. 3.4.5) the functions  $\Lambda_{j,\mathbf{v}}$  are obtained by dyadic translations and changes of scale:

$$\Lambda_{j,\mathbf{v}}(\mathbf{t}) = \Lambda(2^j(\mathbf{t} - \mathbf{v})), \quad \mathbf{t} \in T,$$

from the same function  $\Lambda$ :

$$\Lambda(\mathbf{t}) := \max\left(0, 1 - \max_{t_i < 0} |t_i| - \max_{t_i > 0} t_i\right), \quad \mathbf{t} = (t_1, \dots, t_d) \in [-1, 1]^d.$$

Since  $\Lambda_{j,v}$  are affine on each simplex  $\Delta$  in  $T_j$ , it is clear that  $\Lambda_{j,v} \in \mathbb{H}_\alpha^o([0, 1]^d)$  for  $0 < \alpha < 1$ . Thus linear combinations  $\sum_{v \in V_j} h_v \Lambda_{j,v}$  with  $h_v \in \mathbb{H}$  are elements from  $\mathbb{H}_\alpha^o([0, 1]^d)$ . For each  $j$  such sums are continuous on  $T$ , affine on each simplex  $\Delta$  of  $T_j$  and vanishing at the vertices of  $W_{j-1}$ . Račkauskas and Suquet[33] prove that subspaces containing such functions form Schauder decomposition of  $\mathbb{H}_\alpha^o([0, 1]^d)$ .

**Theorem 3 (Račkauskas and Suquet [33])** *The space  $\mathbb{H}_\alpha^o([0, 1]^d)$  has the Schauder decomposition*

$$\mathbb{H}_\alpha^o([0, 1]^d) = \bigoplus_{i=0}^{\infty} \mathbf{V}_i,$$

where  $\mathbf{V}_0$  is the space of  $\mathbb{H}$ -valued functions continuous on  $T$  and affine on each simplex  $\Delta$  of  $T_0$  and for  $i \geq 1$ ,  $\mathbf{V}_i$  is the space of  $\mathbb{H}$ -valued functions continuous on  $T$ , affine on each simplex  $\Delta$  of  $T_i$  and vanishing at the vertices of  $W_{i-1}$ .

Each element  $x \in \mathbb{H}_\alpha^o([0, 1]^d)$  then has unique representation

$$x = \sum_{i=0}^{\infty} \sum_{v \in V_i} \lambda_{i,v}(x) \Lambda_{i,v},$$

with the  $\mathbb{H}$ -valued coefficients  $\lambda_{j,v}(x)$  defined as

$$\begin{aligned} \lambda_{0,v}(x) &= x(\mathbf{v}), \quad \mathbf{v} \in V_0; \\ \lambda_{j,v}(x) &= x(\mathbf{v}) - \frac{1}{2} \left( x(\mathbf{v}^-) + x(\mathbf{v}^+) \right), \quad \mathbf{v} \in V_j, \quad j \geq 1, \end{aligned}$$

where  $\mathbf{v}^-$  and  $\mathbf{v}^+$  are defined as follows. Each  $\mathbf{v} \in V_j$  admits a unique representation  $\mathbf{v} = (v_1, \dots, v_d)$  with  $v_i = k_i/2^j$ , ( $1 \leq i \leq d$ ). The points  $\mathbf{v}^- = (v_1^-, \dots, v_d^-)$  and  $\mathbf{v}^+ = (v_1^+, \dots, v_d^+)$  are defined by

$$v_i^- = \begin{cases} v_i - 2^{-j}, & \text{if } k_i \text{ is odd;} \\ v_i, & \text{if } k_i \text{ is even} \end{cases} \quad v_i^+ = \begin{cases} v_i + 2^{-j}, & \text{if } k_i \text{ is odd;} \\ v_i, & \text{if } k_i \text{ is even.} \end{cases}$$

Having specified Schauder decomposition of the space  $\mathbb{H}_\alpha^o([0, 1]^d)$  we adapt theorem 2 specifically for space  $\mathbb{H}_\alpha^o([0, 1]^d)$ . Define  $\mathbf{W}_j = \bigoplus_{i=0}^j \mathbf{V}_i$ . Note that

$\mathbf{W}_j$  corresponds to  $\mathcal{Z}_j$  in theorem 2. Define the projection operators  $E_j$  ( $j \geq 0$ ) onto subspaces  $\mathbf{W}_j$  by

$$E_j x := \sum_{i=0}^j \sum_{v \in V_i} \lambda_{i,v}(x) \Lambda_{i,v}, \quad x \in \mathbb{H}_\alpha^o([0, 1]^d). \quad (1.3)$$

Note that  $E_j$  is actually the operator of affine interpolation at the vertices of  $W_j$ , i.e. value of  $E_j x$  depends only on values of  $x(w)$  for  $w \in W_j$ . Since  $W_j$  is a finite set,  $\mathbf{W}_j$  is clearly isomorphic to the Cartesian product of  $\text{card}(W_j)$  copies of  $\mathbb{H}$ , where  $\text{card}(W_j)$  is the number of elements in  $W_j$ . We exploit this fact later in proving tightness.

Having defined operators  $E_j$ , we give now some alternative representation of  $\|x - E_j x\|_\alpha$ . For any function  $x \in H_\alpha(\mathbb{H})$ , define its sequential seminorm by

$$\|x\|_\alpha^{\text{seq}} := \sup_{j \geq 0} 2^{\alpha j} \max_{v \in V_j} \|\lambda_{j,v}(x)\|.$$

Račkauskas and Suquet [33] show that this seminorm is actually a norm and that it is equivalent to the norm  $\|x\|_\alpha$  on  $\mathbb{H}_\alpha^o([0, 1]^d)$ . Furthermore

$$\|x - E_J x\|_\alpha = \|x - E_J x\|_\alpha^{\text{seq}} := \sup_{j > J} 2^{\alpha j} \max_{v \in V_j} \|\lambda_{j,v}(x)\|, \quad (1.4)$$

is non increasing in  $J$ . Now we can state the adaptation of theorem 2 for the space  $\mathbb{H}_\alpha^o([0, 1]^d)$ .

**Theorem 4** *A subset  $K$  is relatively compact in  $\mathbb{H}_\alpha^o([0, 1]^d)$  if and only if:*

- i. *for each  $j \in \mathbb{N}$ ,  $E_j K$  is relatively compact in  $\mathbf{W}_j$ ;*
- ii.  *$\sup_{x \in K} \sup_{j > J} 2^{\alpha j} \max_{v \in V_j} \|\lambda_{j,v}(x)\| \rightarrow 0$ , as  $J \rightarrow \infty$ .*

### 1.2.3 Tightness criteria

Now we prove the tightness criteria. Note that this is an adaptation of the theorem 2 in [33] for nets  $\{\zeta_n, n \in \mathbb{N}^d\}$ .

**Theorem 5** *Let  $\{\zeta_n, n \in \mathbb{N}^d\}$  and  $\zeta$  be random elements with values in the space  $\mathbb{H}_\alpha^o([0, 1]^d)$ . Assume that the following conditions are satisfied.*

- i) *For each dyadic  $t \in [0, 1]^d$ , the net of  $\mathbb{H}$ -valued random elements  $\zeta_n(t)$  is asymptotically tight on  $\mathbb{H}$ .*

ii) For each  $\varepsilon > 0$

$$\lim_{J \rightarrow \infty} \limsup_{m(\mathbf{n}) \rightarrow \infty} P(\sup_{j \geq J} 2^{\alpha j} \max_{v \in V_j} \|\lambda_{j,v}(\zeta_{\mathbf{n}})\| > \varepsilon) = 0.$$

Then the net  $\zeta_{\mathbf{n}}$  is asymptotically tight in the space  $\mathbb{H}_{\alpha}^o([0, 1]^d)$ .

*Proof.*

For fixed positive  $\eta$ , put  $\eta_l = 2^{-l}$ ,  $l = 1, 2, \dots$  and choose a sequence  $(\varepsilon_l)$  decreasing to zero. By (ii) there is an integer  $J_l$  and index  $\mathbf{n}_0 \in \mathbb{N}^d$  such that for set

$$A_l := \{x : \sup_{j \geq J_l} 2^{\alpha j} \max_{v \in V_j} \|\lambda_{j,v}(x)\| < \varepsilon_l\}$$

$P(\zeta_{\mathbf{n}} \in A_l) > 1 - \eta_l$ , for all  $\mathbf{n}$ ,  $\mathbf{n}_0 \leq \mathbf{n}$ .

Recall now from subsection 1.2.2 that  $\mathbf{W}_j$  is isomorphic to the Cartesian product of  $\text{card}(W_j)$  copies of  $\mathbb{H}$ . Thus from (i) there exists for all  $\mathbf{n}$ ,  $\mathbf{n}_0 \leq \mathbf{n}$  a compact  $K_l \subset \mathbb{H}_{\alpha}^o([0, 1]^d)$ , such that for set  $B_l$

$$B_l := \{x \in \mathbb{H}_{\alpha}^o([0, 1]^d) : E_{J_l}x \in K_l\}$$

$P(\zeta_{\mathbf{n}} \in B_l) > 1 - \eta_l$ . Take  $K$  the closure of  $\cap_{l=1}^{\infty} (A_l \cap B_l)$ . Then  $P(K) > 1 - 2\eta$ , and  $K$  is compact due to theorem 4.

Since in  $\mathbb{R}$  closed bounded sets are compact and vice versa we have following corollary for space  $\mathbb{H}_{\alpha}^o([0, 1]^d)$

**Corollary 6** *Let  $\{\zeta_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}^d\}$  and  $\zeta$  be random elements with values in the space  $\mathbb{H}_{\alpha}^o([0, 1]^d)$ . Assume that the following conditions are satisfied.*

i)  $\lim_{a \rightarrow \infty} P(\sup_{t \in [0, 1]^d} |\zeta_{\mathbf{n}}| > a) = 0$

ii) For each  $\varepsilon > 0$

$$\lim_{J \rightarrow \infty} \limsup_{m(\mathbf{n}) \rightarrow \infty} P(\sup_{j \geq J} 2^{\alpha j} \max_{v \in V_j} |\lambda_{j,v}(\zeta_{\mathbf{n}})| > \varepsilon) = 0.$$

Then the net  $\zeta_{\mathbf{n}}$  is asymptotically tight in the space  $\mathbb{H}_{\alpha}^o([0, 1]^d)$ .

## 1.3 Results in probability

### 1.3.1 Gaussian processes

Limiting random fields considered in this work are mainly Gaussian ones. Recall that a real valued random field  $\{G(\mathbf{t}), \mathbf{t} \in [0, 1]^d\}$  is called Gaussian

if its finite-dimensional distributions are multivariate normal. Mean zero real valued Gaussian processes can be uniquely defined by their covariance function  $\mathbf{E} G(\mathbf{t})G(\mathbf{s})$ . The reverse problem is also of interest: when is a given function  $g : [0, 1]^d \times [0, 1]^d \rightarrow \mathbb{R}$  the covariance function of a certain Gaussian random field? The answer is a classical result which can be found in Khoshnevisan [18] for example. For convenience we state it here.

**Theorem 7** *If the function  $g : [0, 1]^d \times [0, 1]^d \rightarrow \mathbb{R}$  is symmetric and positive definite, i.e. for  $n$ -tuple of reals  $x_1, \dots, x_n$  and  $n$ -tuple of vectors  $\mathbf{t}_1, \dots, \mathbf{t}_n$  from  $[0, 1]^d$ , expression  $\sum_{i,j=1}^n x_i g(\mathbf{t}_i, \mathbf{t}_j) x_j \geq 0$ , then there exists a zero mean Gaussian random field  $\{G(\mathbf{t}), \mathbf{t} \in [0, 1]^d\}$  with covariance function  $\mathbf{E} G(\mathbf{t})G(\mathbf{s}) = g(\mathbf{t}, \mathbf{s})$ .*

For  $d = 1$  and  $g(s, t) = s \wedge t$ , we get a Brownian motion. Its generalization for  $d > 1$  is called Brownian sheet, a zero mean Gaussian process with covariance function  $g(\mathbf{s}, \mathbf{t}) = (s_1 \wedge t_1) \dots (s_d \wedge t_d)$ . As Brownian motion is usual limiting process in invariance principles for one parameter summation processes, Brownian sheet is limiting process for multiparameter summation processes.

We now define Hilbert space valued Brownian sheet. Recall that zero mean Gaussian random variables in Hilbert space are uniquely defined through their covariance operator. Covariance operator of  $\mathbb{H}$ -valued random variable  $X$  is linear operator  $\Gamma : \mathbb{H} \rightarrow \mathbb{H}$  satisfying

$$\langle \Gamma x, y \rangle = \mathbf{E} \langle X, x \rangle \langle X, y \rangle, \quad x, y \in \mathbb{H}.$$

Linear operator  $\Gamma$  is the covariance operator of some  $\mathbb{H}$ -valued random variable if it is

1. Symmetric:  $\langle \Gamma x, y \rangle = \langle \Gamma y, x \rangle$  for all  $x, y \in \mathbb{H}$ .
2. Positive:  $\langle \Gamma x, x \rangle \geq 0$ , for all  $x \in \mathbb{H}$ .
3. Nuclear: operator  $\Gamma$  is compact and for every orthonormal base  $\{e_n\} \subset \mathbb{H}$

$$\sum_n \langle \Gamma e_n, e_n \rangle < \infty.$$

Note that if  $\mathbf{E} \|X\|^2 < \infty$  we have

$$\sum_n \langle \Gamma e_n, e_n \rangle = \sum_n \mathbf{E} \langle X, e_n \rangle^2 = \mathbf{E} \|X\|^2 < \infty \quad (1.5)$$

It is well known that in the Hilbert space  $\mathbb{H}$ , every random element  $X$  such that  $\mathbf{E} \|X\|^2 < \infty$  is *pregaussian*, i.e. there is a Gaussian random element  $G$  in  $\mathbb{H}$  with the same covariance operator as  $X$ , see [22, Prop. 9.24]. Let the  $X_i$ 's be i.i.d. copies of  $X$ . If moreover  $\mathbf{E} X = 0$ , then  $n^{-1/2} \sum_{i=1}^n X_i$  converges weakly to  $G$  in  $\mathbb{H}$ , in other words  $X$  satisfies the CLT in  $\mathbb{H}$  [22, Th. 10.5].

Existence of real valued Gaussian processes is given by Kolmogorov theorem. Since it applies also for Cartesian products of Polish spaces it is natural to call  $\mathbb{H}$ -valued random field  $\{G(\mathbf{t}), \mathbf{t} \in [0, 1]^d\}$  Gaussian if for every  $m$ -tuple  $\mathbf{t}_1, \dots, \mathbf{t}_m$  vector  $(G(\mathbf{t}_1), \dots, G(\mathbf{t}_m))$  is  $\mathbb{H}^m$  valued Gaussian random variable.

Define then  $\mathbb{H}$ -valued Brownian sheet with covariance operator  $\Gamma$  as a  $\mathbb{H}$ -valued zero mean Gaussian process indexed by  $[0, 1]^d$  and satisfying

$$\mathbf{E} \langle W(\mathbf{t}), x \rangle \langle W(\mathbf{s}), y \rangle = (t_1 \wedge s_1) \dots (t_d \wedge s_d) \langle \Gamma x, y \rangle \quad (1.6)$$

for  $\mathbf{t}, \mathbf{s} \in [0, 1]^d$  and  $x, y \in \mathbb{H}$ . To check that this definition is valid note at first that for each  $\mathbf{t} \in [0, 1]^d$ ,  $W(\mathbf{t})$  is  $\mathbb{H}$ -valued zero mean Gaussian random variable with covariance operator  $\pi(\mathbf{t})\Gamma$ . Denote by  $\langle \cdot, \cdot \rangle_m$  the scalar product in  $\mathbb{H}^m$  which is defined by

$$\langle h, g \rangle_m := \sum_{i=1}^m \langle h_i, g_i \rangle, \quad h = (h_1, \dots, h_m), \quad g = (g_1, \dots, g_m) \in \mathbb{H}^m.$$

Denote by  $\Gamma_{\mathbf{t}_1, \dots, \mathbf{t}_m}$  the covariance operator of  $(W(\mathbf{t}_1), \dots, W(\mathbf{t}_m))$ . For  $x = (x_1, \dots, x_m) \in \mathbb{H}$  and  $y = (y_1, \dots, y_m) \in \mathbb{H}$  from (1.6) we get

$$\begin{aligned} \langle \Gamma_{\mathbf{t}_1, \dots, \mathbf{t}_m} x, y \rangle_m &= \mathbf{E} \sum_{i=1}^m \langle W(\mathbf{t}_i), x_i \rangle \sum_{j=1}^m \langle W(\mathbf{t}_j), y_j \rangle \\ &= \sum_i \sum_j g(\mathbf{t}_i, \mathbf{t}_j) \langle \Gamma x_i, y_j \rangle \end{aligned}$$

with  $g(\mathbf{t}_i, \mathbf{t}_j) = \prod_{k=1}^d t_{ik} \wedge t_{jk}$ . Since  $\Gamma$  is symmetric we get that  $\Gamma_{\mathbf{t}_1, \dots, \mathbf{t}_m}$  is symmetric also. Denote by  $X$ , a  $\mathbb{H}$ -valued random variable with covariance operator  $\Gamma$ . Then for  $x = (x_1, \dots, x_m) \in \mathbb{H}$

$$\begin{aligned} \langle \Gamma_{\mathbf{t}_1, \dots, \mathbf{t}_m} x, x \rangle_m &= \sum_i \sum_j g(\mathbf{t}_i, \mathbf{t}_j) \langle \Gamma x_i, x_j \rangle \\ &= \sum_i \sum_j g(\mathbf{t}_i, \mathbf{t}_j) \mathbf{E} \langle X, x_i \rangle \langle X, x_j \rangle \\ &= \mathbf{E} \left( \sum_i \sum_j \langle X, x_i \rangle g(\mathbf{t}_i, \mathbf{t}_j) \langle X, x_j \rangle \right) \geq 0. \end{aligned}$$

Thus  $\Gamma_{\mathbf{t}_1, \dots, \mathbf{t}_m}$  is positive. Now since

$$\mathbf{E} \|W(\mathbf{t}_1), \dots, W(\mathbf{t}_d)\|^2 = \sum_{i=1}^m \mathbf{E} \|W(\mathbf{t}_i)\|^2 < \infty$$

(1.5) implies that  $\Gamma_{\mathbf{t}_1, \dots, \mathbf{t}_m}$  is nuclear. Thus there exists  $\mathbb{H}^m$ -valued Gaussian random variable with covariance function  $\Gamma_{\mathbf{t}_1, \dots, \mathbf{t}_m}$  and our definition of  $\mathbb{H}$ -valued Brownian sheet is valid.

It is well known that trajectories of the real valued Brownian sheet are in  $H_\alpha^o([0, 1]^d)$  for  $0 < \alpha < 1/2$ . As the following estimate

$$\mathbf{E} \|W(\mathbf{t} + \mathbf{h}) + W(\mathbf{t} - \mathbf{h}) - 2W(\mathbf{t})\|^2 \leq c|\mathbf{h}| \operatorname{tr} \Gamma,$$

is valid for all  $\mathbf{t} - \mathbf{h}, \mathbf{t}, \mathbf{t} + \mathbf{h} \in [0, 1]^d$ , it follows from Račkauskas and Suquet [27] that  $W(\mathbf{t})$  has a version in  $\mathbb{H}_\alpha^o([0, 1]^d)$  for any  $0 < \alpha < 1/2$ .

### 1.3.2 A variant of continuous mapping theorem

Classical continuous mapping theorem states that if the sequence (or net) of random elements  $X_\alpha$  converges weakly to  $X$ , then for any continuous functional  $g$ , real random variable  $g(X_\alpha)$  converges weakly to  $g(X)$ . This result is widely applied in statistics to obtain limiting distributions. Sometimes though it is too restrictive, since sometimes it is more convenient to use the converging sequence (or net) of continuous functionals. Recall that the net of continuous functionals  $g_\alpha : B \rightarrow R$  where  $(B, \|\cdot\|)$  is a normed Banach space is called equicontinuous if for every  $\varepsilon > 0$  and any  $x, y \in B$  such that  $\|x - y\| < \delta$  we have

$$\sup_\alpha |g_\alpha(x) - g_\alpha(y)| < \varepsilon.$$

Then following theorem holds (it is stated as lemma in [32], we restate it for the case of nets).

**Lemma 8** *Let  $\{\eta_\alpha\}$  be a asymptotically tight net of random elements in the separable Banach space  $B$ , and  $g_\alpha, g$  be a continuous functionals  $B \rightarrow R$ . Assume that  $g_\alpha$  converges pointwise to  $g$  on  $B$ , and that  $g_\alpha$  is equicontinuous. Then*

$$g_\alpha(\eta_\alpha) = g(\eta_\alpha) + o_P(1).$$

*Proof.* By the asymptotic tightness assumption there is for every  $\varepsilon > 0$  a compact subset  $K$  in  $B$  and  $\alpha_0$  such that for every  $\alpha_0 \leq \alpha$ ,  $P(\eta_\alpha \notin K) < \varepsilon$ .

Now by a Arzela-Ascoli theorem the net  $g_\alpha$  is totally bounded on the compact  $K$  with respect to norm of uniform convergence. Since  $g_\alpha$  converges pointwise to  $g$ , this gives us uniform convergence of  $g_\alpha$  to  $g$  on  $K$ . Then for every  $\delta > 0$  there is some  $\alpha_1$  depending on  $\delta$  and  $K$ , such that

$$\sup_{x \in K} |g_\alpha(x) - g(x)| \leq \delta, \quad \alpha_1 \leq \alpha.$$

Take now  $\alpha_2$  such that  $\alpha_1 \leq \alpha_2$  and  $\alpha_0 \leq \alpha_2$ . Then for  $\alpha_2 \leq \alpha$  we have

$$P(|g_\alpha(\eta_\alpha) - g(\eta_\alpha)| \geq \delta) \leq P(\eta_\alpha \notin K) < \varepsilon,$$

which gives us the proof.

The following lemma from [32] provides some practical sufficient conditions to check the equicontinuity of some families of functionals.

**Lemma 9** *Let  $(B, \|\cdot\|)$  be a vector normed space and  $q : B \rightarrow \mathbb{R}$  such that*

(a).  *$q$  is subadditive:  $q(x + y) \leq q(x) + q(y)$ ,  $x, y \in B$ .*

(b).  *$q$  is symmetric:  $q(x) = q(-x)$ ,  $x \in B$ .*

(c). *For some constant  $C$ ,  $q(x) \leq C\|x\|$ ,  $x \in B$ .*

*Then  $q$  satisfies the Lipschitz condition*

$$|q(x + y) - q(x)| \leq C\|y\|, \quad x, y \in B \quad (1.7)$$

*If  $\mathcal{F}$  is any set of functionals  $q$  fulfilling (a), (b), (c) with the same constant  $C$ , then (a), (b), (c) are inherited by  $g(x) := \sup\{q(x), q \in \mathcal{F}\}$  which therefore satisfies (1.7).*

### 1.3.3 Rosenthal inequality

Since the Hilbert space  $\mathbb{H}$  has cotype 2, it satisfies the following vector valued version of Rosenthal's inequality for every  $q \geq 2$ , see [21, Th. 2.6]. For any finite set  $(Y_i)_{i \in I}$  of independent random elements in  $\mathbb{H}$  with zero mean and such that  $\mathbf{E} \|Y_i\|^q < \infty$  for every  $i \in I$ ,

$$\mathbf{E} \left\| \sum_{i \in I} Y_i \right\|^q \leq C'_q \left( \mathbf{E} \left\| \sum_{i \in I} G(Y_i) \right\|^q + \sum_{i \in I} \mathbf{E} \|Y_i\|^q \right), \quad (1.8)$$

where the constant  $C'_q$  depends only on  $q$  and the  $G(Y_i)$  are centered Gaussian independent random elements in  $\mathbb{H}$  such that for every  $i \in I$ ,  $G(Y_i)$  has

the same covariance structure as  $Y_i$ . For real random variables Rosenthal inequality simply reads

$$\mathbf{E} \left| \sum_{i \in I} Y_i \right|^q \leq c \left( \left( \sum_{i \in I} \sigma_i^2 \right)^{q/2} + \sum_{i \in I} \mathbf{E} |Y_i|^q \right), \quad (1.9)$$

where  $\sigma_i^2 = \mathbf{E} Y_i^2$ .

In the i.i.d. case with  $N = \text{card}(I)$ , we note that  $\sum_{i \in I} G(Y_i)$  is Gaussian with the same distribution as  $N^{1/2}G(Y_1)$  and using the equivalence of moments for Gaussian random elements, see [22, Cor. 3.2], we obtain

$$\mathbf{E} \left\| \sum_{i \in I} Y_i \right\|^q \leq C_q'' \left( N^{q/2} \left( \mathbf{E} \|G(Y_1)\|^2 \right)^{q/2} + N \mathbf{E} \|Y_1\|^q \right),$$

where  $C_q''$  depends on  $q$  and does not depend on the distribution of  $Y_1$ . Since  $\mathbb{H}$  has also the type 2, there is a constant  $a$  depending only on  $\mathbb{H}$  such that  $\mathbf{E} \|G(Y_1)\|^2 \leq a \mathbf{E} \|Y_1\|^2$ , see [22, Prop. 9.24]. Finally there is a constant  $C_q$  depending on  $\mathbb{H}$ ,  $q$ , but not on the distribution of the  $Y_i$ 's, such that

$$\mathbf{E} \left\| \sum_{i \in I} Y_i \right\|^q \leq C_q \left( N^{q/2} \left( \mathbf{E} \|Y_1\|^2 \right)^{q/2} + N \mathbf{E} \|Y_1\|^q \right), \quad (N = \text{card}(I)). \quad (1.10)$$

### 1.3.4 Doob inequality

We shall need a generalization of maximal Doob inequality for multiparameter martingales. We use definitions and results from Khoshnevisan [18].

**Definition 2** *Let  $d \in \mathbb{N}$  and consider  $d$  (one parameter) filtrations  $\mathcal{F}^1, \dots, \mathcal{F}^d$ , where  $\mathcal{F}^i = \{\mathcal{F}_k^i, k \geq 1\}$  ( $1 \leq i \leq d$ ). A stochastic process  $M = (M_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}^d)$  is an orthosubmartingale if for each  $1 \leq i \leq d$  ( $M_{\mathbf{n}}, n_i \geq 1$ ) is a one parameter submartingale with respect to the one parameter filtration  $\mathcal{F}^i$  with other coordinates  $n_j \neq n_i$  fixed.*

The classical example of orthosubmartingale is the multiparameter random walk  $\{S_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}^d\}$  defined as

$$S_{\mathbf{n}} = \sum_{1 \leq j \leq n} X_j,$$

where  $\{X_j, \mathbf{j} \in \mathbb{N}^d\}$  is the collection of mean-zero random variables. The process  $\{S_n, \mathbf{n} \in \mathbb{N}^d\}$  is an orthosubmartingale with respect to filtrations

$$\mathcal{F}_k^i = \sigma(X_j, j_i \leq k)$$

where  $\sigma(\dots)$  denotes the  $\sigma$ -field generated by the random variables in parenthesis.

For nonnegative orthosubmartingales the following so called Cairoli's strong  $(p, p)$  inequality holds.

**Theorem 10 (Th. 2.3.1 in Khoshnevisan [18])** *Suppose that  $M = (M_n, \mathbf{n} \in \mathbb{N}^d)$  is a nonnegative orthosubmartingale with respect to one-parameter filtrations  $\mathcal{F}^1, \dots, \mathcal{F}^d$ . Then for all  $\mathbf{n} \in \mathbb{N}^d$  and  $p > 1$*

$$\mathbf{E} \left[ \max_{0 \leq k \leq \mathbf{n}} M_k^p \right] \leq \left( \frac{p}{p-1} \right)^{dp} \mathbf{E} [M_n^p].$$

Following lemma is useful

**Theorem 11 (Lemma 2.1.1 in Khoshnevisan [18])** *Suppose that  $M = (M_n, \mathbf{n} \in \mathbb{N}^d)$  is a nonnegative orthosubmartingale with respect to one-parameter filtrations  $\mathcal{F}^1, \dots, \mathcal{F}^d$ , that  $\Psi : [0, \infty) \rightarrow [0, \infty)$  is convex nondecreasing, and that for all  $\mathbf{n} \in \mathbb{N}^d$ ,  $\mathbf{E} \Psi(M_n) < \infty$ . Then  $(\Psi(M_n), \mathbf{n} \in \mathbb{N}^d)$  is an orthosubmartingale.*

For independent zero mean real random variables  $\{X_j, \mathbf{j} \in \mathbb{N}^d\}$  introduce one parameter filtrations  $\mathcal{F}^i = \mathcal{F}_k^i = \sigma(X_j, j_i \leq k)$ . Then process  $S_n = \sum_{\mathbf{j} \leq \mathbf{n}} X_j$  is orthosubmartingale with respect to filtrations  $\mathcal{F}^i$  and process  $|S_n|$  is nonnegative orthosubmartingale with respect to the same filtrations. Thus we have

$$\mathbf{E} \max_{1 \leq j \leq \mathbf{n}} |S_j|^p \leq \left( \frac{p}{p-1} \right)^{dp} \mathbf{E} |S_n|^p. \quad (1.11)$$

For i.i.d. Hilbert space valued random field  $\{X_j, \mathbf{j} \in \mathbb{N}^d\}$  introduce  $d$  one parameter filtrations,  $\mathcal{F}^i = (\mathcal{F}_k^i, k = 0, 1, \dots)$ ,  $i = 1, \dots, d$ , where  $\mathcal{F}_k^i = \sigma(X_j, \mathbf{j} \in \mathbb{N}^d, j_i \leq k)$ . Since Cairoli's inequality applies for real valued orthosubmartingales we cannot use it directly for  $S_n = \sum_{\mathbf{j} \leq \mathbf{n}} X_j$ , since  $S_n$  is  $\mathbb{H}$  valued stochastic process. On the other hand stochastic process  $(\|S_n\|, \mathbf{n} \in \mathbb{N}^d)$  is real valued so to apply theorem 10 we must show that  $\|S_n\|$  is orthosubmartingale with respect to filtrations  $\mathcal{F}^1, \dots, \mathcal{F}^d$ . Since norm is a continuous functional, the map  $n_i \rightarrow \|S_n\|$  is  $\mathcal{F}_{n_i}^i$ -measurable for each  $i = 1, \dots, d$ .

Assume that  $\mathbf{E} \|X_1\| < \infty$ . Then the  $X_j$ 's are Bochner integrable and according to [40] we can introduce conditional expectations with respect to  $\mathcal{F}^i$ ,  $i = 1, \dots, d$ . Let  $\mathbf{E} X_j = 0$ . From properties of conditional expectation we have for  $i = 1, \dots, d$ ,  $\mathbf{n} \in \mathbb{N}^d$  and  $k = 0, 1, \dots$

$$\mathbf{E} (\|S_{\mathbf{n}}\| | \mathcal{F}_k^i) \geq \|\mathbf{E} (S_{\mathbf{n}} | \mathcal{F}_k^i)\| = \left\| \sum_{j \leq \mathbf{n}} \mathbf{E} (X_j | \mathcal{F}_k^i) \right\| = \|S_{(n_1, \dots, n_{i-1}, k, n_{i+1}, \dots, n_d)}\|.$$

Hence for each  $i = 1, \dots, d$ ,  $n_i \rightarrow S_{\mathbf{n}}$  is a one parameter submartingale with respect to the filtration  $\mathcal{F}^i$ . Applying then theorem 10 we have

$$\mathbf{E} \max_{1 \leq j \leq \mathbf{n}} \|S_j\|^p \leq \left( \frac{p}{p-1} \right)^{dp} \mathbf{E} \|S_{\mathbf{n}}\|^p, \quad (1.12)$$

for all  $\mathbf{n} \in \mathbb{N}^d$  and  $p > 1$ .

# Chapter 2

## Summation processes

We call random process a summation process if its values are defined only by the values of partial sums  $S_k = X_1 + \dots + X_k$ , where  $X_k$ ,  $k = 1, \dots, n$  are random variables. Usually summation process is defined using interpolation arguments. Classical example is polygonal line process indexed by  $[0, 1]$  with vertices  $(k/n, S_k)$ ,  $k = 0, 1, \dots, n$  and  $S_0 = 0$ . This process has continuous and Hölderian trajectories. Sometimes it is convenient to drop the requirement of continuity and to analyze jump process defined as  $\sum_{k=1}^{[nt]} X_k$ . In this section we define summation processes indexed by  $[0, 1]^d$  and give some useful representations. For reasons explained in section 2.2 we analyze separately summation processes based on random variables with the same variance and summation processes based on random variables with different variances. Though the results in this chapter are presented in a context of probability theory, they are derived without using any results from it. This chapter can be viewed as investigation of properties of certain interpolation schema of functions with domain  $[0, 1]^d$ . To improve readability, more technical and longer proofs are given at the end of each subsection.

### 2.1 Uniform variance case

#### 2.1.1 Differences of partial sums

In this and following chapters we deal a lot with differences of partial sums  $S_j = \sum_{1 \leq i \leq j} X_i$ . Let us introduce the notation

$$\Delta_k^{(i)} S_j = S_{(j_1, \dots, j_{i-1}, k, j_{i+1}, \dots, j_d)} - S_{(j_1, \dots, j_{i-1}, k-1, j_{i+1}, \dots, j_d)}$$

Since  $S_{\mathbf{k}}$  is really a function with domain  $\mathbb{N}^d$ , we can say that  $\Delta_{\mathbf{k}}^{(i)}$  is a difference operator acting on  $i$ -th coordinate of the argument of function  $S_{\mathbf{k}}$ . Note that superposition of operators  $\Delta_{\mathbf{k}}^{(i)}$  commute

$$\Delta_{j_i}^{(i)} \Delta_{j_l}^{(l)} S_j = \Delta_{j_l}^{(l)} \Delta_{j_i}^{(i)} S_j.$$

In particular for any  $\mathbf{j} \in \mathbb{N}^d$  we have

$$X_{\mathbf{j}} = \Delta_{j_1}^{(1)} \dots \Delta_{j_d}^{(d)} S_j.$$

### 2.1.2 Definitions and representations

For  $d = 1$  polygonal line and jump processes are given as

$$\xi_n(t) = S_{[nt]} + (nt - [nt])X_{[nt]+1}, \quad (2.1)$$

$$\zeta_n(t) = S_{[nt]}. \quad (2.2)$$

It is natural then to define  $[0, 1]^d$  indexed jump summation process as

$$\zeta_{\mathbf{n}}(\mathbf{t}) = S_{[\mathbf{nt}]}$$

It is not possible to do this for continuous version of  $[0, 1]^d$  indexed summation process, since the relation

$$S_{\mathbf{k}+\mathbf{1}} - S_{\mathbf{k}} = X_{\mathbf{k}+\mathbf{1}}$$

holds only for  $\mathbf{k} = \mathbf{0}$ . The continuous version of summation process for  $d > 1$  still can be defined using analogy. Note that for  $d = 1$  we can write

$$\xi_n(t) = \sum_{1 \leq i \leq n} n \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \cap [0, t] \right| X_i, \quad (2.3)$$

where  $|A|$  denotes Lebesgue measure of the set  $A$ . Define then continuous  $[0, 1]^d$  indexed summation process as

$$\xi_{\mathbf{n}}(\mathbf{t}) = \sum_{\mathbf{i} \leq \mathbf{n}} \pi(\mathbf{n}) \left| \left[ \frac{\mathbf{i}-\mathbf{1}}{\mathbf{n}}, \frac{\mathbf{i}}{\mathbf{n}} \right) \cap [0, \mathbf{t}] \right| X_{\mathbf{i}}. \quad (2.4)$$

where we write

$$[\mathbf{a}, \mathbf{b}] = [a_1, b_1] \times \dots \times [a_d, b_d] \quad (2.5)$$

for  $\mathbf{a}, \mathbf{b} \in [0, 1]^d$ . As in one dimensional case we can see that  $\xi_n(\mathbf{k}/\mathbf{n}) = S_{\mathbf{k}}$ , i. e. the process is a continuous interpolation of the grid  $(\mathbf{k}/\mathbf{n}, S_{\mathbf{k}})$ .

To prove tightness of summation process we must control the difference of the process when the distance between points is small. For case  $d = 1$  if  $s$  and  $t$  are close together, they fall into interval  $[i/n, (i+1)/n]$  for some  $i$ . In this interval we have

$$\begin{aligned}\xi_n(t) &= S_i + (nt - i)X_{i+1} \\ \xi_n(s) &= S_i + (ns - i)X_{i+1}\end{aligned}$$

thus

$$\xi_n(t) - \xi_n(s) = n(t - s)X_{i+1}. \quad (2.6)$$

Thus it is of interest to investigate properties of the summation process in rectangles  $[(i-1)/\mathbf{n}, i/\mathbf{n}]$ . For the case  $d = 1$  the summation process has the property that in interval  $[i/n, (i+1)/n]$  it is the affine interpolation of its values at interval endpoints:

$$\begin{aligned}\xi_n(t) &= S_i + (nt - i)X_{i+1} = S_i + (nt - i)(S_{i+1} - S_i) \\ &= (1 - (nt - i))S_i + (nt - i)S_{i+1} = (1 - \{nt\})S_{[nt]} + \{nt\}S_{[nt]+1} \\ &= (1 - \{nt\})\xi_n(i/n) + \{nt\}\xi_n((i+1)/n),\end{aligned}$$

with the weights coming from

$$nt = (1 - \{nt\})[nt] + \{nt\}([nt] + 1) = [nt] + \{nt\}. \quad (2.7)$$

The summation process  $\xi_n(\mathbf{t})$  retains this property. We show this directly for the case  $d = 2$  and then prove it for general case. Fix  $\mathbf{t} \in [0, 1]^2$  and choose  $\mathbf{i}$  so that  $\mathbf{t} \in [(\mathbf{i}-1)/\mathbf{n}, \mathbf{i}/\mathbf{n}]$ . Necessarily then  $[\mathbf{nt}] = \mathbf{i} - \mathbf{1}$ . In this case the expression  $\pi(\mathbf{n})|[(\mathbf{j}-1)/\mathbf{n}, \mathbf{j}/\mathbf{n}] \cap [0, \mathbf{t}]|$  assumes only following possible values:

$$\pi(\mathbf{n}) \left| \left[ \frac{\mathbf{j}-\mathbf{1}}{\mathbf{n}}, \frac{\mathbf{j}}{\mathbf{n}} \right) \cap [0, \mathbf{t}] \right| = \begin{cases} 1, & \text{for } \mathbf{j} \leq \mathbf{i} - \mathbf{1} \\ 0, & \text{for } \mathbf{j}, \text{ such that either } j_1 \geq i_1, \text{ or } j_2 \geq i_2 \\ \{n_1 t_1\}, & \text{for } \mathbf{j}, \text{ such that } j_1 = i_1 \text{ and } j_2 < i_2 \\ \{n_2 t_2\}, & \text{for } \mathbf{j}, \text{ such that } j_2 = i_2 \text{ and } j_1 < i_1 \\ \pi(\{\mathbf{nt}\}), & \text{for } \mathbf{j} = \mathbf{i}. \end{cases}$$

Thus

$$\xi_n(\mathbf{t}) = S_{i-1} + \{n_1 t_1\} \sum_{j=1}^{i_2-1} X_{i_1 j} + \{n_2 t_2\} \sum_{i=1}^{i_1-1} X_{i i_2} + \pi(\{\mathbf{n}\mathbf{t}\}) X_i. \quad (2.8)$$

We can rewrite this expression using difference operators as

$$\xi_n(\mathbf{t}) = S_{i-1} + \{n_1 t_1\} \Delta_{i_1}^{(1)} S_{i-1} + \{n_2 t_2\} \Delta_{i_2}^{(2)} S_{i-1} + \pi(\{\mathbf{n}\mathbf{t}\}) \Delta_{i_1}^{(1)} \Delta_{i_2}^{(2)} S_{i-1}, \quad (2.9)$$

or alternatively

$$\begin{aligned} \xi_n(\mathbf{t}) &= \pi(\mathbf{1} - \{\mathbf{n}\mathbf{t}\}) S_{i-1} + \pi(\{\mathbf{n}\mathbf{t}\}) S_i \\ &\quad + \{n_1 t_1\} (1 - \{n_2 t_2\}) S_{i_1, i_2-1} + (1 - \{n_1 t_1\}) \{n_2 t_2\} S_{i_1-1, i_2}. \end{aligned} \quad (2.10)$$

Note that by doing so we expressed  $\xi_n(\mathbf{t})$  as linear combination of its values at vertices of rectangle  $[(\mathbf{i} - \mathbf{1})/\mathbf{n}, \mathbf{i}/\mathbf{n}]$ . Furthermore the weights in this combination sum to one and

$$\begin{aligned} \mathbf{n}\mathbf{t} &= \pi(\mathbf{1} - \{\mathbf{n}\mathbf{t}\})[\mathbf{n}\mathbf{t}] + \pi(\{\mathbf{n}\mathbf{t}\})([\mathbf{n}\mathbf{t}] + \mathbf{1}) \\ &\quad + \{n_1 t_1\} (1 - \{n_2 t_2\}) ([n_1 t_1] + 1, [n_2 t_2]) \\ &\quad + (1 - \{n_1 t_1\}) \{n_2 t_2\} ([n_1 t_1], [n_2 t_2] + 1) \\ &= [\mathbf{n}\mathbf{t}] + \{\mathbf{n}\mathbf{t}\}. \end{aligned}$$

Thus in the point in the grid rectangle our summation process is weighted sum of its values on rectangle vertexes with the weights coming from barycentric decomposition of the point as it is in the case  $d = 1$ . Note that though we derived this decomposition for real valued random variables it holds for Banach space valued random variables also. We extend now (2.9) and (2.10) for general  $d$ .

**Proposition 12** For  $\mathbf{t} \in [0, 1)^d$ , denote  $\mathbf{s} = \{\mathbf{n}\mathbf{t}\}$  and represent vertices of the rectangle  $R_{\mathbf{n}, [\mathbf{n}\mathbf{t}] + \mathbf{1}}$  as

$$V(\mathbf{u}) := \frac{[\mathbf{n}\mathbf{t}]}{\mathbf{n}} + \frac{\mathbf{u}}{\mathbf{n}}, \quad \mathbf{u} \in \{0, 1\}^d. \quad (2.11)$$

It is possible to express  $\mathbf{t}$  as a barycenter of these  $2^d$  vertices with weights  $w(\mathbf{u}) \geq 0$  depending on  $\mathbf{t}$ , i.e.,

$$\mathbf{t} = \sum_{\mathbf{u} \in \{0, 1\}^d} w(\mathbf{u}) V(\mathbf{u}), \quad \text{with} \quad \sum_{\mathbf{u} \in \{0, 1\}^d} w(\mathbf{u}) = 1, \quad (2.12)$$

where

$$w(\mathbf{u}) = \prod_{l=1}^d s_l^{u_l} (1 - s_l)^{1-u_l}. \quad (2.13)$$

Using this representation, define the random field  $\xi_n^*$  by

$$\xi_n^*(\mathbf{t}) = \sum_{\mathbf{u} \in \{0,1\}^d} w(\mathbf{u}) S_{[\mathbf{n}\mathbf{t}]+\mathbf{u}}, \quad \mathbf{t} \in [0,1]^d.$$

Then  $\xi_n^*$  coincides with the summation process defined by (2.4), where  $\{X_i, \mathbf{1} \leq i \leq \mathbf{n}\}$  is a collection of Banach space valued random variables. Furthermore  $\xi_n$  admits representation

$$\xi_n(\mathbf{t}) = S_{[\mathbf{n}\mathbf{t}]} + \sum_{l=1}^d \sum_{1 \leq i_1 < i_2 < \dots < i_l \leq d} \left( \prod_{k=1}^l \{n_{i_k} t_{i_k}\} \right) \left( \prod_{k=1}^l \Delta_{[n_{i_k} t_{i_k}]+1}^{(i_k)} \right) S_{[\mathbf{n}\mathbf{t}]}. \quad (2.14)$$

### Proof of proposition 12

For notational convenience write

$$R_{\mathbf{n},\mathbf{j}} := \left[ \frac{\mathbf{j}-1}{\mathbf{n}}, \frac{\mathbf{j}}{\mathbf{n}} \right). \quad (2.15)$$

For fixed  $\mathbf{n} \geq \mathbf{1} \in \mathbb{N}^d$ , any  $\mathbf{t} \neq \mathbf{1} \in [0,1]^d$  belongs to a unique rectangle  $R_{\mathbf{n},\mathbf{j}}$ , defined by (2.15), namely  $R_{\mathbf{n},[\mathbf{n}\mathbf{t}]+\mathbf{1}}$ . Recalling definition  $\mathbf{s} = \{\mathbf{n}\mathbf{t}\}$ , note that

$$\mathbf{t} = \frac{[\mathbf{n}\mathbf{t}]}{\mathbf{n}} + \frac{\mathbf{s}}{\mathbf{n}}. \quad (2.16)$$

For any non empty subset  $L$  of  $\{1, \dots, d\}$ , we denote by  $\{0,1\}^L$  the set of binary vectors indexed by  $L$ . In particular  $\{0,1\}^d$  is an abridged notation for  $\{0,1\}^{\{1,\dots,d\}}$ . Now define the non negative weights

$$w_L(\mathbf{u}) := \prod_{l \in L} s_l^{u_l} (1 - s_l)^{1-u_l}, \quad \mathbf{u} \in \{0,1\}^L$$

and note that when  $L = \{1, \dots, d\}$  these weights coincide with weights  $w(\mathbf{u})$  defined in (2.13), hence we will not write subscript  $L$  in this case. For fixed  $L$ , the sum of all these weights is one since

$$\sum_{\mathbf{u} \in \{0,1\}^L} w_L(\mathbf{u}) = \prod_{l \in L} (s_l + (1 - s_l)) = 1. \quad (2.17)$$

The special case  $L = \{1, \dots, d\}$  gives the second equality in (2.12). From (2.17) we immediately deduce that for any  $K$  non empty and strictly included in  $\{1, \dots, d\}$ , with  $L := \{1, \dots, d\} \setminus K$ ,

$$\sum_{\substack{\mathbf{u} \in \{0,1\}^d, \\ \forall k \in K, u_k=1}} w(\mathbf{u}) = \prod_{k \in K} s_k \sum_{\mathbf{u} \in \{0,1\}^L} s_l^{u_l} (1 - s_l)^{1-u_l} = \prod_{k \in K} s_k. \quad (2.18)$$

Formula (2.18) remains obviously valid in the case where  $K = \{1, \dots, d\}$ .

Now let us observe that

$$\sum_{\mathbf{u} \in \{0,1\}^d} w(\mathbf{u}) V(\mathbf{u}) = \sum_{\mathbf{u} \in \{0,1\}^d} w(\mathbf{u}) \left( \frac{[\mathbf{n}\mathbf{t}]}{\mathbf{n}} + \frac{\mathbf{u}}{\mathbf{n}} \right) = \frac{[\mathbf{n}\mathbf{t}]}{\mathbf{n}} + \sum_{\mathbf{u} \in \{0,1\}^d} w(\mathbf{u}) \frac{\mathbf{u}}{\mathbf{n}}.$$

Comparing with the expression of  $\mathbf{t}$  given by (2.16), we see that the first equality in (2.12) will be established if we check that

$$\mathbf{s}' := \sum_{\mathbf{u} \in \{0,1\}^d} w(\mathbf{u}) \mathbf{u} = \mathbf{s}. \quad (2.19)$$

This is easily seen componentwise using (2.18) because for any fixed  $l \in \{1, \dots, d\}$ ,

$$s'_l = \sum_{\substack{\mathbf{u} \in \{0,1\}^d, \\ u_l=1}} w(\mathbf{u}) = \prod_{k \in \{l\}} s_k = s_l.$$

Next we check that  $\xi_n(\mathbf{t}) = \xi_n^*(\mathbf{t})$  for every  $\mathbf{t} \in [0, 1]^d$ . Introduce the notation

$$D_{\mathbf{t}, \mathbf{u}} := \mathbb{N}^d \cap \left( [0, [\mathbf{n}\mathbf{t}] + \mathbf{u}] \setminus [0, [\mathbf{n}\mathbf{t}]] \right).$$

Then we have

$$\begin{aligned} \xi_n^*(\mathbf{t}) &= \sum_{\mathbf{u} \in \{0,1\}^d} w(\mathbf{u}) \left( S_{[\mathbf{n}\mathbf{t}]} + (S_{[\mathbf{n}\mathbf{t}] + \mathbf{u}} - S_{[\mathbf{n}\mathbf{t}]}) \right) \\ &= S_{[\mathbf{n}\mathbf{t}]} \sum_{\mathbf{u} \in \{0,1\}^d} w(\mathbf{u}) \sum_{i \in D_{\mathbf{t}, \mathbf{u}}} X_i. \end{aligned}$$

Now in view of (2.4) the proof of  $\xi_n(\mathbf{t}) = \xi_n^*(\mathbf{t})$  reduces clearly to that of

$$\sum_{\mathbf{u} \in \{0,1\}^d} w(\mathbf{u}) \sum_{i \in D_{\mathbf{t}, \mathbf{u}}} X_i = \sum_{i \in I} \pi(\mathbf{n}) |R_{\mathbf{n}, i} \cap [0, \mathbf{t}]| X_i, \quad (2.20)$$

where

$$I := \{\mathbf{i} \leq \mathbf{n}; \forall k \in \{1, \dots, d\}, i_k \leq [n_k t_k] + 1 \text{ and} \\ \exists l \in \{1, \dots, d\}, i_l = [n_l t_l] + 1\}. \quad (2.21)$$

Clearly  $I$  is the union of all  $D_{\mathbf{t}, \mathbf{u}}$ ,  $\mathbf{u} \in \{0, 1\}^d$ , so we can rewrite the left hand side of (2.20) under the form  $\sum_{\mathbf{i} \in I} a_{\mathbf{i}} X_{\mathbf{i}}$ . For  $\mathbf{i} \in I$ , put

$$K(\mathbf{i}) := \{k \in \{1, \dots, d\}; i_k = [n_k t_k] + 1\}. \quad (2.22)$$

Then observe that for  $\mathbf{i} \in I$ , the  $\mathbf{u}$ 's such that  $\mathbf{i} \in D_{\mathbf{t}, \mathbf{u}}$  are exactly those which satisfy  $u_k = 1$  for every  $k \in K(\mathbf{i})$ . Using (2.18), this gives

$$\forall \mathbf{i} \in I, \quad a_{\mathbf{i}} = \sum_{\substack{\mathbf{u} \in \{0, 1\}^d, \\ \forall k \in K(\mathbf{i}), u_k = 1}} w(\mathbf{u}) = \prod_{k \in K(\mathbf{i})} s_k. \quad (2.23)$$

On the other hand we have for every  $\mathbf{i} \in I$ ,

$$|R_{\mathbf{n}, \mathbf{i}} \cap [0, \mathbf{t}]| = \prod_{k \in K(\mathbf{i})} \left( t_k - \frac{[n_k t_k]}{n_k} \right) \prod_{k \notin K(\mathbf{i})} \frac{1}{n_k} = \frac{1}{\boldsymbol{\pi}(\mathbf{n})} \prod_{k \in K(\mathbf{i})} s_k = \frac{a_{\mathbf{i}}}{\boldsymbol{\pi}(\mathbf{n})}. \quad (2.24)$$

Thus (2.20) follows. To prove (2.14) note that

$$\xi_{\mathbf{n}}(\mathbf{t}) = S_{[\mathbf{n}\mathbf{t}]} + \sum_{\mathbf{i} \in I} \boldsymbol{\pi}(\mathbf{n}) |R_{\mathbf{n}, \mathbf{i}} \cap [0, \mathbf{t}]| X_{\mathbf{i}} = S_{[\mathbf{n}\mathbf{t}]} + \sum_{\mathbf{i} \in I} \left( \prod_{k \in K(\mathbf{i})} s_k \right) X_{\mathbf{i}},$$

which can be recast as

$$\xi_{\mathbf{n}}(\mathbf{t}) = S_{[\mathbf{n}\mathbf{t}]} + \sum_{l=1}^d T_l(\mathbf{t}) \quad (2.25)$$

with

$$T_l(\mathbf{t}) := \sum_{\substack{\mathbf{i} \in I \\ \text{card}(K(\mathbf{i}))=l}} \left( \prod_{k \in K(\mathbf{i})} s_k \right) X_{\mathbf{i}}. \quad (2.26)$$

Let  $K \subset \{1, \dots, d\}$  and  $I_K = \{\mathbf{i} \in I; K(\mathbf{i}) = K\}$ . Then

$$I_K = \{\mathbf{i} \leq \mathbf{n}; i_k = [n_k t_k] + 1, \text{ for } k \in K \text{ and} \\ i_k \leq [n_k t_k], \text{ for } k \in \{1, \dots, d\} \setminus K\},$$

Take for example  $K = \{1\}$  and notice that

$$\sum_{\mathbf{i} \in I_K} X_{\mathbf{i}} = \sum_{i_2=1}^{[nt_2]} \cdots \sum_{i_d=1}^{[nt_d]} X_{([nt_1]+1, i_2, \dots, i_d)} = \Delta_{[nt_1]+1}^{(1)} S_{[nt]}.$$

Then it should be clear that

$$\sum_{\mathbf{i} \in I_K} X_{\mathbf{i}} = \left( \prod_{k \in K} \Delta_{[n_k t_k]+1}^{(k)} \right) S_{[nt]}.$$

Now observe that

$$T_l(\mathbf{t}) = \sum_{\substack{K \subset \{1, \dots, d\} \\ \text{card}(K)=l}} \sum_{\mathbf{i} \in I_K} \left( \prod_{k \in K} s_k \right) X_{\mathbf{i}} = \sum_{\substack{K \subset \{1, \dots, d\} \\ \text{card}(K)=l}} \left( \prod_{k \in K} s_k \right) \sum_{\mathbf{i} \in I_K} X_{\mathbf{i}}.$$

Recalling that  $s_k = \{n_k t_k\}$ , this leads to

$$T_l(\mathbf{t}) = \sum_{\substack{K \subset \{1, \dots, d\} \\ \text{card}(K)=l}} \left( \prod_{k \in K} \{n_k t_k\} \right) \left( \prod_{k \in K} \Delta_{[n_k t_k]+1}^{(k)} \right) S_{[nt]}. \quad (2.27)$$

Finally we obtain the representation (2.14) and complete the proof.

### 2.1.3 Estimate of sequential Hölder norm

Using the results from previous sections we give now the estimate of sequential norm of  $\boldsymbol{\pi}(\mathbf{n})^{-1/2} \xi_{\mathbf{n}}$  in terms of  $m$ -indexed sums. We use this result later for proving the tightness of process  $\boldsymbol{\pi}(\mathbf{n})^{-1/2} \xi_{\mathbf{n}}$ . Recall that sequential norm of  $x \in \mathbb{H}_{\alpha}^o([0, 1]^d)$  is defined as

$$\|x\|_{\alpha}^{\text{seq}} = \sup_{j \geq 0} 2^{\alpha j} \max_{\mathbf{v} \in V_j} \|\lambda_{j, \mathbf{v}}(x)\|.$$

Recall from 1.2.2 that a dyadic point  $\mathbf{v} \in V_j$  can be expressed as  $\mathbf{v} = (k_1 2^{-j}, \dots, k_d 2^{-j})$  with at least one  $k_i$  odd. Denote by  $K = \{j_1, \dots, j_l\}$  the set of indices for which coordinates of  $2^j \mathbf{v}$  are odd. Then

$$\xi_{\mathbf{n}}(\mathbf{v}) - \xi_{\mathbf{n}}(\mathbf{v}^+) = \sum_{i=1}^{\text{card}(K)} \xi_{\mathbf{n}} \left( \mathbf{v} + 2^{-j} \sum_{k=1}^{i-1} e_{j_k} \right) - \xi_{\mathbf{n}} \left( \mathbf{v} + 2^{-j} \sum_{k=1}^i e_{j_k} \right)$$

similarly

$$\xi_n(\mathbf{v}) - \xi_n(\mathbf{v}^-) = \sum_{i=1}^{\text{card}(K)} \xi_n \left( \mathbf{v} - 2^{-j} \sum_{k=1}^{i-1} e_{j_k} \right) - \xi_n \left( \mathbf{v} - 2^{-j} \sum_{k=1}^i e_{j_k} \right)$$

thus we can express double difference

$$\lambda_{j,v}(\xi_n) = \xi_n(\mathbf{v}) - \frac{1}{2}(\xi_n(\mathbf{v}^+) + \xi_n(\mathbf{v}^-))$$

as differences of process  $\xi_n$  where only one coordinate is changing. Define

$$\Delta_n^{(1)}(t, t'; \mathbf{s}) := \|\xi_n(t', s_2, \dots, s_d) - \xi_n(t, s_2, \dots, s_d)\|,$$

for the change in the first coordinate and similarly  $\Delta_n^{(j)}(t, t'; \mathbf{s})$  for the change in the  $j$ -th coordinate. Then

$$\max_{\mathbf{v} \in V_j} \|\lambda_{j,v}(\xi_n)\| \leq \sum_{m=1}^d \max_{\substack{0 \leq k < 2^j \\ 0 \leq \ell \leq 2^j}} \Delta_n^{(m)}(t_{k+1}, t_k; \mathbf{s}_\ell) \quad (2.28)$$

with  $t_k = k2^{-j}$ ,  $\ell = (l_2, \dots, l_d)$ ,  $\mathbf{2}^j = (2^j, \dots, 2^j)$  (vector of dimension  $d-1$ ) and  $\mathbf{s}_\ell = \ell \mathbf{2}^{-j}$ .

Let us first examine the the case  $d = 1$ . Then

$$\max_{\mathbf{v} \in V_j} \|\lambda_{j,v}(\xi_n)\| \leq \max_{0 \leq k < 2^j} |\xi_n(t_{k+1}) - \xi_n(t_k)|.$$

Denote by “log” the logarithm *with basis 2* ( $\log 2 = 1$ ). If  $j > \log n$  then  $t_{k+1} - t_k = 2^{-j} < 1/n$  and from definition (2.1) we get

$$\xi_n(t_{k+1}) - \xi_n(t_k) = n2^{-j} X_{[nt_k]+1}$$

if  $[nt_{k+1}] = [nt_k]$ . For  $j \leq \log n$  we get

$$\xi_n(t_{k+1}) - \xi_n(t_k) = \sum_{i=[nt_k]+1}^{[nt_{k+1}]} X_i$$

if  $n = 2^l$  with  $l > j$ . With little additional work it is possible to refine these

expressions to get the estimate

$$\begin{aligned} \max_{0 \leq k < 2^j} |\xi_n(t_{k+1}) - \xi_n(t_k)| &\leq \mathbf{1}(j \leq \log n) \max_{0 \leq k < 2^j} \sum_{i=[nt_k]+1}^{[nt_{k+1}]} |X_i| \\ &\quad + \mathbf{1}(j > \log n) n 2^{-j+1} \max_{1 \leq i \leq n} |X_i|. \end{aligned}$$

For the case  $d > 1$  it is possible to get similar estimate given by the following lemma.

**Lemma 13** For  $m = 1, \dots, d$  and any  $t', t \in [0, 1], t' > t$ ,

$$\begin{aligned} \sup_{s \in [0, 1]^{d-1}} \Delta_n^{(m)}(t, t'; \mathbf{s}) &\leq 3^d \mathbf{1}\left(t' - t \geq \frac{1}{n_m}\right) \psi_n^{(m)}(t', t) \\ &\quad + 3^d \min\left(1, n_m(t' - t)\right) Z_n^{(m)}, \end{aligned}$$

where

$$\psi_n^{(m)}(t', t) := \max_{\mathbf{1}_{-m} \leq \mathbf{k}_{-m} \leq \mathbf{n}_{-m}} \left\| \sum_{k_m=[n_m t]+1}^{[n_m t']} \Delta_{k_m}^{(m)} S_{k_m} \right\|, \quad (2.29)$$

$$Z_n^{(m)} := \max_{1 \leq \mathbf{k} \leq \mathbf{n}} \|\Delta_{k_1}^{(m)} S_{k_1}\|. \quad (2.30)$$

Thus

$$\begin{aligned} \max_{\substack{0 \leq k < 2^j \\ 0 \leq \ell \leq 2^j}} \Delta_n^{(m)}(t_{k+1}, t_k; \mathbf{s}_\ell) &\leq \max_{0 \leq k < 2^j} \left[ 3^d \mathbf{1}(t_{k+1} - t_k \geq 1/n_m) \psi_n^{(m)}(t_{k+1}, t_k) \right. \\ &\quad \left. + 3^d \min\{1, n_m(t_{k+1} - t_k)\} Z_n^{(m)} \right] \end{aligned}$$

It is possible to further refine this expression by considering certain values of  $j$ . For  $j > \log n_m$ , we have  $2^j > n_m$ , whence  $(t_{k+1} - t_k) = 2^{-j} < 1/n_m$  thus in this case

$$\max_{\substack{0 \leq k < 2^j \\ 0 \leq \ell \leq 2^j}} \Delta_n^{(m)}(t_{k+1}, t_k; \mathbf{s}_\ell) \leq 3^d \max_{0 \leq k < 2^j} n_m(t_{k+1} - t_k) Z_n^{(m)}.$$

Now since  $2^{-j} < n_m^{-1}$  we have

$$2^{j\alpha} n_m(t_{k+1} - t_k) = 2^{-j(1-\alpha)} n_m < n_m^\alpha,$$

giving us

$$2^{j\alpha} \max_{\substack{0 \leq k < 2^j \\ 0 \leq \ell \leq 2^j}} \Delta_{\mathbf{n}}^{(m)}(t_{k+1}, t_k; \mathbf{s}_\ell) \leq 3^d n_m^\alpha Z_{\mathbf{n}}^{(m)} \quad (2.31)$$

for  $j > \log n_m$ . On the other hand, for  $j \leq \log n_m$ , we have  $2^{\alpha j} \leq n_m^\alpha$ , whence

$$2^{j\alpha} \max_{\substack{0 \leq k < 2^j \\ 0 \leq \ell \leq 2^j}} \Delta_{\mathbf{n}}^{(m)}(t_{k+1}, t_k; \mathbf{s}_\ell) \leq 3^d \left[ \max_{j \leq \log n_m} \max_{0 \leq k < 2^j} \psi_{\mathbf{n}}^{(m)}(t_{k+1}, t_k) + n_m^\alpha Z_{\mathbf{n}}^{(m)} \right] \quad (2.32)$$

for  $j \leq \log n_m$ . Reporting now (2.31) and (2.32) to (2.28) we get

$$\|\xi_{\mathbf{n}}\|_{\alpha}^{\text{seq}} \leq 3^d \sum_{m=1}^d \left( \max_{j \leq \log n_m} 2^{\alpha j} \max_{0 \leq k < 2^j} \psi_{\mathbf{n}}^{(m)}(t_{k+1}, t_k) + n_m^\alpha Z_{\mathbf{n}}^{(m)} \right). \quad (2.33)$$

Recalling the definition of  $E_J$  from (1.3) and noting that we can restrict the domain of  $j$  in inequality (2.32) we also get the estimate for  $\|\xi_{\mathbf{n}} - E_J \xi_{\mathbf{n}}\|$

$$\|\xi_{\mathbf{n}} - E_J \xi_{\mathbf{n}}\|_{\alpha}^{\text{seq}} \leq 3^d \sum_{m=1}^d \left( \max_{J \leq j \leq \log n_m} 2^{\alpha j} \max_{0 \leq k < 2^j} \psi_{\mathbf{n}}^{(m)}(t_{k+1}, t_k) + n_m^\alpha Z_{\mathbf{n}}^{(m)} \right). \quad (2.34)$$

From these inequalities we see that the sequential norm of process  $\xi_{\mathbf{n}}$  can be controlled by only first differences of the process when only one coordinate changes.

### Proof of lemma 13

We prove this lemma in case of  $m = 1$ , proof for other cases is identical. Put  $\mathbf{u} := (t, \mathbf{s})$ ,  $\mathbf{u}' := (t', \mathbf{s})$ , so  $u_1 = t$ ,  $u'_1 = t'$  and  $\mathbf{u}_{2:d} = \mathbf{u}'_{2:d} = \mathbf{s}$ . Denote

$$T_l(\mathbf{t}) = \sum_{1 \leq i_1 < i_2 < \dots < i_l \leq d} \left( \prod_{k=1}^l \{n_{i_k} t_{i_k}\} \right) \left( \prod_{k=1}^l \Delta_{[n_{i_k} t_{i_k}] + 1}^{(i_k)} \right) S_{[nt]}.$$

Then from representation 2.14 we have

$$\xi_{\mathbf{n}}(\mathbf{u}') - \xi_{\mathbf{n}}(\mathbf{u}) = S_{[nu']} - S_{[nu]} + \sum_{l=1}^d (T_l(\mathbf{u}') - T_l(\mathbf{u})). \quad (2.35)$$

To estimate this  $\xi_{\mathbf{n}}$ 's increment we discuss according to the different possible configurations.

Case 1.  $0 < t' - t < 1/n_1$ .

Case 1.a.  $[n_1 t'] = [n_1 t]$ , whence  $[n\mathbf{u}'] = [n\mathbf{u}]$ . Consider first the increment  $T_1(\mathbf{u}') - T_1(\mathbf{u})$  and note that by (2.27) with  $l = 1$ ,

$$T_1(\mathbf{u}) = \sum_{1 \leq k \leq d} \{n_k u_k\} \Delta_{[n_k u_k]+1}^{(k)} S_{[n\mathbf{u}]}.$$

Because  $\mathbf{u}_{2:d} = \mathbf{u}'_{2:d}$  and  $[n\mathbf{u}'] = [n\mathbf{u}]$ , all the terms indexed by  $k \geq 2$  disappear in the difference  $T_1(\mathbf{u}') - T_1(\mathbf{u})$ . Note also that  $\{n_1 t'\} - \{n_1 t\} = n_1(t' - t)$ . This leads to the factorization

$$T_1(\mathbf{u}') - T_1(\mathbf{u}) = n_1(t' - t) \Delta_{[n_1 t]+1}^{(1)} S_{[n\mathbf{u}]}.$$

For  $l \geq 2$ ,  $T_l(\mathbf{u})$  is expressed by (2.27) as

$$T_l(\mathbf{u}) = \sum_{1 \leq i_1 < \dots < i_l \leq d} \{n_{i_1} u_{i_1}\} \dots \{n_{i_l} u_{i_l}\} \Delta_{[n_{i_1} u_{i_1}]+1}^{(i_1)} \dots \Delta_{[n_{i_l} u_{i_l}]+1}^{(i_l)} S_{[n\mathbf{u}]}.$$

As above, all the terms for which  $i_1 \geq 2$  disappear in the difference  $T_l(\mathbf{u}') - T_l(\mathbf{u})$  and we obtain

$$T_l(\mathbf{u}') - T_l(\mathbf{u}) = n_1(t' - t) \sum_{1 < i_2 < \dots < i_l \leq d} \{n_{i_2} s_{i_2}\} \dots \{n_{i_l} s_{i_l}\} \Delta_{[n_1 t]+1}^{(1)} \Delta_{[n_{i_2} s_{i_2}]+1}^{(i_2)} \dots \Delta_{[n_{i_l} s_{i_l}]+1}^{(i_l)} S_{[n\mathbf{u}]}.$$

Since  $\{n_{i_2} s_{i_2}\} \dots \{n_{i_l} s_{i_l}\} < 1$  and

$$\begin{aligned} \left\| \Delta_{[n_1 t]+1}^{(1)} \Delta_{[n_{i_2} s_{i_2}]+1}^{(i_2)} \dots \Delta_{[n_{i_l} s_{i_l}]+1}^{(i_l)} S_{[n\mathbf{u}]} \right\| &= \left\| \Delta_{[n_1 t]+1}^{(1)} \sum_{\mathbf{i} \in I} \varepsilon_{\mathbf{i}} S_{\mathbf{i}} \right\| \\ &\leq \sum_{\mathbf{i} \in I} \left\| \Delta_{[n_1 t]+1}^{(1)} S_{\mathbf{i}} \right\|, \end{aligned}$$

where  $\varepsilon_{\mathbf{i}} = \pm 1$  and  $I$  is some appropriate subset of  $[0, \mathbf{n}] \cap \mathbb{N}^d$  with  $2^{l-1}$  elements. Hence with  $Z_n$  defined by (2.30), we obtain for  $l \geq 2$

$$\|T_l(\mathbf{u}') - T_l(\mathbf{u})\| \leq n_1(t' - t) \binom{d-1}{l-1} 2^{l-1} Z_n.$$

Clearly this estimate holds true also for  $l = 1$ , so going back to (2.35) and

recalling that in the case under consideration  $[\mathbf{n}\mathbf{u}'] = [\mathbf{n}\mathbf{u}]$ , we obtain

$$\|\xi_n(\mathbf{u}') - \xi_n(\mathbf{u})\| \leq \sum_{l=1}^d n_1(t' - t) \binom{d-1}{l-1} 2^{l-1} Z_n = 3^{d-1} n_1(t' - t) Z_n. \quad (2.36)$$

*Case 1.b.*  $n_1 t < [n_1 t'] \leq n_1 t'$ . Using chaining to exploit the result of case 1.a, we obtain

$$\begin{aligned} \|\xi_n(\mathbf{u}') - \xi_n(\mathbf{u})\| &\leq \left\| \xi_n(\mathbf{u}') - \xi_n\left(\frac{[n_1 t']}{n_1}, \mathbf{s}\right) \right\| + \left\| \xi_n\left(\frac{[n_1 t']}{n_1}, \mathbf{s}\right) - \xi_n(\mathbf{u}) \right\| \\ &\leq 3^{d-1} (n_1 t' - [n_1 t']) Z_n + 3^{d-1} ([n_1 t'] - n_1 t) Z_n \\ &= 3^{d-1} n_1 (t' - t) Z_n. \end{aligned} \quad (2.37)$$

*Case 2.*  $t' - t \geq 1/n_1$ . Then  $[n_1 t] \leq n_1 t < [n_1 t] + 1 \leq [n_1 t'] \leq n_1 t'$  and putting

$$t_1 := \frac{[n_1 t]}{n_1}, \quad t'_1 := \frac{[n_1 t']}{n_1}, \quad \mathbf{v} := (t_1, \mathbf{s}), \quad \mathbf{v}' := (t'_1, \mathbf{s}),$$

we get the upper bound

$$\begin{aligned} \|\xi_n(\mathbf{u}') - \xi_n(\mathbf{u})\| &\leq \|\xi_n(\mathbf{u}') - \xi_n(\mathbf{v}')\| + \|\xi_n(\mathbf{v}') - \xi_n(\mathbf{v})\| \\ &\quad + \|\xi_n(\mathbf{v}) - \xi_n(\mathbf{u})\|, \end{aligned}$$

where the first and third terms fall within the case 1 since  $t' - t'_1 < 1/n_1$  and  $t - t_1 < 1/n_1$ . As  $n_1 v_1 = n_1 t_1 = [n_1 t]$ , we have

$$[n\mathbf{v}] = ([n_1 t_1], [\mathbf{n}_{2:d} \mathbf{s}]) = [n\mathbf{u}] \quad \text{and} \quad \{n_1 v_1\} = \{[n_1 t]\} = 0,$$

so the representation (2.14) for  $\xi_n(\mathbf{v})$  may be recast as

$$\xi_n(\mathbf{v}) = S_{[n\mathbf{u}]} + \sum_{l=1}^{d-1} \sum_{2 \leq i_1 < i_2 < \dots < i_l \leq d} \left( \prod_{k=1}^l \{n_{i_k} v_{i_k}\} \right) \left( \prod_{k=1}^l \Delta_{[n_{i_k} v_{i_k}] + 1}^{(i_k)} \right) S_{[n\mathbf{u}]}.$$

Clearly the same representation holds for  $\xi_n(\mathbf{v}')$ , by just replacing  $\mathbf{u}$  by  $\mathbf{u}'$ . Now since  $\Delta$ 's are interchangeable and

$$S_{[n\mathbf{u}']} - S_{[n\mathbf{u}]} = \sum_{i=[nt]+1}^{[nt']} \Delta_i^{(1)} S_{(i, [\mathbf{n}_{2:d} \mathbf{s}])},$$

we get

$$\|\xi_n(\mathbf{v}') - \xi_n(\mathbf{v})\| \leq \psi_n(t', t) \sum_{l=0}^{d-1} \binom{d-1}{l} 2^l = 3^{d-1} \psi_n(t', t),$$

with  $\psi_n(t', t)$  defined by (2.29). Using case 1 to bound  $\|\xi_n(\mathbf{u}') - \xi_n(\mathbf{v}')\|$  and  $\|\xi_n(\mathbf{v}) - \xi_n(\mathbf{u})\|$ , we obtain

$$\begin{aligned} \|\xi_n(t', s) - \xi_n(t, s)\| &\leq 3^{d-1} \{n_1 t'\} Z_n + 3^{d-1} \psi_n(t', t) + 3^{d-1} \{n_1 t\} Z_n \\ &\leq 3^{d-1} \psi_n(t', t) + 2 \cdot 3^{d-1} Z_n. \end{aligned} \quad (2.38)$$

Combining (2.36), (2.37) and (2.38) we complete the proof of lemma 13.  $\square$

## 2.2 Unequal variance

Let us examine a simple example. Take collection  $\{X_{n,k}, k = 1, \dots, 2n\}$  of random variables i.i.d. for each  $n$  with zero mean and variance  $\mathbf{E} X_{n,k}^2 = 1/(2n)$ . Then the sum  $S_{n,2n} = X_{n,1} + \dots + X_{n,2n}$  converges to standard normal and the polygonal line process  $\xi_n(t) = S_{n,[2nt]} + (2nt - [2nt])X_{[2nt]+1}$  converges to Brownian motion. Now introduce collection  $\{Y_{n,k}, k = 1, \dots, 2n\}$  by taking  $Y_{n,k} = 1/\sqrt{2}X_{n,k}$  for  $k \leq n$ , and  $Y_{n,k} = \sqrt{3/2}X_{n,k}$  for  $k = n+1, \dots, 2n$ . The sum  $S'_{n,2n} = Y_{n,1} + \dots + Y_{n,2n}$  still converges to standard normal. But for polygonal line process  $\xi'_n(t) = S'_{n,[2nt]} + (2nt - [2nt])Y_{[2nt]+1}$  we then have  $\mathbf{E} (\xi_n(1/2)')^2 = 1/4$  for all  $n$ . Thus if functional central limit theorem holds the variance of limiting process at  $1/2$  is  $1/4$ . Yet Brownian motion variance at  $1/2$  is  $1/2$ , thus the limiting process (if it exists) in this case is not the Brownian motion. From classical result of Prokhorov [24] we know that for triangular arrays it is possible to use different construction of summation process so that the functional central limit theorem holds and the limiting process is always Brownian motion. Furthermore in the case of i.i.d. random variables both definitions coincide. We propose similar definition for  $[0, 1]^d$  indexed summation processes, which though does not solve the problem completely as in case  $d = 1$ , is nevertheless an improvement on using the definition (2.4).

### 2.2.1 Definitions and representations

Let us first review the case of  $[0, 1]$  indexed summation process, i.e. the classical result of Prokhorov [24] for triangular arrays. Suppose we have collection  $\{X_{n,k}, 1 \leq k \leq k_n, k_n, n \in \mathbb{N}\}$  of random variables. Let  $\mathbf{E} X_{n,k}^2 =$

$\sigma_{n,k}^2$  and assume that  $\sum_{k=1}^{k_n} \sigma_{n,k}^2 = 1$ . Define

$$S_n(k) = X_{n,1} + \cdots + X_{n,k}$$

and

$$b_n(k) = \sigma_{n,1}^2 + \cdots + \sigma_{n,k}^2.$$

Then classical definition of summation process is

$$\xi_n(t) = S_n(k) + (t - b_n(k))\sigma_{n,k+1}^{-2}X_{n,k+1}, \text{ for } b_n(k) \leq t < b_n(k+1). \quad (2.39)$$

Define

$$u_n(t) = \min(k : b_n(k) < t),$$

then

$$\mathbf{E} \xi_n(t)^2 = \sum_{k=1}^{u_n(t)} \sigma_{n,k}^2 + \frac{(t - b_n(k))^2}{\sigma_{n,k+1}^2}.$$

If we assume

$$\max_{1 \leq k \leq k_n} \sigma_{n,k}^2 \rightarrow 0, \text{ as } n \rightarrow \infty$$

which is necessary for central limit theorem to apply, we have

$$\mathbf{E} \xi_n(t)^2 \rightarrow t. \quad (2.40)$$

Define triangular array with multidimensional index as

$$(X_{n,\mathbf{k}}, \mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n), \mathbf{n} \in \mathbb{N}^d,$$

where for each  $\mathbf{n}$  the random variables  $X_{n,\mathbf{k}}$  are independent. The expression  $\mathbf{k}_n$  is the element from  $\mathbb{N}^d$  with multidimensional index:  $\mathbf{k}_n = (k_n^1, \dots, k_n^d)$ . Assume that  $\mathbf{E} X_{n,\mathbf{k}} = 0$  and that  $\sigma_{n,\mathbf{k}}^2 := \mathbf{E} X_{n,\mathbf{k}}^2 < \infty$ , for  $\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n$ ,  $\mathbf{n} \in \mathbb{N}^d$ . Define for each  $\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n$

$$S_n(\mathbf{k}) := \sum_{j \leq \mathbf{k}} X_{n,j}, \quad b_n(\mathbf{k}) := \sum_{j \leq \mathbf{k}} \sigma_{n,j}^2.$$

We require that the sum of all variances is one, i.e.  $b_n(\mathbf{k}_n) = 1$  and that  $m(\mathbf{k}_n) \rightarrow \infty$ , as  $m(\mathbf{n}) \rightarrow \infty$ . Note that these requirements are the same as for one-dimensional triangular array.

If  $\pi(\mathbf{k}) = 0$ , let  $S_n(\mathbf{k}) = 0$ ,  $b_n(\mathbf{k}) = 0$ . For  $i = 1, \dots, d$  introduce the

notations

$$\begin{aligned} b_i(k) &:= b_{\mathbf{n}}(k_{\mathbf{n}}^1, \dots, k_{\mathbf{n}}^{i-1}, k, k_{\mathbf{n}}^{i+1}, \dots, k_{\mathbf{n}}^d), \\ \Delta b_i(k) &:= b_i(k) - b_i(k-1) \end{aligned} \quad (2.41)$$

and

$$\mathbf{B}(\mathbf{k}) = (b_1(k_1), \dots, b_d(k_d)), \quad \Delta \mathbf{B}(\mathbf{k}) := (\Delta b_1(k_1), \dots, \Delta b_d(k_d)) \quad (2.42)$$

for its vector counterparts. Note that these variables depend on  $\mathbf{n}$  and  $\mathbf{k}_{\mathbf{n}}$ . For  $\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_{\mathbf{n}}$  let

$$Q_{\mathbf{n}, \mathbf{k}} := \left[ b_1(k_1 - 1), b_1(k_1) \right) \times \dots \times \left[ b_d(k_d - 1), b_d(k_d) \right). \quad (2.43)$$

Due to definition of  $b_i(k)$  we have  $Q_{\mathbf{n}, \mathbf{j}} \cap Q_{\mathbf{n}, \mathbf{k}} = \emptyset$ , if  $\mathbf{k} \neq \mathbf{j}$ . Also  $\cup_{\mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} Q_{\mathbf{n}, \mathbf{k}} = [0, 1)^d$  and  $\sum_{\mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} |Q_{\mathbf{n}, \mathbf{k}}| = 1$  with  $|Q_{\mathbf{n}, \mathbf{k}}| = \pi(\Delta \mathbf{B}(\mathbf{k}))$ . Thus any  $\mathbf{t} \in [0, 1)^d$  falls into unique rectangle  $Q_{\mathbf{n}, \mathbf{k}}$ , for some  $\mathbf{k}$ . In that case trivial equality

$$t_i = b_i(k_i - 1) + \frac{t_i - b_i(k_i - 1)}{\Delta b_i(k_i)} \Delta b_i(k_i)$$

gives

$$\mathbf{t} = \mathbf{B}(\mathbf{k} - \mathbf{1}) + \frac{\mathbf{t} - \mathbf{B}(\mathbf{k} - \mathbf{1})}{\Delta \mathbf{B}(\mathbf{k})} \Delta \mathbf{B}(\mathbf{k})$$

with

$$0 \leq \frac{\mathbf{t} - \mathbf{B}(\mathbf{k} - \mathbf{1})}{\Delta \mathbf{B}(\mathbf{k})} < \mathbf{1}.$$

This corresponds to decomposition

$$\mathbf{t} = \frac{[\mathbf{n}\mathbf{t}]}{\mathbf{n}} + \frac{\{\mathbf{n}\mathbf{t}\}}{\mathbf{n}}.$$

It is natural then that summation process defined on the grid  $Q_{\mathbf{n}, \mathbf{k}}$  as

$$\Xi_{\mathbf{n}}(\mathbf{t}) = \sum_{\mathbf{1} \leq \mathbf{j} \leq \mathbf{n}} |Q_{\mathbf{n}, \mathbf{j}}|^{-1} |Q_{\mathbf{n}, \mathbf{j}} \cap [0, \mathbf{t}]| X_{\mathbf{n}, \mathbf{j}}, \quad (2.44)$$

similar to (2.4) assumes the same representations as process  $\xi_{\mathbf{n}}(\mathbf{t})$ . For  $t \in$

$[0, 1]$  and  $\mathbf{t} \in [0, 1]^d$ , write

$$u_i(t) := \max\{j \geq 0 : b_i(j) \leq t\}, \quad \mathbf{U}(\mathbf{t}) := (u_1(t_1), \dots, u_d(t_d)).$$

Then following proposition holds.

**Proposition 14** For  $\mathbf{t} \in [0, 1]^d$ , denote

$$\mathbf{s} = \frac{\mathbf{t} - \mathbf{B}(\mathbf{U}(\mathbf{t}))}{\Delta \mathbf{B}(\mathbf{U}(\mathbf{t}) + \mathbf{1})}$$

and write vertices of the rectangle  $R_{\mathbf{n}, \mathbf{U}(\mathbf{t}) + \mathbf{1}}$  as

$$V(\mathbf{u}) := \mathbf{B}(\mathbf{U}(\mathbf{t})) + \mathbf{u} \Delta \mathbf{B}(\mathbf{U}(\mathbf{t}) + \mathbf{1}). \quad \mathbf{u} \in \{0, 1\}^d. \quad (2.45)$$

It is possible to express  $\mathbf{t}$  as a barycenter of these  $2^d$  vertices with weights  $w(\mathbf{u}) \geq 0$  depending on  $\mathbf{t}$ , i.e.,

$$\mathbf{t} = \sum_{\mathbf{u} \in \{0, 1\}^d} w(\mathbf{u}) V(\mathbf{u}), \quad \text{where} \quad \sum_{\mathbf{u} \in \{0, 1\}^d} w(\mathbf{u}) = 1, \quad (2.46)$$

where

$$w(\mathbf{u}) = \prod_{l=1}^d s_l^{u_l} (1 - s_l)^{1-u_l}.$$

Using this representation, define the random field  $\Xi_n^*$  by

$$\Xi_n^*(\mathbf{t}) = \sum_{\mathbf{u} \in \{0, 1\}^d} w(\mathbf{u}) S_n(\mathbf{U}(\mathbf{t}) + \mathbf{u}), \quad \mathbf{t} \in [0, 1]^d.$$

Then  $\Xi_n^*$  coincides with the summation process defined by (2.44). Furthermore  $\Xi_n(\mathbf{t})$  admits representation

$$\begin{aligned} \Xi_n(\mathbf{t}) &= S_n(\mathbf{U}(\mathbf{t})) + \\ &\sum_{l=1}^d \sum_{1 \leq i_1 < i_2 < \dots < i_l \leq d} \left( \prod_{k=1}^l \frac{t_{i_k} - b_{i_k}(u_{i_k})}{\Delta b_{i_k}(u_{i_k}(t_{i_k}) + 1)} \right) \left( \prod_{k=1}^l \Delta_{u_{i_k}(t_{i_k}) + 1}^{(i_k)} \right) S_n(\mathbf{U}(\mathbf{t})). \end{aligned} \quad (2.47)$$

*Proof.* The proof is the same as in proposition 12 with the change of notation:  $\{\mathbf{n}\mathbf{t}\}$  changed to  $\mathbf{U}(\mathbf{t})$  and  $\{\mathbf{n}\mathbf{t}\}$  to  $(\mathbf{t} - \mathbf{B}(\mathbf{U}(\mathbf{t}))) / \Delta \mathbf{B}(\mathbf{U}(\mathbf{t}) + \mathbf{1})$ .

### 2.2.2 Estimate of sequential norm

We give now the estimate of sequential norm of  $\Xi_n$ . The estimate is similar to the one given in section 2.1.3 for process  $\xi_n$ . As in section 2.1.3 we can write

$$\max_{v \in V_j} \|\lambda_{j,v}(\Xi_n)\| \leq \sum_{m=1}^d \max_{\substack{0 \leq k < 2^j \\ 0 \leq \ell \leq 2^j}} \Delta_n^{(m)}(t_{k+1}, t_k; \mathbf{s}_\ell) \quad (2.48)$$

with  $t_k = k2^{-j}$ ,  $\ell = (l_2, \dots, l_d)$ ,  $\mathbf{2}^j = (2^j, \dots, 2^j)$  (vector of dimension  $d-1$ ),  $\mathbf{s}_\ell = \ell \mathbf{2}^{-j}$ , and  $\Delta_n^{(m)}(t, t'; \mathbf{s})$  defined for  $m = 1$  as

$$\Delta_n^{(1)}(t, t'; \mathbf{s}) := |\Xi_n(t', \mathbf{s}) - \Xi_n(t, \mathbf{s})|$$

and similarly for other coordinates for  $m > 1$ . Introduce set  $D_j = \{2(l-1)2^{-j}; 1 \leq l \leq 2^{j-1}\}$  and notation  $r^- = r - 2^{-j}$  and  $r^+ = r + 2^{-j}$  for  $r \in D_j$ . Then

$$\max_{v \in V_j} \|\lambda_{j,v}(\Xi_n)\| \leq \sum_{m=1}^d \max \left\{ \max_{\substack{r \in D_j \\ 0 \leq \ell \leq 2^j}} \Delta_n^{(m)}(r, r^-; \mathbf{s}_\ell), \max_{\substack{r \in D_j \\ 0 \leq \ell \leq 2^j}} \Delta_n^{(m)}(r^+, r; \mathbf{s}_\ell) \right\}. \quad (2.49)$$

At first glance this separation seems unnecessary, especially since the treatment of both  $\Delta_n^{(m)}(r, r^-; \mathbf{s}_\ell)$  and  $\Delta_n^{(m)}(r, r^+; \mathbf{s}_\ell)$  is identical, but this simplifies the proofs later on. Similar lemma to 13 holds.

**Lemma 15** For  $m = 1, \dots, d$  and any  $r \in D_j$

$$\begin{aligned} \sup_{\mathbf{s} \in [0,1]^{d-1}} \Delta_n^{(m)}(r, r^-; \mathbf{s}) &\leq 3^d \mathbf{1}(u_m(r) > u_m(r^-) + 1) \psi_n^{(m)}(r, r^-) \\ &\quad + 3^d 2^{-j\alpha} Z_n^{(m)}, \end{aligned}$$

where

$$\psi_n(r, r^-)^{(m)} := \max_{\mathbf{k}_{-m} \leq (\mathbf{k}_n)_{-m}} \left| \sum_{k_m = u_m(r^-) + 2}^{u_m(r)} \Delta_{k_m}^{(m)} S_n(\mathbf{k}) \right| \quad (2.50)$$

$$Z_n^{(m)} := \max_{1 \leq k \leq k_n} \frac{|\Delta_{k_m}^{(m)} S_n(\mathbf{k})|}{(\Delta b_m(k_m))^\alpha}. \quad (2.51)$$

Similarly

**Lemma 16** For  $m = 1, \dots, d$  and any  $r \in D_j$

$$\sup_{\mathbf{s} \in [0,1]^{d-1}} \Delta_{\mathbf{n}}^{(m)}(r, r^+; \mathbf{s}) \leq 3^d \mathbf{1}(u_m(r) > u_m(r^-) + 1) \psi_{\mathbf{n}}^{(m)}(r, r^+) + 3^d 2^{-j\alpha} Z_{\mathbf{n}}^{(m)},$$

where

$$\psi_{\mathbf{n}}(r, r^+)^{(m)} := \max_{\mathbf{k}-m \leq (\mathbf{k}_{\mathbf{n}})-m} \left| \sum_{k_m=u_m(r)+2}^{u_m(r^+)} \Delta_{k_m}^{(m)} S_{\mathbf{n}}(\mathbf{k}) \right|. \quad (2.52)$$

with  $Z_{\mathbf{n}}^{(m)}$  as in (2.51).

Note that only definitions of  $\psi_{\mathbf{n}}$  differs and  $Z_{\mathbf{n}}^{(m)}$  does not depend on  $j$ . We make no distinction for different  $j$  and derive immediately estimate of sequential norm

$$\|\Xi_{\mathbf{n}}\|_{\alpha}^{\text{seq}} \leq 3^d \sum_{m=1}^d \left( \max_{j \geq 0} 2^{j\alpha} \max_{r \in D_j} [\psi_{\mathbf{n}}^{(m)}(r, r^-) + \psi_{\mathbf{n}}^{(m)}(r, r^+)] + Z_{\mathbf{n}}^{(m)} \right) \quad (2.53)$$

and the tail

$$\begin{aligned} & \|\Xi_{\mathbf{n}} - E_J \Xi_{\mathbf{n}}\|_{\alpha}^{\text{seq}} \\ & \leq 3^d \sum_{m=1}^d \left( \max_{j \geq J} 2^{j\alpha} \max_{r \in D_j} [\psi_{\mathbf{n}}^{(m)}(r, r^-) + \psi_{\mathbf{n}}^{(m)}(r, r^+)] + Z_{\mathbf{n}}^{(m)} \right). \end{aligned} \quad (2.54)$$

### Proof of lemma 15

We prove this for  $m = 1$  since the proof is the same for other  $m$ , subsequently we drop the superscript in definitions  $Z_{\mathbf{n}}$  and  $\psi_{\mathbf{n}}$ . The proof is similar to proof of lemma 13. Denote by  $\mathbf{v} = (r, \mathbf{s})$ , and  $\mathbf{v}^- = (r^-, \mathbf{s})$ . Recall representation (2.47) and write

$$T_l(\mathbf{t}) = \sum_{1 \leq i_1 < i_2 < \dots < i_l \leq d} \left( \prod_{k=1}^l \frac{t_{i_k} - b_{i_k}(u_{i_k})}{\Delta b_{i_k}(u_{i_k}(t_{i_k}) + 1)} \right) \left( \prod_{k=1}^l \Delta_{u_{i_k}(t_{i_k})+1}^{(i_k)} \right) S_{\mathbf{n}}(\mathbf{U}(\mathbf{t}))$$

for  $l = 1, \dots, d$ . Then

$$\begin{aligned} \Xi_n(r, s_\ell) - \Xi_n(r^-, s_\ell) &= S_n(\mathbf{U}(\mathbf{v})) - S_n(\mathbf{U}(\mathbf{v}^-)) \\ &\quad + \sum_{l=1}^d (T_l(\mathbf{v}) - T_l(\mathbf{v}^-)). \end{aligned}$$

To estimate this increment we discuss according to following configurations  
*Case 1.*  $u_1(r) = u_1(r^-)$ . Consider first the increment  $T_1(\mathbf{v}) - T_1(\mathbf{v}^-)$  and note that by (2.27) with  $l = 1$ ,

$$T_1(\mathbf{v}) = \sum_{1 \leq k \leq d} \frac{v_k - b_k(u_k(v_k))}{\Delta b_k(u_k(v_k) + 1)} \Delta_{u_k(v_k)+1}^{(k)} S_n(\mathbf{U}(\mathbf{v})).$$

Because  $\mathbf{v}_{2:d} = \mathbf{v}_{2:d}^-$  and  $\mathbf{U}(\mathbf{v}) = \mathbf{U}(\mathbf{v}^-)$ , all terms indexed by  $k \geq 2$  disappear in difference  $T_1(\mathbf{v}) - T_1(\mathbf{v}^-)$ . This leads to the factorisation

$$T_1(\mathbf{v}) - T_1(\mathbf{v}^-) = \frac{r - r^-}{\Delta b_1(u_1(r) + 1)} \Delta_{u_1(r)+1}^{(1)} S_n(\mathbf{U}(\mathbf{v})). \quad (2.55)$$

For  $l \geq 2$ ,  $T_l(\mathbf{v})$  is expressed by (2.27) as

$$\begin{aligned} T_l(\mathbf{v}) &= \sum_{1 \leq i_1 < \dots < i_l \leq d} \frac{v_{i_1} - b_{i_1}(u_{i_1}(v_{i_1}))}{\Delta b_{i_1}(u_{i_1}(v_{i_1}) + 1)} \cdots \frac{v_{i_l} - b_{i_l}(u_{i_l}(v_{i_l}))}{\Delta b_{i_l}(u_{i_l}(v_{i_l}) + 1)} \\ &\quad \Delta_{u_{i_1}(v_{i_1})+1}^{(i_1)} \cdots \Delta_{u_{i_l}(v_{i_l})+1}^{(i_l)} S_n(\mathbf{U}(\mathbf{v})). \end{aligned}$$

In the difference  $T_l(\mathbf{v}) - T_l(\mathbf{v}^-)$  all the terms for which  $i_1 \geq 2$  again disappear and we obtain

$$\begin{aligned} T_l(\mathbf{v}) - T_l(\mathbf{v}^-) &= \frac{r - r^-}{\Delta b_1(u_1(r) + 1)} \\ &\quad \sum_{1 < i_2 < \dots < i_l \leq d} \frac{v_{i_2} - b_{i_2}(u_{i_2}(v_{i_2}))}{\Delta b_{i_1}(u_{i_1}(v_{i_1}) + 1)} \cdots \frac{v_{i_l} - b_{i_l}(u_{i_l}(v_{i_l}))}{\Delta b_{i_l}(u_{i_l}(v_{i_l}) + 1)} \\ &\quad \Delta_{u_1(r)+1}^{(1)} \Delta_{u_{i_2}(v_{i_2})+1}^{(i_2)} \cdots \Delta_{u_{i_l}(v_{i_l})+1}^{(i_l)} S_n(\mathbf{U}(\mathbf{v})). \end{aligned} \quad (2.56)$$

Since  $u_1(r) = u_1(r^-)$ , we have  $b_1(u_1(r)) \leq r < r^- < b_1(u_1(r) + 1)$ , thus

$$\frac{r - r^-}{\Delta b_1(u_1(r) + 1)} \leq \left( \frac{r - r^-}{\Delta b_1(u_1(r) + 1)} \right)^\alpha.$$

Now

$$\frac{v_{i_2} - b_{i_2}(u_{i_2}(v_{i_2}))}{\Delta b_{i_1}(u_{i_1}(v_{i_1}) + 1)} \cdots \frac{v_{i_l} - b_{i_l}(u_{i_l}(v_{i_l}))}{\Delta b_{i_l}(u_{i_l}(v_{i_l}) + 1)} < 1$$

and

$$\begin{aligned} |\Delta_{u_1(r)+1}^{(1)} \Delta_{u_{i_2}(v_{i_2})+1}^{(i_2)} \cdots \Delta_{u_{i_l}(v_{i_l})+1}^{(i_l)} S_n(\mathbf{U}(\mathbf{v}))| &= |\Delta_{u_1(r)+1}^{(1)} \sum_{\mathbf{i} \in I} \varepsilon_{\mathbf{i}} S_n(\mathbf{i})| \\ &\leq \sum_{\mathbf{i} \in I} |\Delta_{u_1(r)+1}^{(1)} S_n(\mathbf{i})|, \end{aligned} \quad (2.57)$$

where  $\varepsilon_{\mathbf{i}} = \pm 1$  and  $I$  is some appropriate subset of  $[0, \mathbf{n}] \cap \mathbb{N}^d$  with  $2^{l-1}$  elements. Recall that  $Z_n$  is defined as

$$Z_n = \max_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \frac{|\Delta_{k_1}^{(1)} S_n(\mathbf{k})|}{(\Delta b_1(k_1))^\alpha}.$$

Now noting that  $r - r^- = 2^{-j}$  and  $\Delta b_1(k_1)$  depends only on  $k_1$ , we obtain for  $l \geq 2$

$$|T_l(\mathbf{u}') - T_l(\mathbf{u})| \leq 2^{-j\alpha} \binom{d-1}{l-1} 2^{l-1} Z_n.$$

Thus

$$|\Xi_n(\mathbf{v}) - \Xi_n(\mathbf{v}^-)| \leq \sum_{l=1}^d 2^{-j\alpha} \binom{d-1}{l-1} 2^{l-1} Z_n = 3^{d-1} 2^{-j\alpha} Z_n. \quad (2.58)$$

*Case 2.*  $u_1(r) = u_1(r^-) + 1$ . In this case we have  $b_1(u_1(r^-)) \leq r^- < b_1(u_1(r)) \leq r$ . Using previous definitions we can write

$$\begin{aligned} |\Xi_n(\mathbf{v}) - \Xi_n(\mathbf{v}^-)| &\leq |\Xi_n(\mathbf{v}) - \Xi_n(b_1(u_1(r)), \mathbf{s}_\ell)| \\ &\quad + |\Xi_n(b_1(u_1(r)), \mathbf{s}_\ell) - \Xi_n(\mathbf{v}^-)|. \end{aligned}$$

Now

$$\frac{r - b_1(u_1(r))}{\Delta b_1(u_1(r) + 1)} \leq \left( \frac{r - b_1(u_1(r))}{\Delta b_1(u_1(r) + 1)} \right)^\alpha \leq \frac{2^{-j\alpha}}{(\Delta b_1(u_1(r) + 1))^\alpha}$$

and similarly

$$\frac{b_1(u_1(r)) - r^-}{\Delta b_1(u_1(r^-) + 1)} \leq \frac{2^{-j\alpha}}{(\Delta b_1(u_1(r^-) + 1))^\alpha}.$$

Combining these inequalities with (2.55) and (2.56) we get as in (2.36)

$$|\Xi_n(\mathbf{v}) - \Xi_n(\mathbf{v}^-)| \leq 2 \cdot 3^{d-1} 2^{-j\alpha} Z_n.$$

Case 3.  $u_1(r) > u_1(r^-) + 1$ . Put

$$\mathbf{u} = (b_1(u_1(r)), \mathbf{s}_\ell), \quad \mathbf{u}^- = (b_1(u_1(r^-)) + 1, \mathbf{s}_\ell).$$

Then

$$\begin{aligned} |\Xi_n(\mathbf{v}) - \Xi_n(\mathbf{v}^-)| &\leq |\Xi_n(\mathbf{v}) - \Xi_n(\mathbf{u})| + |\Xi_n(\mathbf{u}) - \Xi_n(\mathbf{u}^-)| \\ &\quad + |\Xi_n(\mathbf{u}^-) - \Xi_n(\mathbf{v}^-)|. \end{aligned}$$

Since  $\mathbf{U}(\mathbf{u})_{2:d} = \mathbf{U}(\mathbf{u}^-)_{2:d} = \mathbf{U}(\mathbf{v})_{2:d}$ , we have

$$\begin{aligned} \Xi_n(\mathbf{u}) &= S_n(\mathbf{U}(\mathbf{u})) + \sum_{l=1}^{d-1} \sum_{2 \leq i_1 < i_2 < \dots < i_l \leq d} \left( \prod_{k=1}^l \frac{v_{i_k} - b_{i_k}(u_{i_k}(v_{i_k}))}{\Delta b_{i_k} u_{i_k}(v_{i_k}) + 1} \right) \\ &\quad \left( \prod_{k=1}^l \Delta_{u_{i_k}(v_{i_k})+1}^{(i_k)} \right) S_n(\mathbf{U}(\mathbf{u})) \end{aligned}$$

and similar representation holds for  $\Xi_n(\mathbf{u}^-)$ . We have

$$S_n(\mathbf{U}(\mathbf{u})) - S_n(\mathbf{U}(\mathbf{u}^-)) = \sum_{i=u_1(r^-)+2}^{u_1(r)} \Delta_i^{(1)} S_n((i, \mathbf{U}(\mathbf{s}_\ell))).$$

Recalling the definition

$$\psi_n(r, r^-) = \max_{\mathbf{k}_{2:d} \leq \mathbf{k}_n, 2:d} \left| \sum_{i=u_1(r^-)+2}^{u_1(r)} \Delta_i^{(1)} S_n((i, \mathbf{k}_{2:d})) \right|,$$

similar to (2.57) and (2.58) we get

$$|\Xi_n(\mathbf{u}) - \Xi_n(\mathbf{u}^-)| \leq \psi_n(r, r^-) \sum_{l=0}^{d-1} 2^l \leq 3^{d-1} \psi_n(r, r^-).$$

We can bound  $|\Xi_n(\mathbf{v}) - \Xi_n(\mathbf{u})|$  and  $|\Xi_n(\mathbf{u}^-) - \Xi_n(\mathbf{v})|$  as in case 2. Thus we get

$$|\Xi_n(r, \mathbf{s}_\ell) - \Xi_n(r^-, \mathbf{s}_\ell)| \leq 3^{d-1} \psi_n(r, r^-) + 2 \cdot 3^{d-1} 2^{-j\alpha} Z_n, \quad (2.59)$$

which gives us the proof of the lemma.

**Proof of the lemma 16**

Proof is identical to the proof of lemma 15. Instead of analyzing configurations when  $u_1(r) \geq u_1(r^-)$ , analyze configurations when  $u_1(r^+) \geq u_1(r)$ .

## Chapter 3

# Functional central limit theorems

Functional central limit theorems deal with weak convergence of summation processes. Classical approach is to prove first the convergence of finite-dimensional distributions usually using central limit theorem, and then to show that the summation process is tight. Due to Prokhorov theorem this then gives the weak convergence and we say that functional central limit theorem is proved. Functional central limit theorem is called the invariance principle if necessary and sufficient conditions for the convergence are given. In this section we prove invariance principle for i.i.d. Hilbert space valued random variables in Hölder space. We also prove functional central limit theorem for real valued independent but non-identically distributed random variables. Usually proving tightness is harder task, but that is not necessarily so as we show for the triangular array.

For better readability shorter proofs are given straight after theorems in this chapter. The end of the proof is noted by the symbol  $\square$ .

### 3.1 Invariance principle

#### 3.1.1 Finite dimensional distributions

Recall the definition of summation process  $\xi_n$ :

$$\xi_n(t) = \sum_{i \leq n} \pi(n) \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \cap [0, t] \right| X_i.$$

The so called jump summation process is defined by  $\zeta_n(\mathbf{t}) = S_{[n\mathbf{t}]}$  which can be put alternatively as

$$\zeta_n(\mathbf{t}) = \sum_{j \leq \mathbf{n}} \mathbf{1}(j/\mathbf{n} \in [0, \mathbf{t}]) X_j.$$

The following theorem holds.

**Theorem 17** *Let  $\{X_i, 1 \leq i \leq \mathbf{n}\}$  be a collection of  $\mathbb{H}$ -valued random variables. Assume that all variables have finite second moment and uniform variance  $\sigma^2 = \mathbf{E} \|X_i\|^2 < \infty$  for all  $i \leq \mathbf{n}$ . Then if  $\mathbf{E} \langle X_i, X_j \rangle = 0$  for  $1 \leq i \neq j \leq \mathbf{n}$ ,*

$$\|\pi(\mathbf{n})^{-1/2}(\xi_n(\mathbf{t}) - \zeta_n(\mathbf{t}))\| \xrightarrow[\mathbf{m}(\mathbf{n}) \rightarrow \infty]{\text{Pr}} 0, \quad (3.1)$$

for each  $\mathbf{t} \in [0, 1]^d$ .

*Proof.* For each  $\mathbf{t}$  we can write  $\pi(\mathbf{n})^{-1/2}(\xi_n(\mathbf{t}) - \zeta_n(\mathbf{t})) = \sum_{i \leq \mathbf{n}} \alpha_i X_i$ , where

$$\alpha_i := \pi(\mathbf{n})^{1/2}(|[(i-1)/\mathbf{n}, i/\mathbf{n}] \cap [0, \mathbf{t}]| - \pi(\mathbf{n})^{-1} \mathbf{1}(i/\mathbf{n} \in [0, \mathbf{t}])).$$

Then

$$\mathbf{E} \|\pi(\mathbf{n})^{-1/2}(\xi_n - \zeta_n)\|^2 = \sum_{i \leq \mathbf{n}} \sum_{j \leq \mathbf{n}} \alpha_i \alpha_j \mathbf{E} \langle X_i, X_j \rangle = \sigma^2 \sum_{i \leq \mathbf{n}} \alpha_i^2.$$

Now  $|\alpha_i| < 1$ , and vanishes if  $[(i-1)/\mathbf{n}, i/\mathbf{n}] \subset [0, \mathbf{t}]$ , or  $[(i-1)/\mathbf{n}, i/\mathbf{n}] \cap [0, \mathbf{t}] = \emptyset$ . Actually  $\alpha_i \neq 0$  if and only if  $i \in I$ , where  $I$  is defined as

$$I := \{i \leq \mathbf{n}; \forall k \in \{1, \dots, d\}, i_k \leq [n_k t_k] + 1 \text{ and} \\ \exists l \in \{1, \dots, d\}, i_l = [n_l t_l] + 1\}.$$

For any Borel set  $A \subset [0, 1]^d$  define for  $\varepsilon > 0$

$$A^\varepsilon := \{y \in \mathbb{R}^d, \exists x \in A; |x - y| < \varepsilon\}, \quad A^{-\varepsilon} := \mathbb{R}^d \setminus (\mathbb{R}^d \setminus A)^\varepsilon.$$

Put  $\varepsilon_n := \mathbf{m}(\mathbf{n})^{-1}$  and  $\beta_n(\mathbf{t}) := |[0, \mathbf{t}]^{\varepsilon_n} \setminus [0, \mathbf{t}]^{-\varepsilon_n}|$ . Then

$$\sum_{i \leq \mathbf{n}} \alpha_i^2 = \sum_{i \in I} \alpha_i^2 \leq \beta_n(\mathbf{t}),$$

and this upper bound tends to zero since the Lebesgue measure of  $[0, \mathbf{t}]^{\varepsilon_n} \setminus [0, \mathbf{t}]^{-\varepsilon_n}$  is clearly  $O(\varepsilon_n) = O(\mathbf{m}(\mathbf{n})^{-1})$ . Combined with the estimate  $P(\|Y\| > r) \leq r^{-2} \mathbf{E} \|Y\|^2$ , for any random variable  $Y$ , the theorem follows.  $\square$

This theorem coupled with Slutsky's lemma, implies then that for i.i.d.  $\mathbb{H}$ -valued random variables the limits of finite dimensional distributions of both processes  $\pi(\mathbf{n})^{-1/2}\xi_n$  and  $\pi(\mathbf{n})^{-1/2}\zeta_n$  coincide. Note that for fixed  $\mathbf{t}$

$$\zeta_n(\mathbf{t}) = \sum_{j \in J(\mathbf{n})} X_j,$$

where

$$J(\mathbf{n}) := \{\mathbf{j} \in \mathbb{N}^d : \mathbf{j}/\mathbf{n} \in [\mathbf{0}, \mathbf{t}]\}.$$

If  $X_j$  are zero mean i.i.d.  $\mathbb{H}$ -valued random variables, satisfying  $\mathbf{E} \|X_1\|^2 < \infty$  and  $G$  is the Gaussian random variable with the same covariance operator as  $X_1$ ,  $\{X_j\}$  satisfy CLT in  $\mathbb{H}$  [22, Th. 10.5]., i.e.

$$\pi(\mathbf{n})^{-1/2} \sum_{j \leq n} X_j \rightarrow G, \text{ as } \pi(\mathbf{n}) \rightarrow \infty.$$

By denoting  $l(\mathbf{n})$  the number of elements in the set  $J(\mathbf{n})$ , we then get

$$l(\mathbf{n})^{-1/2} \sum_{j \in J(\mathbf{n})} X_j \xrightarrow{\mathbb{H}} G, \text{ as } l(\mathbf{n}) \rightarrow \infty.$$

So it is easier to deal with the limits of finite-dimensional distributions of  $\zeta_n$ . Now

$$\begin{aligned} \frac{l(\mathbf{n})}{\pi(\mathbf{n})} &= \frac{1}{\pi(\mathbf{n})} \sum_{j \leq n} \mathbf{1}(\mathbf{j}/\mathbf{n} \in [\mathbf{0}, \mathbf{t}]) \\ &= P(U_n \in [\mathbf{0}, \mathbf{t}]) \xrightarrow{m(\mathbf{n}) \rightarrow \infty} |[\mathbf{0}, \mathbf{t}]| = t_1 \dots t_d, \end{aligned}$$

with  $U_n$  - random variable uniformly distributed on the points  $\mathbf{j}/\mathbf{n}$ . Recalling definition of  $\mathbb{H}$ -valued Brownian sheet we get

$$\pi(\mathbf{n})^{-1/2} \zeta_n(\mathbf{t}) \xrightarrow[m(\mathbf{n}) \rightarrow \infty]{\mathbb{H}} W(\mathbf{t}). \quad (3.2)$$

It turns out that this convergence also holds for any vector  $(\zeta_n(\mathbf{t}_1), \dots, \zeta_n(\mathbf{t}_q))$  under the same conditions.

**Theorem 18** *The convergence*

$$\pi(\mathbf{n})^{-1/2} (\zeta_n(\mathbf{t}_1), \dots, \zeta_n(\mathbf{t}_q)) \xrightarrow[m(\mathbf{n}) \rightarrow \infty]{D} (W(\mathbf{t}_1), \dots, W(\mathbf{t}_q))$$

holds for each  $q \geq 1$  and each  $\mathbf{t}_1, \dots, \mathbf{t}_q \in [0, 1]^d$ , if  $X_j$  are zero mean i.i.d.

$\mathbb{H}$ -valued random variables, satisfying  $\mathbf{E} \|X_1\|^2 < \infty$ .

This in view of theorem 17 gives the following.

**Theorem 19** *Let  $\{X_i, 1 \leq i \leq n\}, n \in \mathbb{N}$  be a collection of  $\mathbb{H}$ -valued i.i.d. random variables. Assume that  $\mathbf{E} X_1 = 0$ ,  $\mathbf{E} \|X_1\|^2 < \infty$ . Then for each  $q$ -tuple  $\mathbf{t}_1, \dots, \mathbf{t}_q \in [0, 1]^d$*

$$\pi(n)^{-1/2} (\xi_n(\mathbf{t}_1), \dots, \xi_n(\mathbf{t}_q)) \xrightarrow{D} (W(\mathbf{t}_1), \dots, W(\mathbf{t}_q)).$$

### Proof of theorem 18

For convenience write  $\tilde{\zeta}_n := \pi(n)^{-1/2} \zeta_n$ . Equip  $\mathbb{H}^q$  with product topology. Then the net  $(\tilde{\zeta}_n(\mathbf{t}_1), \dots, \tilde{\zeta}_n(\mathbf{t}_q))$  is tight in  $\mathbb{H}^q$  since the nets  $(\tilde{\zeta}_n(\mathbf{t}_i))$  are tight in  $\mathbb{H}$  due to (3.2).

Denote by  $\langle \cdot, \cdot \rangle_q$  the scalar product in  $\mathbb{H}^q$  which is defined by

$$\langle h, g \rangle_q := \sum_{i=1}^q \langle h_i, g_i \rangle, \quad h = (h_1, \dots, h_q), \quad g = (g_1, \dots, g_q) \in \mathbb{H}^q.$$

Accounting the above mentioned tightness, to prove the theorem we have to check that for each  $h \in \mathbb{H}^q$ , the following weak convergence holds

$$V_n := \left\langle (\tilde{\zeta}_n(\mathbf{t}_1), \dots, \tilde{\zeta}_n(\mathbf{t}_q)), h \right\rangle_q \xrightarrow[m(n) \rightarrow \infty]{\mathbb{R}} \left\langle (W(\mathbf{t}_1), \dots, W(\mathbf{t}_q)), h \right\rangle_q. \quad (3.3)$$

This will be done through Lindeberg theorem. The first step is to establish the convergence of the variance  $b_n := \mathbf{E} V_n^2$  using the decomposition

$$V_n = \sum_{k=1}^q \langle \tilde{\zeta}_n(\mathbf{t}_k), h_k \rangle = \pi(n)^{-1/2} \sum_{i \leq n} \sum_{k=1}^q \mathbf{1}(i/n \in [0, \mathbf{t}_k]) \langle X_i, h_k \rangle.$$

Denoting by  $\Gamma$  the covariance operator of  $X_1$ , we get

$$\begin{aligned} b_n &= \frac{1}{\pi(n)} \sum_{i \leq n} \sum_{j \leq n} \sum_{k=1}^q \sum_{l=1}^q \mathbf{1}(i/n \in [0, \mathbf{t}_k]) \mathbf{1}(j/n \in [0, \mathbf{t}_l]) \mathbf{E} (\langle X_i, h_k \rangle \langle X_j, h_l \rangle) \\ &= \sum_{k=1}^q \sum_{l=1}^q \langle \Gamma h_k, h_l \rangle \frac{1}{\pi(n)} \sum_{i \leq n} \mathbf{1}(i/n \in [0, \mathbf{t}_k] \cap [0, \mathbf{t}_l]) \\ &= \sum_{k=1}^q \sum_{l=1}^q \langle \Gamma h_k, h_l \rangle P(U_n \in [0, \mathbf{t}_k] \cap [0, \mathbf{t}_l]), \end{aligned}$$

where the discrete random variable  $U_n$  is uniformly distributed on the grid  $i/n, 1 \leq i \leq n$ . Under this form it is clear that when  $m(n)$  goes to infinity,

$b_n$  converges to  $b$  given by

$$b := \sum_{k=1}^q \sum_{l=1}^q \langle \Gamma h_k, h_l \rangle |[0, \mathbf{t}_k] \cap [0, \mathbf{t}_l]| = \mathbf{E} \left( \sum_{k=1}^q \langle W(\mathbf{t}_k), h_k \rangle \right)^2.$$

When  $b = 0$ , the convergence (3.3) is obvious. When  $b > 0$ , let us introduce the real random variables

$$Y_{n,i} := \sum_{k=1}^q \pi(\mathbf{n})^{-1/2} \mathbf{1}(i/\mathbf{n} \in [0, \mathbf{t}_k]) \langle X_i, h_k \rangle,$$

which have both zero mean and finite variance and note that  $V_n = \sum_{i \leq n} Y_{n,i}$ . To obtain (3.3) we have to check, by Lindeberg theorem, that for each  $\varepsilon > 0$ ,

$$L(\mathbf{n}) := \frac{1}{b_n} \sum_{i \leq n} \mathbf{E} \left( Y_{n,i}^2 \mathbf{1}(|Y_{n,i}| > \varepsilon b_n^{1/2}) \right) \xrightarrow{m(\mathbf{n}) \rightarrow \infty} 0. \quad (3.4)$$

Now we have

$$\begin{aligned} Y_{n,i}^2 &= \frac{1}{\pi(\mathbf{n})} \sum_{k=1}^q \sum_{l=1}^q \mathbf{1}(i/\mathbf{n} \in [0, \mathbf{t}_k]) \mathbf{1}(i/\mathbf{n} \in [0, \mathbf{t}_l]) \langle X_i, h_k \rangle \langle X_i, h_l \rangle \\ &\leq \frac{1}{\pi(\mathbf{n})} \sum_{k=1}^q \sum_{l=1}^q \|X_i\|^2 \|h_k\| \|h_l\| \\ &= \frac{1}{\pi(\mathbf{n})} \left( \sum_{k=1}^q \|h_k\| \right)^2 \|X_i\|^2 = \frac{c_h}{\pi(\mathbf{n})} \|X_i\|^2. \end{aligned}$$

Recalling that the number of terms in  $\sum_{i \leq n}$  is exactly  $\pi(\mathbf{n})$  and choosing  $m(\mathbf{n})$  large enough to have  $b_n > b/2$ , we obtain :

$$L(\mathbf{n}) \leq \frac{2}{b} \mathbf{E} \left( \|X_1\|^2 \mathbf{1} \left( \|X_1\|^2 > \frac{b\varepsilon^2}{2c_h} \pi(\mathbf{n}) \right) \right),$$

which gives (3.4) by square integrability of  $X_1$ .

### 3.1.2 Necessity

Suppose we have weak convergence of summation process  $\pi(\mathbf{n})^{-1/2} \xi_n$  to  $\mathbb{H}$ -valued Brownian sheet  $W_d$ . Since the function  $w_\alpha(\cdot, \delta)$  is continuous on  $\mathbb{H}_\alpha^o([0, 1]^d)$ , by continuous mapping theorem it follows that

$$\lim_{m(\mathbf{n}) \rightarrow \infty} P(w_\alpha((n_1 \dots n_d)^{1/2} \xi_n, \delta) > a) = P(w_\alpha(W_d, \delta) > a) \quad (3.5)$$

for each continuity point  $a$  of distribution function of the random variable  $w_\alpha(W_d, \delta)$ . Since paths of  $W_d$  lie in  $\mathbb{H}_\alpha^o([0, 1]^d)$ ,

$$P(w_\alpha(W_d, \delta) > t) \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \quad (3.6)$$

Thus oscillations of process  $\boldsymbol{\pi}(\mathbf{n})^{-1/2}\xi_{\mathbf{n}}$  should be small. Recall that  $\xi_{\mathbf{n}}(\mathbf{k}/\mathbf{n}) = S_{\mathbf{k}}$ . For arbitrary  $\delta > 0$  and  $\mathbf{n}$  such that  $|\mathbf{1}/\mathbf{n}| = m(\mathbf{n})^{-1} < \delta$ , we have

$$\begin{aligned} & P\left(w_\alpha((n_1 \dots n_d)^{-1/2}\xi_{\mathbf{n}}, \delta) > t\right) \\ & \geq P\left((n_1 \dots n_d)^{-1/2} \max_{|\frac{\mathbf{k}-\mathbf{l}}{\mathbf{n}}| = |\frac{\mathbf{1}}{\mathbf{n}}|} \frac{\|S_{\mathbf{k}} - S_{\mathbf{l}}\|}{|(\mathbf{k} - \mathbf{l})/\mathbf{n}|^\alpha} > t\right). \end{aligned}$$

On the other hand since

$$X_{\mathbf{k}} = \Delta_{k_1}^{(1)} \dots \Delta_{k_d}^{(d)} S_{\mathbf{k}},$$

we get

$$\|X_{\mathbf{k}}\| = \left\| \Delta_{k_1}^{(1)} \sum_{i \in I} \varepsilon_i S_i \right\| \leq \left\| \sum_{i \in I} \Delta_{k_1}^{(1)} S_i \right\|$$

where  $\varepsilon_i = \pm 1$  and  $I$  is some appropriate subset of  $[0, \mathbf{n}] \cap \mathbb{N}^d$  with  $2^{d-1}$  elements. Thus

$$\max_{1 \leq \mathbf{k} \leq \mathbf{n}} \frac{\|X_{\mathbf{k}}\|}{n_1^{-\alpha}} \leq 2^{d-1} \max_{|\frac{\mathbf{k}-\mathbf{l}}{\mathbf{n}}| = |\frac{\mathbf{1}}{\mathbf{n}}|} \frac{\|S_{\mathbf{k}} - S_{\mathbf{l}}\|}{|(\mathbf{k} - \mathbf{l})/\mathbf{n}|^\alpha}.$$

Now with  $p = (1/2 - \alpha)^{-1}$

$$\begin{aligned} & P\left((n_1 \dots n_d)^{-1/2} \max_{1 \leq \mathbf{k} \leq \mathbf{n}} \frac{\|X_{\mathbf{k}}\|}{n_1^{-\alpha}} > t\right) \\ & = P\left(n_1^{-1/p} n_2^{-1/2} \dots n_d^{-1/2} \max_{1 \leq \mathbf{k} \leq \mathbf{n}} \|X_{\mathbf{k}}\| > t\right) \end{aligned}$$

and we see that (3.6) gives us

$$n_1^{-1/p} n_2^{-1/2} \dots n_d^{-1/2} \max_{1 \leq \mathbf{k} \leq \mathbf{n}} \|X_{\mathbf{k}}\| \xrightarrow[m(\mathbf{n}) \rightarrow \infty]{\text{Pr}} 0. \quad (3.7)$$

If we assume that  $\{X_{\mathbf{k}}, \mathbf{k} \leq \mathbf{n}\}$  are independent and identically distributed, we have for each  $t > 0$

$$\begin{aligned} P\left(n_1^{-1/p} n_2^{-1/2} \dots n_d^{-1/2} \max_{1 \leq \mathbf{k} \leq \mathbf{n}} \|X_{\mathbf{k}}\| > t\right) &= \\ &= 1 - \left(1 - P\left(\|X_1\| > t n_1^{1/p} n_2^{1/2} \dots n_d^{1/2}\right)\right)^{n_1 n_2 \dots n_d}. \end{aligned} \quad (3.8)$$

Thus (3.7) is equivalent to

$$n_1 \dots n_d P\left(\|X_1\| > n_1^{1/p} n_2^{1/2} \dots n_d^{1/2}\right) \xrightarrow{\mathbf{m}(\mathbf{n}) \rightarrow \infty} 0. \quad (3.9)$$

Note that (3.7) is as well equivalent to

$$\boldsymbol{\pi}(\mathbf{n}) P\left(\|X_1\| > n_m^{1/p} \boldsymbol{\pi}(\mathbf{n}_{-m})^{1/2}\right) \xrightarrow{\mathbf{m}(\mathbf{n}) \rightarrow \infty} 0, \quad (3.10)$$

for any  $m \in \{2, \dots, d\}$ .

### 3.1.3 Tightness

In this subsection we prove the tightness of summation process  $\boldsymbol{\pi}(\mathbf{n})^{-1/2} \xi_{\mathbf{n}}$  in the space  $\mathbb{H}_{\alpha}^0([0, 1]^d)$  for the mean-zero i.i.d. collection of random variables  $\{X_{\mathbf{j}}, \mathbf{j} \in \mathbb{N}^d\}$ . We use tightness criteria, theorem 5, so we have to check two conditions. The first about asymptotic tightness of the net  $\xi_{\mathbf{n}}$  at each point  $\mathbf{t}$  readily follows from finite-dimensional convergence, which requires that  $\mathbf{E} \|X_1\|^2 < \infty$ .

Recalling the estimate (2.34) from the section 2.1.3 and the relation (1.4) from the section 1.2.3 it follows that

$$\lim_{J \rightarrow \infty} \limsup_{\mathbf{m}(\mathbf{n}) \rightarrow \infty} P\left(\sup_{j \geq J} 2^{\alpha j} \max_{v \in V_j} |\lambda_{j,v}(\boldsymbol{\pi}(\mathbf{n})^{-1/2} \xi_{\mathbf{n}})| > \varepsilon\right) = 0$$

holds if

$$n_m^{\alpha} \boldsymbol{\pi}(\mathbf{n})^{-1/2} Z_{\mathbf{n}}^{(m)} \xrightarrow[\mathbf{m}(\mathbf{n}) \rightarrow \infty]{\text{Pr}} 0 \quad (3.11)$$

and

$$\lim_{J \rightarrow \infty} \limsup_{\mathbf{m}(\mathbf{n}) \rightarrow \infty} P\left(\max_{J \leq j \leq \log n_1} 2^{\alpha j} \boldsymbol{\pi}(\mathbf{n})^{-1/2} \max_{0 \leq k < 2^j} \psi_{\mathbf{n}}(t_{k+1}, t_k) > \varepsilon'\right) = 0 \quad (3.12)$$

hold for each  $m = 1, \dots, d$ . It turns out that condition

$$\pi(\mathbf{n})P\left(\|X_1\| > n_m^{1/p}\pi(\mathbf{n}_{-m})^{1/2}\right) \xrightarrow{m(\mathbf{n}) \rightarrow \infty} 0, \quad m = 1, \dots, d, \quad (3.13)$$

is sufficient for (3.11) and (3.12) to hold. Since (3.13) ensures that  $\mathbf{E}\|X_1\|^2 < \infty$ , then condition (3.13) alone is sufficient for tightness of summation process  $\pi(\mathbf{n})^{1/2}\xi_{\mathbf{n}}$ .

### Proof of (3.11)

We prove (3.11) for  $m = 1$ , since the proof is the same for other  $m$ . For this reason we drop superscript <sup>(1)</sup> from  $Z_{\mathbf{n}}$ . Note first that really

$$Z_{\mathbf{n}} = \max_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{n}} \left\| \sum_{i_2=1}^{k_2} \dots \sum_{i_d=1}^{k_d} X_{(k_1, i_2, \dots, i_d)} \right\|.$$

Fix  $\varepsilon > 0$  and associate to any  $\delta \in (0, 1)$  the truncated random variables  $\widetilde{X}_j$  and  $X'_j$  defined as

$$\begin{aligned} \widetilde{X}_j &:= X_j \mathbf{1} \left( \|X_j\| \leq \delta n_1^{1/p} (n_2 \dots n_d)^{1/2} \right), \\ X'_j &:= \widetilde{X}_j - \mathbf{E} \widetilde{X}_j, \quad \mathbf{1} \leq j \leq \mathbf{n}. \end{aligned}$$

Substituting  $X_j$  by  $\widetilde{X}_j$ , respectively  $X'_j$ , in the definition of  $Z_{\mathbf{n}}$  we obtain  $\widetilde{Z}_{\mathbf{n}}$ , respectively  $Z'_{\mathbf{n}}$ . Introducing the complementary events

$$E_{\mathbf{n}} := \left\{ \forall \mathbf{k} \leq \mathbf{n}, \|X_{\mathbf{k}}\| \leq \delta n_1^{1/p} (n_2 \dots n_d)^{1/2} \right\}, \quad E_{\mathbf{n}}^c := \Omega \setminus E_{\mathbf{n}},$$

we have

$$P(Z_{\mathbf{n}} > \varepsilon n_1^{1/p} (n_2 \dots n_d)^{1/2}) \leq P(\{Z_{\mathbf{n}} > \varepsilon n_1^{1/p} (n_2 \dots n_d)^{1/2}\} \cap E_{\mathbf{n}}) + P(E_{\mathbf{n}}^c).$$

Clearly  $Z_{\mathbf{n}} = \widetilde{Z}_{\mathbf{n}}$  on the event  $E_{\mathbf{n}}$ . By identical distribution of the  $X_{\mathbf{k}}$ 's,

$$P(E_{\mathbf{n}}^c) \leq n_1 \dots n_d P(\|X_1\| > \delta n_1^{1/p} (n_2 \dots n_d)^{1/2})$$

and this upper bound goes to zero when  $m(\mathbf{n})$  goes to infinity by condition (3.13). This leads to

$$\begin{aligned} & \limsup_{m(\mathbf{n}) \rightarrow \infty} P(Z_{\mathbf{n}} > \varepsilon n_1^{1/p} (n_2 \dots n_d)^{1/2}) \\ & \leq \limsup_{m(\mathbf{n}) \rightarrow \infty} P(\tilde{Z}_{\mathbf{n}} > \varepsilon n_1^{1/p} (n_2 \dots n_d)^{1/2}). \end{aligned} \quad (3.14)$$

Because  $n_1^{-1/p} (n_2 \dots n_d)^{1/2} \|\mathbf{E} \tilde{X}_1\| \rightarrow 0$  as  $m(\mathbf{n}) \rightarrow \infty$  by lemma 23, the right-hand side of (3.14) does not exceed

$$\limsup_{m(\mathbf{n}) \rightarrow \infty} P(n_1^{-1/p} (n_2 \dots n_d)^{-1/2} Z'_{\mathbf{n}} > \varepsilon).$$

Using the extension of Doob inequality (1.12), we obtain with  $q > p$

$$\begin{aligned} & P(n_1^{-1/p} (n_2 \dots n_d)^{-1/2} Z'_{\mathbf{n}} > \varepsilon) \\ & \leq n_1 P \left( \max_{\mathbf{1}_{2:d} \leq \mathbf{k}_{2:d} \leq \mathbf{n}_{2:d}} \left\| \sum_{i_{2:d}=\mathbf{1}_{2:d}}^{\mathbf{k}_{2:d}} X'_{(1,i_2,\dots,i_d)} \right\| > \varepsilon n_1^{1/p} (n_2 \dots n_d)^{1/2} \right) \\ & \leq \varepsilon^{-q} n_1^{1-q/p} (n_2 \dots n_d)^{-q/2} \mathbf{E} \left\| \sum_{i_{2:d}=\mathbf{1}_{2:d}}^{\mathbf{n}_{2:d}} X'_{(1,i_2,\dots,i_d)} \right\|^q. \end{aligned}$$

Applying Rosenthal inequality (1.10) together with the estimates (3.34), (3.35), provided in subsection 3.1.5 below, we obtain

$$\begin{aligned} & P(n_1^{-1/p} (n_2 \dots n_d)^{-1/2} Z'_{\mathbf{n}} > \varepsilon) \\ & \leq \varepsilon^{-q} n_1^{1-q/p} (n_2 \dots n_d)^{-q/2} C_q \left( (n_2 \dots n_d)^{q/2} (\mathbf{E} \|X'_1\|^2)^{q/2} \right. \\ & \quad \left. + n_2 \dots n_d \mathbf{E} \|X'_1\|^q \right) \\ & \leq C_q \varepsilon^{-q} \left( n_1^{1-q/p} (\mathbf{E} \|X_1\|^2)^{q/2} + \frac{2^{q+1} c_{p,m}}{q-p} \delta^{q-p} \right). \end{aligned}$$

Combined with (3.14) this gives

$$\limsup_{m(\mathbf{n}) \rightarrow \infty} P(n_1^{-1/p} (n_2 \dots n_d)^{-1/2} Z_{\mathbf{n}} > \varepsilon) \leq c \delta^{q-p},$$

where the constant  $c$  depends on  $\varepsilon$ ,  $p$  and  $q$ . Since  $q > p$  and  $\delta$  may be chosen arbitrarily small in  $(0, 1)$ , the convergence (3.11) follows.

**Proof of (3.12)**

Again we give the proof for  $m = 1$  and subsequently drop the superscript. For notational simplification, let us agree to denote by  $\varepsilon'$  the successive splittings of  $\varepsilon$ , i.e.  $\varepsilon' = c\varepsilon$  where the constant  $c \in (0, 1)$  may decrease from one formula to following one. Notations  $\tilde{\psi}_{\mathbf{n}}(t_{k+1}, t_k)$  and  $\psi'_{\mathbf{n}}(t_{k+1}, t_k)$  mean that  $X_j$  are substituted by  $\tilde{X}_j$  and  $X'_j$  respectively in the definition of  $\psi_{\mathbf{n}}(t_{k+1}, t_k)$ . Accordingly we introduce the notations  $\tilde{P}(J, \mathbf{n}; \varepsilon')$  and  $P'(J, \mathbf{n}; \varepsilon')$  where

$$P(J, \mathbf{n}; \varepsilon) = P\left(\max_{J \leq j \leq \log n_1} 2^{\alpha j} \boldsymbol{\pi}(\mathbf{n})^{-1/2} \max_{0 \leq k < 2^j} \psi_{\mathbf{n}}(t_{k+1}, t_k) > \varepsilon'\right) \quad (3.15)$$

Splitting  $\Omega$  in complementary events

$$E_n := \left\{ \forall \mathbf{k} \leq \mathbf{n}, \|X_{\mathbf{k}}\| \leq \delta n_1^{1/p} (n_2 \dots n_d)^{1/2} \right\}, \quad E_n^c := \Omega \setminus E_n,$$

like in previous subsection we obtain

$$P(J, \mathbf{n}; \varepsilon') \leq \tilde{P}(J, \mathbf{n}; \varepsilon) + n_1 \dots n_d P(\|X_1\| \geq \delta n_1^{1/p} (n_2 \dots n_d)^{1/2}).$$

Then (3.12) is reduced by condition (3.13) to

$$\lim_{J \rightarrow \infty} \limsup_{\mathbf{m}(\mathbf{n}) \rightarrow \infty} \tilde{P}(J, \mathbf{n}; \varepsilon') = 0. \quad (3.16)$$

The number of variables  $\tilde{X}_{\mathbf{k}}$  to be centered in the sum  $\tilde{\psi}_{\mathbf{n}}(t_{k+1}, t_k)$  is at most  $n_1(t_{k+1} - t_k)n_2 \dots n_d \leq n_1 2^{-J} n_2 \dots n_d$  and (3.32) yields

$$\begin{aligned} \max_{J \leq j \leq \log n_1} 2^{\alpha j} (n_1 \dots n_d)^{-1/2} \|\mathbf{E} \tilde{X}_1\| &\leq n_1^{\alpha-1/2} (2\delta^{1-p} c_{p,m}) n_1^{1/p-1} (n_2 \dots n_d)^{-1} \\ &= 2\delta^{1-p} c_{p,m} (n_1 \dots n_d)^{-1}. \end{aligned}$$

Therefore

$$\limsup_{\mathbf{m}(\mathbf{n}) \rightarrow \infty} \max_{J \leq j \leq \log n_1} 2^{\alpha j} (n_1 \dots n_d)^{-1/2} n_1 2^{-J} n_2 \dots n_d \|\mathbf{E} \tilde{X}_1\| \leq \delta^{1-p} c_p 2^{-J+1}.$$

This upper bound going to zero when  $J$  goes to infinity, (3.16) is reduced to

$$\lim_{J \rightarrow \infty} \limsup_{\mathbf{m}(\mathbf{n}) \rightarrow \infty} P'(J, \mathbf{n}; \varepsilon') = 0. \quad (3.17)$$

We have with  $q > p$

$$\begin{aligned} P'(J, \mathbf{n}; \varepsilon') &\leq \sum_{j=J}^{\log n_1} P\left(2^{\alpha j}(n_1 \dots n_d)^{-1/2} \max_{0 \leq k < 2^j} \psi'_n(t_{k+1}, t_k) > \varepsilon'\right) \\ &\leq \sum_{j=J}^{\log n_1} 2^{q\alpha j} (n_1 \dots n_d)^{-q/2} \varepsilon'^{-q} 2^j \mathbf{E} \psi'_n(t_{k+1}, t_k)^q. \end{aligned} \quad (3.18)$$

Denote  $u_k = \lfloor n_1 t_k \rfloor$  and observe that  $u_{k+1} - u_k \leq n_1 2^{-j}$ . By (1.12),

$$\mathbf{E} \psi'_n(t_{k+1}, t_k)^q \leq \mathbf{E} \left\| \sum_{i_1=1+u_k}^{u_{k+1}} \sum_{i_{2:d}=\mathbf{1}_{2:d}}^{n_{2:d}} X'_i \right\|^q.$$

Estimating this last  $q$ -moment by Rosenthal inequality (1.10) with a number of summands  $N \leq (n_1 2^{-j}) n_2 \dots n_d$ , we obtain

$$\begin{aligned} \mathbf{E} \psi'_n(t_{k+1}, t_k)^q &\leq C_q \left( (n_1 2^{-j})^{q/2} (n_2 \dots n_d)^{q/2} \mathbf{E} \|X'_1\|^2 + n_1 2^{-j} n_2 \dots n_d \mathbf{E} \|X'_1\|^q \right) \\ &\leq C_q \mathbf{E} \|X_1\|^2 2^{-jq/2} (n_1 \dots n_d)^{q/2} \\ &\quad + \frac{2^{q+1} C_q C_{p,m}}{q-p} \delta^{q-p} 2^{-j} n_1^{q/p} (n_2 \dots n_d)^{q/2}. \end{aligned}$$

Reporting this estimate into (3.18) we obtain

$$P'(J, \mathbf{n}; \varepsilon') \leq \Sigma_1(J, \mathbf{n}; \varepsilon') + \Sigma_2(J, \mathbf{n}; \varepsilon')$$

with  $\Sigma_1$  and  $\Sigma_2$  explicited and bounded as follows. First

$$\begin{aligned} \Sigma_1(J, \mathbf{n}; \varepsilon') &:= \frac{C_q}{\varepsilon'^q} \mathbf{E} \|X_1\|^2 \sum_{J \leq j \leq \log n_1} 2^{(1+q(\alpha-1/2))j} \\ &\leq \frac{C_q}{\varepsilon'^q} \mathbf{E} \|X_1\|^2 \sum_{j=J}^{\infty} 2^{-(q/p-1)j} \\ &= \frac{C_q}{\varepsilon'^q} \mathbf{E} \|X_1\|^2 \frac{2^{-(q/p-1)J}}{1 - 2^{-(q/p-1)}}. \end{aligned}$$

Hence

$$\lim_{J \rightarrow \infty} \limsup_{m(\mathbf{n}) \rightarrow \infty} \Sigma_1(J, \mathbf{n}; \varepsilon') = 0.$$

Next

$$\begin{aligned}\Sigma_2(J, \mathbf{n}; \varepsilon') &:= \frac{2^{q+1}C_q c_{p,m}}{(q-p)\varepsilon'^q} \delta^{q-p} n_1^{-q\alpha} \sum_{J \leq j \leq \log n_1} 2^{jq\alpha} \\ &\leq \frac{2^{q+1}C_q c_{p,m}}{(q-p)\varepsilon'^q} \delta^{q-p} n_1^{-q\alpha} \frac{n_1^{q\alpha}}{2^{q\alpha} - 1}.\end{aligned}$$

Noting that  $m = m(\mathbf{n})$  and  $\limsup_{m \rightarrow \infty} c_{p,m} = c_p$ , we obtain

$$\limsup_{m(\mathbf{n}) \rightarrow \infty} \Sigma_2(J, \mathbf{n}; \varepsilon') \leq \frac{2^{q+1}C_q c_p}{(q-p)(2^{q\alpha} - 1)\varepsilon'^q} \delta^{q-p}.$$

Recalling (3.15) and summing up all the successive reductions leads to

$$\limsup_{J \rightarrow \infty} \limsup_{m(\mathbf{n}) \rightarrow \infty} P(J, \mathbf{n}; \varepsilon) \leq \frac{2^{q+1}C_q c_p}{(q-p)(2^{q\alpha} - 1)\varepsilon'^q} \delta^{q-p}.$$

Since  $P(J, \mathbf{n}; \varepsilon)$  does not depend on  $\delta$  which may be chosen arbitrarily small, the left-hand side is null and this gives (3.12).

### 3.1.4 Corollaries

We state now the invariance principle in the space  $\mathbb{H}_\alpha^o([0, 1]^d)$ .

**Theorem 20** *For  $0 < \alpha < 1/2$ , set  $p = p(\alpha) := 1/(1/2 - \alpha)$ . For  $d \geq 2$ , let  $\{X_i; \mathbf{i} \in \mathbb{N}^d, \mathbf{i} \geq \mathbf{1}\}$  be an i.i.d. collection of square integrable centered random elements in the separable Hilbert space  $\mathbb{H}$  and  $\xi_n$  be the summation process defined by*

$$\xi_n(\mathbf{t}) = \sum_{\mathbf{i} \leq \mathbf{n}} \pi(\mathbf{n}) \left| \left[ \frac{\mathbf{i} - \mathbf{1}}{\mathbf{n}}, \frac{\mathbf{i}}{\mathbf{n}} \right] \cap [0, \mathbf{t}] \right| X_i. \quad (3.19)$$

*Let  $W$  be a  $\mathbb{H}$ -valued Brownian sheet with the same covariance operator as  $X_1$ . Then the convergence*

$$\pi(\mathbf{n})^{-1/2} \xi_n \xrightarrow[m(\mathbf{n}) \rightarrow \infty]{\mathbb{H}_\alpha^o([0, 1]^d)} W \quad (3.20)$$

*holds if and only if*

$$\pi(\mathbf{n}) P\left(\|X_1\| > n_m^{1/p} \pi(\mathbf{n}_{-m})^{1/2}\right) \xrightarrow[m(\mathbf{n}) \rightarrow \infty]{} 0, \quad (3.21)$$

*for  $m = 1, \dots, d$ .*

*Proof.* It is really nothing left to prove, since necessity is proved in subsection 3.1.2, the convergence of finite-dimensional distributions follow since (3.21) ensures that  $\mathbf{E} \|X_1\|^2 < \infty$  and tightness is proved in subsection 3.1.3.  $\square$

Though condition (3.21) looks rather technical it turns out that it is equivalent to the finiteness of the weak  $p$ -moment of  $X_1$ , i.e.

$$\sup_{t>0} t^p P(\|X_1\| > t) < \infty. \quad (3.22)$$

We prove this for  $m = 1$  as the proof is the same for all  $m$ . Note that (3.21) is equivalent to

$$v_1^p (v_2 \cdots v_d)^2 P(\|X_1\| > v_1 v_2 \cdots v_d) \xrightarrow{m(\mathbf{v}) \rightarrow \infty} 0 \quad (3.23)$$

and in return (3.23) is equivalent to the convergence

$$F(m) \xrightarrow{m \rightarrow \infty} 0, \quad (3.24)$$

where

$$F(m) := \sup_{m(\mathbf{v}) \geq m} v_1^p (v_2 \cdots v_d)^2 P(\|X_1\| > v_1 v_2 \cdots v_d).$$

Now introducing the function  $g(t) := P(\|X_1\| > t)$  and the sets

$$H_{t,m} := \{\mathbf{v} \in \mathbb{R}^d; \mathbf{v} \geq m, v_1 v_2 \cdots v_d = t\},$$

we have

$$F(m) = \sup_{t \geq m^d} \sup_{\mathbf{v} \in H_{t,m}} v_1^{p-2} t^2 g(t) = \sup_{t \geq m^d} t^2 g(t) \sup_{\mathbf{v} \in H_{t,m}} v_1^{p-2}.$$

When  $t \geq m^d$ ,  $H_{t,m}$  is non empty and on this set,  $v_1 = t(v_2 \cdots v_d)^{-1}$  is maximal for  $v_2 = \cdots = v_d = m$ , so

$$t^2 g(t) \sup_{\mathbf{v} \in H_{t,m}} v_1^{p-2} = t^p g(t) m^{-(d-1)(p-2)}.$$

Finally

$$F(m) = m^{-(d-1)(p-2)} \sup_{t \geq m^d} t^p g(t).$$

Recalling that  $d > 1$  and  $p > 2$ , this reduces the convergence (3.24) to the finiteness of  $\sup_{t \geq m_0^d} t^p g(t)$  for some  $m_0 > 0$ . As  $t^p g(t)$  is bounded on any interval  $[0, a]$  for  $a < \infty$ , this finiteness is equivalent to (3.22). Thus we proved the following theorem.

**Theorem 21** *The convergence*

$$\pi(\mathbf{n})^{-1/2} \xi_{\mathbf{n}} \xrightarrow[\mathbf{m}(\mathbf{n}) \rightarrow \infty]{\mathbb{H}_{\alpha}^o([0,1]^d)} W$$

holds if and only if

$$\sup_{t>0} t^{p(\alpha)} P(\|X_1\| > t) < \infty, \quad p(\alpha) = (1/2 - \alpha)^{-1}.$$

As condition (3.22) is weaker than  $\mathbf{E} \|X_1\|^p < \infty$ , then theorem 20 improves when  $\mathbb{H} = \mathbb{R}$ , Erickson's [13] result for  $\mathcal{Q}_d$ :

$$(n_1 \cdots n_d)^{-1/2} \xi_{\mathbf{n}} \xrightarrow[\mathbf{m}(\mathbf{n}) \rightarrow \infty]{\mathbb{H}_{\alpha}^o([0,1]^d)} W,$$

if  $0 < \alpha < 1/2$  and  $\mathbf{E} |X_1|^q < \infty$ , where  $q > dp(\alpha)$ .

Considering the convergence of random fields  $(\xi_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}^d)$  along the fixed path  $\mathbf{n} = (n, \dots, n) \in \mathbb{N}^d$ ,  $n \in \mathbb{N}$  we obtain the following result.

**Theorem 22** *The convergence*

$$n^{-d/2} \xi_{(n, \dots, n)} \xrightarrow[n \rightarrow \infty]{\mathbb{H}_{\alpha}^o([0,1]^d)} W \quad (3.25)$$

holds if and only if

$$\lim_{t \rightarrow \infty} t^{\frac{2d}{d-2\alpha}} P(\|X_1\| > t) = 0. \quad (3.26)$$

*Proof.* Looking back at the proofs in previous subsections and having in mind the extra assumption that  $n_1 = n_2 = \dots = n_d = n$ , it should be clear that the weak  $\mathbb{H}_{\alpha}^o([0,1]^d)$  convergence of  $n^{-d/2} \xi_{(n, \dots, n)}$  to  $W$  is equivalent to the condition obtained by reporting this equality of the  $n_i$ 's in (3.9), namely to

$$n^d P(\|X_1\| > n^{1/p+(d-1)/2}) \xrightarrow[n \rightarrow \infty]{} 0. \quad (3.27)$$

It is easily checked that in (3.27) the integer  $n$  can be replaced by a positive real number  $s$  and then putting  $t = s^{1/p+(d-1)/2}$ , we obtain the equivalence of (3.27) with

$$\lim_{t \rightarrow \infty} t^{\frac{2pd}{2+p(d-1)}} P(\|X_1\| > t) = 0. \quad (3.28)$$

Finally recalling that  $p = p(\alpha) = 2/(1 - 2\alpha)$ , we get

$$\frac{2pd}{2 + p(d-1)} = \frac{2d}{d - 2\alpha},$$

which reported in (3.28) gives (3.26) and completes the proof.  $\square$

Since  $2d/(d-2\alpha) < 2d/(d-1)$  we see that  $\mathbf{E} \|X_1\|^{2d/(d-1)} < \infty$  yields (3.26). In particular  $\mathbf{E} \|X_1\|^4 < \infty$  gives the convergence (3.25) for any  $d \geq 2$  and any  $0 < \alpha < 1/2$ . This contrasts with the corresponding result for Hölder convergence of the usual Donsker-Prokhorov polygonal line processes where necessarily  $\mathbf{E} |X_1|^q < \infty$  for any  $q < p(\alpha)$  as follows from (24).

Of course, Theorem 22 is only a striking special case and similar results can be obtained adapting the proof of Theorem 20 for summation processes with index going to infinity along some various paths or surfaces.

As passing from  $n$  to  $n+1$  brings  $O(n^{d-1})$  new summands in the summation process of Theorem 22, one may be tempted to look for similar weakening of the assumption in the Hölderian FCLT for  $d=1$ , when restricting for subsequences. In fact even so, the situation is quite different: it is easy to see that for any increasing sequence of integers  $n_k$  such that  $\sup_{k \geq 1} n_{k+1}/n_k < \infty$ , the convergence to zero of  $n_k^{p(\alpha)} P(|X_1| > n_k)$  when  $k$  tends to infinity implies (24). As  $n_k^{p(\alpha)} P(|X_1| > n_k) = o(1)$  is a necessary condition for  $(\xi_{n_k})_{k \geq 1}$  to satisfy the FCLT in  $H_\alpha([0, 1]^d)$  when  $d=1$ , there is no hope to obtain this FCLT for  $(\xi_{n_k})_{k \geq 1}$  under some condition weaker than (24).

### 3.1.5 Truncated variables

In this subsection we complete the technical details about the estimates of moment of truncated variables used above. Such estimates are obtained under the assumption:

$$n_1 \cdots n_d P\left(\|X_1\| > n_1^{1/p} n_2^{1/2} \cdots n_d^{1/2}\right) \xrightarrow{m(n) \rightarrow \infty} 0. \quad (3.29)$$

Let  $\delta \in (0, 1)$  be an arbitrary number. Define

$$\widetilde{X}_j := X_j \mathbf{1}\left(\|X_j\| \leq \delta n_1^{1/p} (n_2 \cdots n_d)^{1/2}\right), \quad (3.30)$$

$$X'_j := \widetilde{X}_j - \mathbf{E} \widetilde{X}_j, \quad 1 \leq j \leq n. \quad (3.31)$$

Though we give the proof for (3.30) definition of truncated variable, the estimates hold for any permutation of indexes  $1, \dots, d$  in (3.30) combined with the same permutation in (3.29).

Denote for  $m \geq 0$

$$\begin{aligned} c(m) &:= \sup_{u \geq m} \sup_{v_2, \dots, v_d \geq m} uv_2 \cdots v_d P(\|X_1\| > u^{1/p} (v_2 \cdots v_d)^{1/2}) \\ c_p &:= \sup_{t \geq 0} t^{d/(d/2-\alpha)} P(\|X_1\| > t). \end{aligned}$$

Evidently condition (3.29) yields  $c(m) \rightarrow 0$  as  $m \rightarrow \infty$  and  $c_p < \infty$ . Set

$$c_{p,m} := \max\{c_p; c(m)\}.$$

**Lemma 23** *With  $m = m(\mathbf{n})$  and any  $q > p$*

$$\|\mathbf{E} \widetilde{X}_1\| \leq 2\delta^{1-p} c_{p,m} n_1^{1/p-1} (n_2 \dots n_d)^{-1/2}; \quad (3.32)$$

$$\mathbf{E} \|\widetilde{X}_1\|^q \leq \frac{2c_{p,m}}{q-p} \delta^{q-p} n_1^{q/p-1} (n_2 \dots n_d)^{q/2-1}; \quad (3.33)$$

$$\mathbf{E} \|X'_1\|^2 \leq \mathbf{E} \|X_1\|^2; \quad (3.34)$$

$$\mathbf{E} \|X'_1\|^q \leq \frac{2^{q+1} c_{p,m}}{q-p} \delta^{q-p} n_1^{q/p-1} (n_2 \dots n_d)^{q/2-1}. \quad (3.35)$$

*Proof.* To check (3.32), we observe first that since  $\mathbf{E} X_1 = 0$ ,

$$\begin{aligned} \|\mathbf{E} \widetilde{X}_1\| &= \|\mathbf{E} X_1 - \mathbf{E} X_1 \mathbf{1}(\|X_1\| > \delta n_1^{1/p} (n_2 \dots n_d)^{1/2})\| \\ &\leq \int_{\delta n_1^{1/p} (n_2 \dots n_d)^{1/2}}^{\infty} P(\|X_1\| > t) dt \\ &\quad + \delta n_1^{1/p} (n_2 \dots n_d)^{1/2} P(\|X_1\| > \delta n_1^{1/p} (n_2 \dots n_d)^{1/2}). \end{aligned}$$

Next we have

$$\begin{aligned} &\int_{\delta n_1^{1/p} (n_2 \dots n_d)^{1/2}}^{\infty} P(\|X_1\| > t) dt \\ &= \delta n_1^{1/p-1} (n_3 \dots n_d)^{-1/2} \int_{n_2^{1/2}}^{\infty} v^2 n_1 n_3 \dots n_d \\ &\quad P(\|X_1\| > \delta v n_1^{1/p} (n_3 \dots n_d)^{1/2}) \frac{dv}{v^2} \\ &\leq \delta n_1^{1/p-1} (n_3 \dots n_d)^{-1/2} b(m, \delta) \int_{n_2^{1/2}}^{\infty} v^{-2} dv \\ &\leq \delta b(m, \delta) n_1^{1/p-1} (n_2 \dots n_d)^{-1/2}, \end{aligned}$$

where

$$b(m, \delta) := \sup_{u \geq m} \sup_{v_{2:d} \geq m} u v_2 \dots v_d P(\|X_1\| > \delta u^{1/p} (v_2 \dots v_d)^{1/2}).$$

We complete the proof of (3.32) noting that

$$\begin{aligned}
b(m; \delta) &= \delta^{-p} \sup_{u \geq \delta^p m} \sup_{v_{2:d} \geq m} uv_2 \dots v_d P(\|X_1\| > u^{1/p} (v_2 \dots v_d)^{1/2}) \\
&= \delta^{-p} \max \left\{ \sup_{m \geq u \geq \delta^p m} \sup_{v_{2:d} \geq m} uv_2 \dots v_d P(\|X_1\| > u^{1/p} (v_2 \dots v_d)^{1/2}); \right. \\
&\quad \left. \sup_{u \geq m} \sup_{v_{2:d} \geq m} uv_2 \dots v_d P(\|X_1\| > u^{1/p} (v_2 \dots v_d)^{1/2}) \right\} \\
&\leq \delta^{-p} c_{p,m}, \tag{3.36}
\end{aligned}$$

since

$$\begin{aligned}
&\sup_{u \leq m} \sup_{v_{2:d} \geq m} uv_2 \dots v_d P(\|X_1\| > u^{1/p} (v_2 \dots v_d)^{1/2}) \\
&\leq \sup_{u \leq m} \sup_{v_{2:d} \geq m} uv_2 \dots v_d c_p (u^{1/p} (v_2 \dots v_d)^{1/2})^{-d/(d/2-\alpha)} \\
&= c_p \sup_{u \leq m} u^{2\alpha(d-1)/(d-2\alpha)} \sup_{v_{2:d} \geq m} (v_2 \dots v_d)^{-2\alpha/(d-2\alpha)} = c_p.
\end{aligned}$$

Next we have

$$\begin{aligned}
\mathbf{E} \|\widetilde{X}_1\|^q &\leq \int_0^{\delta n_1^{1/p} (n_2 \dots n_d)^{1/2}} t^{q-1} P(\|X_1\| > t) dt \\
&= \int_0^{\delta (n_2 \dots n_d)^{1/2}} t^{q-1} P(\|X_1\| > t) dt \\
&\quad + \int_{\delta (n_2 \dots n_d)^{1/2}}^{\delta n_1^{1/p} (n_2 \dots n_d)^{1/2}} t^{q-1} P(\|X_1\| > t) dt.
\end{aligned}$$

By Chebyshev inequality  $P(\|X_1\| > t) \leq t^{-2}$ , hence the first integral does not exceed  $(q-2)^{-1} \delta^{q-2} (n_2 \dots n_d)^{q/2-1}$ . As  $\int_1^{n_1^{1/p}} \leq n_1^{q/p-1}$ , the second integral does not exceed

$$\begin{aligned}
&\delta^q (n_2 \dots n_d)^{q/2-1} \int_1^{n_1^{1/p}} n_2 \dots n_d u^p P(\|X_1\| > \delta u (n_2 \dots n_d)^{1/2}) u^{q-p-1} du \\
&\leq \delta^q (n_2 \dots n_d)^{q/2-1} \sup_{v_{2:d} \geq m} \sup_{1 \leq u \leq n_1} uv_2 \dots v_d P(\|X_1\| > \delta u^{1/p} (v_2 \dots v_d)^{1/2}) n_1^{q/p-1} \\
&\leq \frac{1}{q-p} \max\{b'(m, \delta); b(m, \delta)\} \delta^q n_1^{q/p-1} (n_2 \dots n_d)^{q/2-1},
\end{aligned}$$

where

$$\begin{aligned} b'(m, \delta) &:= \sup_{v_2:d \geq m} \sup_{1 \leq u \leq m} uv_2 \dots v_d P(\|X_1\| > \delta u^{1/p} (v_2 \dots v_d)^{1/2}) \\ &\leq \delta^{-2d/(d/2-\alpha)} c_p \leq \delta^{-p} c_p, \end{aligned}$$

recalling that  $0 < \delta < 1$  and  $p = (1/2 - \alpha)^{-1}$ . Accounting (3.36) inequality (3.33) now follows.

To check (3.34), let us denote by  $(e_k, k \in \mathbb{N})$  some orthonormal basis of the separable Hilbert space  $\mathbb{H}$ . Then we have

$$\|X'_1\|^2 = \sum_{k=0}^{\infty} \left| \langle \widetilde{X}_1 - \mathbf{E} \widetilde{X}_1, e_k \rangle \right|^2 = \sum_{k=0}^{\infty} \left| \langle \widetilde{X}_1, e_k \rangle - \mathbf{E} \langle \widetilde{X}_1, e_k \rangle \right|^2,$$

whence

$$\begin{aligned} \mathbf{E} \|X'_1\|^2 &= \sum_{k=0}^{\infty} \text{Var}(\langle \widetilde{X}_1, e_k \rangle) \leq \sum_{k=0}^{\infty} \mathbf{E} \left| \langle \widetilde{X}_1, e_k \rangle \right|^2 \\ &= \mathbf{E} \sum_{k=0}^{\infty} \left| \langle \widetilde{X}_1, e_k \rangle \right|^2 = \mathbf{E} \|\widetilde{X}_1\|^2 \leq \mathbf{E} \|X_1\|^2, \end{aligned}$$

which gives (3.34).

Finally we note that (3.35) is obviously obtained from (3.33) since the convexity inequality  $\|X'_1\|^q \leq 2^{q-1} \|\widetilde{X}_1\|^q + 2^{q-1} \|\mathbf{E} \widetilde{X}_1\|^q$  together with  $\mathbf{E} \|\widetilde{X}_1\| \leq (\mathbf{E} \|\widetilde{X}_1\|^q)^{1/q}$  gives  $\mathbf{E} \|X'_1\|^q \leq 2^q \mathbf{E} \|\widetilde{X}_1\|^q$ .  $\square$

## 3.2 Triangular array

### 3.2.1 Finite dimensional distributions

In this subsection we find the limits of finite-dimensional distributions of process  $\Xi_n$ . We show that the convergence to Brownian sheet is only a special case. In general case the limiting process is Gaussian, if the limit exists, but that is not always the case. As in the i.i.d. case we show that it is more convenient to analyze the jump version  $Z_n$  of process  $\Xi_n$ . Then we give some examples for which convergence to the Brownian sheet fails. Finally we give the conditions and assumptions, under which the finite-dimensional distributions converge to some Gaussian process.

Recall definitions  $\mathbf{B}(\mathbf{k})$  and  $b_l(k_l)$  from equations (2.41) and (2.42) in the

section 2.2.1. Define then the jump process as

$$Z_n(\mathbf{t}) = \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \mathbf{1}(\mathbf{B}(\mathbf{k}) \in [0, \mathbf{t}]) X_{n,\mathbf{k}}.$$

The following result holds.

**Lemma 24** *Assume*

$$\max_{1 \leq l \leq d} \max_{1 \leq k_l \leq k_n^l} \Delta b_l(k_l) \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty. \quad (3.37)$$

*Then*

$$\mathbf{E} |\Xi_n(\mathbf{t}) - Z_n(\mathbf{t})|^2 \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty \quad (3.38)$$

*and subsequently*

$$|\Xi_n(\mathbf{t}) - Z_n(\mathbf{t})| \xrightarrow{P} 0, \text{ as } m(\mathbf{n}) \rightarrow \infty.$$

*Proof.* For each  $\mathbf{t}$  we have

$$|\Xi_n(\mathbf{t}) - Z_n(\mathbf{t})| = \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \alpha_{n,\mathbf{k}} X_{n,\mathbf{k}},$$

where

$$\alpha_{n,\mathbf{k}} = |Q_{n,\mathbf{k}}|^{-1} |Q_{n,\mathbf{k}}| - \mathbf{1}\{\mathbf{B}(\mathbf{k}) \in [0, \mathbf{t}]\}.$$

Now  $|\alpha_{n,\mathbf{k}}| < 1$ , and vanishes if  $Q_{n,\mathbf{k}} \subset [0, \mathbf{t}]$ , or  $Q_{n,\mathbf{k}} \cap [0, \mathbf{t}] = \emptyset$ . Actually  $\alpha_{n,\mathbf{k}} \neq 0$  if and only if  $\mathbf{k} \in I$ , where  $I$  is defined as

$$I := \{\mathbf{i} \leq \mathbf{n}; \forall k \in \{1, \dots, d\}, i_k \leq b_k(u_k(t_k) + 1) \text{ and} \\ \exists l \in \{1, \dots, d\}, i_l = b_l(u_l(t_l) + 1)\}.$$

Thus

$$\mathbf{E} |\Xi_n(\mathbf{t}) - \zeta_n(\mathbf{t})|^2 = \sum_{\mathbf{k} \in I} \alpha_{n,\mathbf{k}} \sigma_{n,\mathbf{k}}^2 \leq \sum_{\mathbf{k} \in I} \sigma_{n,\mathbf{k}}^2 \leq \sum_{l=1}^d \Delta b_l(u_l(t_l) + 1).$$

Now due to (3.37) we have

$$\mathbf{E} |\Xi_n(\mathbf{t}) - Z_n(\mathbf{t})|^2 \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty,$$

which coupled with the estimate  $P(|Y| > r) \leq r^{-2} \mathbf{E} |Y|^2$ , for any random

variable  $Y$  gives us

$$|\Xi_n(\mathbf{t}) - Z_n(\mathbf{t})| \xrightarrow{P} 0, \text{ as } m(\mathbf{n}) \rightarrow \infty. \quad \square$$

The condition (3.37) ensures that the grid gets thinner and thinner as  $m(\mathbf{n})$  approach the infinity. It is more restrictive than the condition of asymptotic negligibility. Define now

$$\mu_n(\mathbf{t}) = \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \mathbf{1}\{\mathbf{B}(\mathbf{k}) \in [0, \mathbf{t}]\} \sigma_{n,\mathbf{k}}^2. \quad (3.39)$$

It is clear that  $\mu_n(\mathbf{t}) = \mathbf{E} Z_n(\mathbf{t})^2$ . If the limiting finite-dimensional distributions of  $Z_n$  were those of Brownian sheet, then  $\mu_n(\mathbf{t})$  would converge to  $\boldsymbol{\pi}(\mathbf{t})$  for each  $\mathbf{t}$ . Consider the following example of triangular array.

**Example 1** For  $\mathbf{n} = (n, n)$  and  $\mathbf{k}_n = (2n, 2n)$  take  $X_{n,\mathbf{k}} = a_{n,\mathbf{k}} Y_{\mathbf{k}}$ , with  $\{Y_{\mathbf{k}}, \mathbf{k} \leq \mathbf{k}_n\}$  i.i.d. random variables with standard normal distribution, and

$$a_{n,\mathbf{k}}^2 = \begin{cases} \frac{1}{10n^2}, & \text{for } \mathbf{k} \leq (n, n) \\ \frac{3}{10n^2}, & \text{otherwise.} \end{cases} \quad (3.40)$$

Thus defined triangular array satisfies the condition (3.37), but simple algebra shows that for such an array

$$\begin{aligned} \mu_n(\mathbf{t}) \rightarrow \nu(\mathbf{t}) := & \frac{1}{10} \left( \frac{5}{2} t_1 \wedge 1 \right) \left( \frac{5}{2} t_2 \wedge 1 \right) + \frac{(5t_1 - 2) \vee 0}{10} \left( \frac{5}{2} t_2 \vee 1 \right) \\ & + \frac{(5t_2 - 2) \vee 0}{10} \left( \frac{5}{2} t_1 \vee 1 \right) + \frac{((5t_1 - 2) \vee 0)((5t_2 - 2) \vee 0)}{30}. \end{aligned}$$

Furthermore for the following example  $\mu_n(\mathbf{t})$  does not converge for any  $\mathbf{t}$ .

**Example 2** For  $\mathbf{n} = (n, n)$  and  $\mathbf{k}_n = (n, n)$  take  $X_{n,\mathbf{k}} = b_{n,\mathbf{k}} Y_{\mathbf{k}}$  with  $\{Y_{\mathbf{k}}, \mathbf{k} \leq \mathbf{k}_n\}$  i.i.d. random variables with standard normal distribution, and

$$b_{n,\mathbf{k}}^2 = \begin{cases} \pi(\mathbf{k}_n)^{-1}, & \text{for } \mathbf{n} = (2l - 1, 2l - 1), \ l \in \mathbb{N} \\ a_{n,\mathbf{k}}^2, & \text{for } \mathbf{n} = (2l, 2l), \ l \in \mathbb{N} \end{cases}$$

where  $a_{n,\mathbf{k}}$  are defined as in (3.40).

Nevertheless we can get some fruitful results by making following assumption.

**Assumption 1** *There exists a function  $\mu : [0, 1]^d \rightarrow \mathbb{R}$  such that*

$$\forall \mathbf{t} \in [0, 1]^d, \quad \lim_{m(\mathbf{n}) \rightarrow \infty} \mu_{\mathbf{n}}(\mathbf{t}) = \mu(\mathbf{t}). \quad (3.41)$$

With this assumption we can state the following result whose proof is postponed for a while.

**Theorem 25** *Given assumption 1 there exists a Gaussian process  $\{G(\mathbf{t}), \mathbf{t} \in [0, 1]^d\}$  with covariance function  $\mathbf{E} G(\mathbf{t})G(\mathbf{s}) = \mu(\mathbf{t} \wedge \mathbf{s})$ . Furthermore if*

$$\max_{1 \leq l \leq d} \max_{1 \leq k_l \leq k_n^l} \Delta b_l(k_l) \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty \quad (3.42)$$

and for every  $\varepsilon > 0$

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq k \leq k_n} \mathbf{E} X_{n,k}^2 \mathbf{1}\{|X_{n,k}| \geq \varepsilon\} = 0, \quad (3.43)$$

then for any collection of  $m$  points  $\mathbf{t}_1, \dots, \mathbf{t}_m \in [0, 1]^d$

$$(Z_n(\mathbf{t}_1), \dots, Z_n(\mathbf{t}_m)) \xrightarrow{D} (G(\mathbf{t}_1), \dots, G(\mathbf{t}_m)).$$

For the process  $\Xi_n$ , the following theorem holds.

**Theorem 26** *If there exists a function  $\mu : [0, 1]^d \rightarrow \mathbb{R}$  such that*

$$\forall \mathbf{t} \in [0, 1]^d, \quad \lim_{m(\mathbf{n}) \rightarrow \infty} \mathbf{E} \Xi_n^2(\mathbf{t}) = \mu(\mathbf{t}) \quad (3.44)$$

and

$$\max_{1 \leq l \leq d} \max_{1 \leq k_l \leq k_n^l} \Delta b_l(k_l) \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty \quad (3.45)$$

and for every  $\varepsilon > 0$

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq k \leq k_n} \mathbf{E} X_{n,k}^2 \mathbf{1}\{|X_{n,k}| \geq \varepsilon\} = 0, \quad (3.46)$$

then given any collection of  $m$  points  $\mathbf{t}_1, \dots, \mathbf{t}_m \in [0, 1]^d$

$$(\Xi_n(\mathbf{t}_1), \dots, \Xi_n(\mathbf{t}_m)) \xrightarrow{D} (G(\mathbf{t}_1), \dots, G(\mathbf{t}_m)),$$

where  $G$  is a Gaussian process satisfying  $\mathbf{E} G(\mathbf{t})G(\mathbf{s}) = \mu(\mathbf{t} \wedge \mathbf{s})$  for  $\mathbf{t}, \mathbf{s} \in [0, 1]^d$ .

*Proof.* Since (3.45) is satisfied lemma 24 gives us

$$|\mathbf{E} \Xi_n^2(\mathbf{t}) - \mathbf{E} Z_n^2(\mathbf{t})| \xrightarrow{P} 0, \text{ as } m(\mathbf{n}) \rightarrow \infty,$$

thus the limits  $\mu(\mathbf{t})$  in (3.41) and (3.44) coincide. The theorem 25 then gives us the existence of the process  $G$  and the same theorem combined again with lemma 24 gives us the proof.  $\square$

For triangular arrays with certain variance structure, the limiting process is always a Brownian sheet. Take double indexed triangular array  $\{X_{n,ij}, 1 \leq i \leq I_n, 1 \leq j \leq J_n\}$  with  $\mathbf{E} X_{n,ij}^2 = a_i b_j$ , where  $a_i, b_j > 0$  and  $\sum a_i = 1 = \sum b_j$ . Recalling notations (2.41) we get

$$b_1(k) = \sum_{i=1}^k a_i, \quad b_2(l) = \sum_{j=1}^l b_j$$

so our grid rectangle is now

$$Q_{n,kl} = \left[ \sum_{i=1}^{k-1} a_i, \sum_{i=1}^k a_i \right) \times \left[ \sum_{j=1}^{l-1} b_j, \sum_{j=1}^l b_j \right).$$

We see that grid points on x-axis are defined only by  $a_i$  and on y-axis by  $b_j$ . Now the variance of jump process in this case will be

$$\begin{aligned} \mu_n(\mathbf{t}) &= \sum_{i=1}^{I_n} \sum_{j=1}^{J_n} \mathbf{1} \left( \left( \sum_{k=1}^i a_k, \sum_{l=1}^j b_l \right) \in [0, t_1] \times [0, t_1] \right) a_i b_j \\ &= \sum_{i=1}^{I_n} \mathbf{1} \left( \sum_{k=1}^i a_k \in [0, t_1] \right) a_i \sum_{j=1}^{J_n} \mathbf{1} \left( \sum_{l=1}^j b_l \in [0, t_2] \right) b_j \end{aligned}$$

and if we assume the condition (3.45) which in this case translates to

$$\max_{1 \leq i \leq I_n} a_i \rightarrow 0, \quad \max_{1 \leq j \leq J_n} b_j \rightarrow 0,$$

we see that

$$\mu_n(\mathbf{t}) \rightarrow t_1 t_2,$$

which is the variance of Brownian sheet. Thus assuming the Lindeberg condition (3.46), theorem 25 implies that limiting finite-dimensional distributions in this case are those of Brownian sheet.

**Proof of theorem 25**

Define

$$g(\mathbf{t}, \mathbf{s}) = \lim_{m(\mathbf{n}) \rightarrow \infty} \mu_{\mathbf{n}}(\mathbf{t} \wedge \mathbf{s}).$$

If we prove that  $g(\mathbf{t}, \mathbf{s})$  is positive definite, then the existence of zero mean Gaussian process  $\{G(\mathbf{t}), \mathbf{t} \in [0, 1]^d\}$  with covariance function  $\mathbf{E} G(\mathbf{t})G(\mathbf{s}) = g(\mathbf{t}, \mathbf{s})$  is ensured by theorem 7. Take  $p \in \mathbb{N}$ ,  $v_1, \dots, v_p \in \mathbb{R}$  and  $\mathbf{t}_1, \dots, \mathbf{t}_p \in [0, 1]^d$ . Note that for any  $\mathbf{t}, \mathbf{s}, \mathbf{r} \in [0, 1]^d$  we have

$$\mathbf{1}\{\mathbf{r} \in [0, \mathbf{t} \wedge \mathbf{s}]\} = \mathbf{1}\{\mathbf{r} \in [0, \mathbf{t}] \cap [0, \mathbf{s}]\} = \mathbf{1}\{\mathbf{r} \in [0, \mathbf{t}]\} \mathbf{1}\{\mathbf{r} \in [0, \mathbf{s}]\}. \quad (3.47)$$

Then

$$\begin{aligned} \sum_{i=1}^p \sum_{j=1}^p v_i \mu_{\mathbf{n}}(\mathbf{t}_i \wedge \mathbf{t}_j) v_j &= \sum_{i=1}^p \sum_{j=1}^p v_i v_j \sum_{\mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} \mathbf{1}\{B_{\mathbf{n}}(\mathbf{k}) \in [0, \mathbf{t}_i \wedge \mathbf{t}_j]\} \sigma_{\mathbf{n}, \mathbf{k}}^2 \\ &= \sum_{\mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} \sigma_{\mathbf{n}, \mathbf{k}}^2 \left( \sum_{i=1}^p v_i \mathbf{1}\{B_{\mathbf{n}}(\mathbf{k}) \in [0, \mathbf{t}_i]\} \right)^2 \geq 0. \end{aligned}$$

Since this holds for each  $\mathbf{n}$ , taking the limit as  $m(\mathbf{n}) \rightarrow \infty$  gives the positive definiteness of  $g(\mathbf{t}, \mathbf{s})$ . So the first part of the theorem is proved.

Now fix  $\mathbf{t}_1, \dots, \mathbf{t}_r \in [0, 1]^d$  and  $v_1, \dots, v_r$  real, and set

$$V_{\mathbf{n}} = \sum_{p=1}^r v_p Z_{\mathbf{n}}(\mathbf{t}_p) = \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} \alpha_{\mathbf{n}, \mathbf{k}} X_{\mathbf{n}, \mathbf{k}},$$

where

$$\alpha_{\mathbf{n}, \mathbf{k}} = \sum_{p=1}^r v_p \mathbf{1}(B(\mathbf{k}) \in [0, \mathbf{t}_p]).$$

Now using (3.47) we get

$$\begin{aligned} b_{\mathbf{n}} &:= \mathbf{E} V_{\mathbf{n}}^2 = \sum_{\mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} \alpha_{\mathbf{n}, \mathbf{k}}^2 \sigma_{\mathbf{n}, \mathbf{k}}^2 \\ &= \sum_{\mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} \sum_p \sum_q v_p v_q \mathbf{1}(B(\mathbf{k}) \in [0, \mathbf{t}_p]) \mathbf{1}(B(\mathbf{k}) \in [0, \mathbf{t}_q]) \sigma_{\mathbf{n}, \mathbf{k}}^2 \\ &= \sum_p \sum_q v_p v_q \mu_{\mathbf{n}}(\mathbf{t}_p \wedge \mathbf{t}_q). \end{aligned}$$

Letting  $m(\mathbf{n})$  tend to infinity and using assumption 1, we obtain

$$b_n \xrightarrow{m(\mathbf{n}) \rightarrow \infty} \sum_p \sum_q v_p v_q \mu(\mathbf{t}_p \wedge \mathbf{t}_q) = \mathbf{E} \left( \sum_p v_p G(\mathbf{t}_p) \right)^2 =: b.$$

If  $b = 0$ , then  $V_n$  converges to zero in distribution since  $\mathbf{E} V_n^2$  tends to zero. In this special case we also have  $\sum_p v_p G(\mathbf{t}_p) = 0$  almost surely, thus the convergence of finite dimensional distributions holds.

Assume now, that  $b > 0$ . For convenience put  $Y_{n,k} = \alpha_{n,k} X_{n,k}$  and  $v = \sum_p \sum_q v_p v_q$ . Conditions (3.42) and (3.43) ensures that triangular array  $X_{n,j}$  satisfies the conditions for central limit theorem: infinitesimal negligibility and Lindeberg condition. The same is true for triangular array  $\{Y_{n,k}\}$ . We have

$$Y_{n,k}^2 \leq v X_{n,k}^2,$$

thus  $Y_{n,k}$  satisfies the condition of infinitesimal negligibility. For  $m(\mathbf{n})$  large enough to have  $b_n > b/2$ , we get

$$\begin{aligned} & \frac{1}{\mathbf{E} V_n^2} \sum_{1 \leq k \leq k_n} \mathbf{E} \left( Y_{n,k}^2 \mathbf{1}_{\{|Y_{n,k}|^2 > \varepsilon^2 \mathbf{E} V_n^2\}} \right) \\ & \leq \frac{2v}{b} \sum_{1 \leq k \leq k_n} \mathbf{E} \left( X_{n,k}^2 \mathbf{1}_{\{|X_{n,k}|^2 > \frac{b\varepsilon^2}{2v}\}} \right). \end{aligned}$$

Thus Lindeberg condition for  $V_n$  is also satisfied and that gives us the convergence of finite dimensional distributions and the proof of the theorem.

### 3.2.2 Tightness

To prove tightness of process  $\Xi_n$  only certain moment conditions are required. There is no need for additional variance structure assumptions as proving the convergence of finite-dimensional distributions. This is quite clear, since due to results from section 2.2.2 and corollary 6, the process  $\Xi_n$  is tight if

$$\lim_{a \rightarrow \infty} P \left( \sup_{t \in [0,1]^d} |\Xi_n(\mathbf{t})| > a \right) = 0 \quad (3.48)$$

and for every  $\varepsilon > 0$  and  $m = 1, \dots, d$ ,

$$\lim_{m(\mathbf{n}) \rightarrow \infty} P\left(Z_{\mathbf{n}}^{(m)} > \varepsilon\right) = 0; \quad (3.49)$$

$$\lim_{J \rightarrow \infty} \lim_{m(\mathbf{n}) \rightarrow \infty} P\left(\sup_{j \geq J} 2^{-j\alpha} \max_{r \in D_j} \psi_{\mathbf{n}}(r, r^-)^{(m)} > \varepsilon\right) = 0; \quad (3.50)$$

$$\lim_{J \rightarrow \infty} \lim_{m(\mathbf{n}) \rightarrow \infty} P\left(\sup_{j \geq J} 2^{-j\alpha} \max_{r \in D_j} \psi_{\mathbf{n}}(r, r^+)^{(m)} > \varepsilon\right) = 0, \quad (3.51)$$

recalling respectively the definitions (2.51), (2.50) and (2.52).

Using Doob inequality (1.12) we have

$$\begin{aligned} P\left(\sup_{t \in [0,1]^d} |\Xi_{\mathbf{n}}(t)| > a\right) &= P\left(\max_{\mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} |S_{\mathbf{n}}(\mathbf{k})| > a\right) \\ &\leq a^{-2} \mathbf{E} S_{\mathbf{n}}(\mathbf{k}_{\mathbf{n}})^2 = a^{-2} \rightarrow 0, \text{ as } a \rightarrow \infty, \end{aligned}$$

thus (3.48) is satisfied leaving us with checking (3.49) to (3.51). As the expressions in the probability involve only sums we can use similar techniques as in proving tightness of process  $\xi_{\mathbf{n}}$ . We give now two sets of conditions.

**Theorem 27** For  $0 < \alpha < 1/2$ , set  $p(\alpha) := 1/(1/2 - \alpha)$ . If

$$\max_{1 \leq l \leq d} \max_{1 \leq k_l \leq k_{\mathbf{n}}^l} \Delta b_l(k_l) \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty. \quad (3.52)$$

and for some  $q > p(\alpha)$

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} \sigma_{\mathbf{n}, \mathbf{k}}^{-2q\alpha} \mathbf{E} |X_{\mathbf{n}, \mathbf{k}}|^q = 0, \quad (3.53)$$

then the net  $\{\Xi_{\mathbf{n}}, \mathbf{n} \in \mathbb{N}\}$  is asymptotically tight in the space  $H_{\alpha}^o([0, 1]^d)$ .

Introduce for every  $\tau > 0$ , the truncated random variables:

$$X_{\mathbf{n}, \mathbf{k}, \tau} := X_{\mathbf{n}, \mathbf{k}} \mathbf{1}\{|X_{\mathbf{n}, \mathbf{k}}| \leq \tau \sigma_{\mathbf{n}, \mathbf{k}}^{2\alpha}\}.$$

**Theorem 28** Assume that

$$\max_{1 \leq l \leq d} \max_{1 \leq k_l \leq k_{\mathbf{n}}^l} \Delta b_l(k_l) \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty. \quad (3.54)$$

and that the following conditions hold.

(a). For every  $\varepsilon > 0$ ,

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} P(|X_{n,\mathbf{k}}| \geq \varepsilon \sigma_{n,\mathbf{k}}^{2\alpha}) = 0. \quad (3.55)$$

(b). For every  $\varepsilon > 0$ ,

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} \mathbf{E} X_{n,\mathbf{k}}^2 \mathbf{1}\{|X_{n,\mathbf{k}}| \geq \varepsilon\} = 0. \quad (3.56)$$

(c). For some  $q > 1/(1/2 - \alpha)$ ,

$$\lim_{\tau \rightarrow 0} \lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} \sigma_{n,\mathbf{k}}^{-2q\alpha} \mathbf{E} |X_{n,\mathbf{k},\tau}|^q = 0. \quad (3.57)$$

Then the net  $\{\Xi_n, \mathbf{n} \in \mathbb{N}\}$  is asymptotically tight in the space  $H_\alpha^0([0, 1]^d)$ .

### Proof of the theorem 27

We only need to check that (3.52) and (3.53) ensure (3.49), (3.50), (3.51). We check only the case  $m = 1$ , since the proof is the same for other  $m$ , thus in following proofs we drop the superscript  $m$ .

*Proof of (3.49).* Using Markov and Doob (1.12) inequalities for  $q > 1/(1/2 - \alpha)$  we get

$$\begin{aligned} P\left(Z_n > \varepsilon\right) &\leq \sum_{k=1}^{k_n^1} P\left(\max_{\mathbf{k}_{2:d} \leq \mathbf{k}_{n,2:d}} |\Delta_k^{(1)} S_n(\mathbf{k})| > \varepsilon (\Delta b_1(k))^\alpha\right) \\ &\leq \sum_{k=1}^{k_n^1} \varepsilon^{-q} (\Delta b_1(k))^{-q\alpha} \mathbf{E} \left(\max_{\mathbf{k}_{2:d} \leq \mathbf{k}_{n,2:d}} |\Delta_k^{(1)} S_n(\mathbf{k})|\right)^q \\ &\leq \sum_{k=1}^{k_n^1} \varepsilon^{-q} (\Delta b_1(k))^{-q\alpha} \mathbf{E} |\Delta_k^{(1)} S_n(\mathbf{k}_n)|^q. \end{aligned}$$

Rosenthal inequality (1.10) gives

$$P\left(Z_n > \varepsilon\right) \leq c \sum_{k=1}^{k_n^1} \varepsilon^{-q} (\Delta b_1(k))^{-q\alpha} \left( (\Delta b_1(k))^{q/2} + \sum_{k_2=1}^{k_n^2} \cdots \sum_{k_d=1}^{k_n^d} \mathbf{E} |X_{n,\mathbf{k}}|^q \right). \quad (3.58)$$

We have

$$\begin{aligned} \sum_{k=1}^{k_n^1} (\Delta b_1(k))^{q(1/2-\alpha)} &\leq \left( \max_{1 \leq k \leq k_n^1} \Delta b_1(k) \right)^{q(1/2-\alpha)-1} \sum_{k=1}^{k_n^1} \Delta b_1(k) \\ &= \left( \max_{1 \leq k \leq k_n^1} \Delta b_1(k) \right)^{q(1/2-\alpha)-1} \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty, \end{aligned} \quad (3.59)$$

due to (3.52) and the fact that  $q > (1/2 - \alpha)^{-1}$ . Also

$$\begin{aligned} \sum_{k=1}^{k_n^1} (\Delta b_1(k))^{-q\alpha} \sum_{k_2=1}^{k_n^2} \cdots \sum_{k_d=1}^{k_n^d} \mathbf{E} |X_{n,k}|^q &= \sum_{\mathbf{k} \leq \mathbf{k}_n} (\Delta b_1(k_1))^{-q\alpha} \mathbf{E} |X_{n,k}|^q \\ &\leq \sum_{\mathbf{k} \leq \mathbf{k}_n} \sigma_{n,k}^{-2q\alpha} \mathbf{E} |X_{n,k}|^q \rightarrow 0, \end{aligned}$$

as  $m(\mathbf{n}) \rightarrow \infty$ , due to (3.53), since  $(\Delta b_1(k_1))^{-q\alpha} \leq \sigma_{n,k}^{-2q\alpha}$  for all  $1 \leq \mathbf{k} \leq \mathbf{k}_n$ . Reporting these estimates to (3.58) we see that (3.52) and (3.53) imply (3.49).

*Proof of (3.50) and (3.51).* We check only (3.50) since (3.51) is treated similarly. Define

$$\Pi(J, \mathbf{n}, \varepsilon) = P\left(\sup_{j \geq J} 2^{-j\alpha} \max_{r \in D_j} \psi_{\mathbf{n}}(r, r^-)^{(m)} > \varepsilon\right)$$

then

$$\Pi(J, \mathbf{n}, \varepsilon) \leq \sum_{j \geq J} P(2^{\alpha j} \max_{r \in D_j} \psi_{\mathbf{n}}(r, r^-) > \varepsilon) \leq \sum_{j \geq J} \sum_{r \in D_j} \varepsilon^{-q} 2^{\alpha j q} \mathbf{E} |\psi_{\mathbf{n}}(r, r^-)|^q.$$

Doob (1.12) and then Rosenthal (1.9) inequalities give us

$$\begin{aligned} \mathbf{E} \psi_{\mathbf{n}}(r, r^-)^q &\leq \mathbf{E} \left| \sum_{\mathbf{k}_{2:d} \leq \mathbf{k}_{n,2:d}} \left( \sum_{k_1=u_1(r^-)+2}^{u_1(r)} X_{n,k} \right) \right|^q \\ &\leq c \left( \left( \sum_{k_1=u_1(r^-)+2}^{u_1(r)} \sum_{\mathbf{k}_{2:d} \leq \mathbf{k}_{n,2:d}} \sigma_{n,k}^2 \right)^{q/2} \right. \\ &\quad \left. + \sum_{k_1=u_1(r^-)+2}^{u_1(r)} \sum_{\mathbf{k}_{2:d} \leq \mathbf{k}_{n,2:d}} \mathbf{E} |X_{n,k}|^q \right). \end{aligned}$$

Due to definition of  $u_1(r)$

$$\sum_{k_1=u_1(r^-)+2}^{u_1(r)} \sum_{\mathbf{1} \leq \mathbf{k}_{2:d} \leq \mathbf{k}_{\mathbf{n},2:d}} \sigma_{\mathbf{n},\mathbf{k}}^2 = \sum_{k_1=u_1(r^-)+2}^{u_1(r)} \Delta b_1(k_1) \leq r - r^- = 2^{-j},$$

thus

$$\begin{aligned} \Pi(J, \mathbf{n}, \varepsilon) &\leq \frac{C}{\varepsilon^q} \sum_{j \geq J} 2^{(q\alpha+1-q/2)j} \\ &\quad + \frac{C}{\varepsilon^q} \sum_{j \geq J} \sum_{r \in D_j} 2^{q\alpha j} \sum_{k_1=u_1(r^-)+2}^{u_1(r)} \sum_{\mathbf{k}_{2:d} \leq (\mathbf{k}_{\mathbf{n}})_{2:d}} \mathbf{E} |X_{\mathbf{n},\mathbf{k}}|^q. \end{aligned} \quad (3.60)$$

Denote by  $I(J, \mathbf{n}, q)$  the second sum without the constant  $C\varepsilon^{-q}$ . By changing the order of summation we get

$$I(J, \mathbf{n}, q) = \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_{\mathbf{n}}} \mathbf{E} |X_{\mathbf{n},\mathbf{k}}|^q \sum_{j \geq J} 2^{q\alpha j} \sum_{r \in D_j} \mathbf{1}\{u_1(r^-) + 1 < k_1 \leq u_1(r)\}. \quad (3.61)$$

The proof further proceeds as in [29]. Consider for fixed  $k_1$  the condition

$$u_1(r^-) + 1 < k_1 < u_1(r). \quad (3.62)$$

Suppose that there exists  $r \in D_j$  satisfying (3.62) and take another  $r' \in D_j$ . Since  $u_1$  is non decreasing, if  $r' < r^-$  we have  $u_1(r') < u_1(r^-) + 1 < k$ , and thus  $r'$  cannot satisfy (3.62). If  $r' > r$ , then  $r'^- > r$ , whence  $k \leq u_1(r) \leq u_1(r'^-) < u_1(r'^-) + 1$  and again it follows that  $r'$  cannot satisfy (3.62). Thus there will exist at most only one  $r$  satisfying (3.62). If such  $r$  exists we have

$$r^- \leq \sum_{i=1}^{u_1(r^-)+1} \Delta b_1(i) < \sum_{i=1}^{k_1} \Delta b_1(i) \leq \sum_{i=1}^{u_1(r)} \Delta b_1(i) \leq r.$$

Thus  $\Delta b_1(k_1) \leq 2^{-j}$ . So

$$\forall k_1 = 1, \dots, k_{\mathbf{n}}^1, \sum_{r \in D_j} \mathbf{1}\{u_1(r^-) + 1 < k_1 \leq u_1(r)\} \leq \mathbf{1}\{\Delta b_1(k_1) \leq 2^{-j}\}$$

so

$$\sum_{j \geq J} 2^{q\alpha j} \sum_{r \in D_j} \mathbf{1}\{u_1(r^-) + 1 < k_1 \leq u_1(r)\} \leq \frac{2^{q\alpha}}{2^{q\alpha} - 1} (\Delta b_1(k_1))^{-\alpha q} \quad (3.63)$$

(we can sum only those  $j$ , for which  $\Delta b_1(k_1) \leq 2^{-j}$ , because for larger  $j$ ,  $r$  and  $r^-$  will be closer together and will fall in the same  $R_{n,k}$ ).

Reporting estimate (3.63) to (3.61) we get

$$I(J, \mathbf{n}, q) \leq C \sum_{k \leq k_n} (\Delta b_1(k_1))^{-q\alpha} \mathbf{E} |X_{n,k}|^q \leq \sum_{k \leq k_n} \sigma_{n,k}^{-2q\alpha} \mathbf{E} |X_{n,k}|^q$$

and substituting this into inequality (3.60) we get

$$\Pi(J, \mathbf{n}; \varepsilon) \leq C_1 \varepsilon^{-q} 2^{-Jq\alpha+1-q/2} + C_2 \sum_{k \leq k_n} \sigma_{n,k}^{-2q\alpha} \mathbf{E} |X_{n,k}|^q.$$

Thus

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \Pi(J, \mathbf{n}; \varepsilon) = 0$$

follows from (3.53), which gives us the proof of the theorem.

### Proof of theorem 28

As in proof of the theorem 27 we check (3.49), (3.50), (3.51) and give a proof only for case  $m = 1$ .

*Proof of (3.49)* Define:

$$S_{n,\tau}(\mathbf{k}) = \sum_{1 \leq j \leq k} X_{n,j,\tau}, \quad S_{n,\tau}(\mathbf{k})' = \sum_{1 \leq j \leq k} (X_{n,j,\tau} - \mathbf{E} X_{n,j,\tau})$$

and

$$A_n = \left\{ \max_{1 \leq k \leq k_n} |X_k| \leq \tau \sigma_{n,k}^{2\alpha} \right\}.$$

Then we can estimate the probability in (3.49) by

$$P(Z_n > \varepsilon) =: P(\mathbf{n}, \varepsilon) \leq P_1(\mathbf{n}, \varepsilon, \tau) + P(A_n^c)$$

where

$$P_1(\mathbf{n}, \varepsilon, \tau) = P\left( \max_{1 \leq k \leq k_n} \frac{|\Delta_{k_1}^{(1)} S_{n,\tau}(\mathbf{k})|}{(\Delta b_1(k_1))^\alpha} > \varepsilon \right). \quad (3.64)$$

Due to (3.55) the probability  $P(A_n^c)$  tends to zero so we need only to study the asymptotics of  $P_1(\mathbf{n}, \varepsilon, \tau)$ .

Using the splitting

$$\Delta_{k_1}^{(1)} S_{n,\tau}(\mathbf{k}) = \Delta_{k_1}^{(1)} S'_{n,\tau}(\mathbf{k}) + \mathbf{E} \Delta_{k_1}^{(1)} S_{n,\tau}(\mathbf{k}),$$

let us begin with some estimate of the expectation term, since  $X_{n,\mathbf{k},\tau}$  are not centered.

We have

$$\mathbf{E} |X_{n,\mathbf{k},\tau}| \leq \mathbf{E}^{1/2} X_{n,\mathbf{k}}^2 P^{1/2}(|X_{n,\mathbf{k}}| > \tau \sigma_{n,\mathbf{k}}^{2\alpha}).$$

By applying Cauchy inequality we get

$$\begin{aligned} \max_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \frac{|\mathbf{E} \Delta_{k_1}^{(1)} S_{n,\tau}(\mathbf{k})|}{(\Delta b_1(k_1))^\alpha} &\leq \max_{1 \leq k_1 \leq k_n^1} \frac{\sum_{k_{2:d}=1}^{k_{n,2:d}} \mathbf{E} |X_{n,j,\tau}|}{(\Delta b_1(k_1))^\alpha} \\ &\leq \max_{1 \leq k_1 \leq k_n^1} \frac{(\Delta b_1(k_1))^{1/2} \left( \sum_{k_{2:d}=1}^{k_{n,2:d}} P(|X_{n,\mathbf{k}}| > \tau \sigma_{n,\mathbf{k}}^{2\alpha}) \right)^{1/2}}{(\Delta b_1(k_1))^\alpha} \\ &\leq \max_{1 \leq k_1 \leq k_n^1} (\Delta b_1(k_1))^{1/2-\alpha} \left( \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} P(|X_{n,\mathbf{k}}| > \tau \sigma_{n,\mathbf{k}}^{2\alpha}) \right)^{1/2}. \end{aligned}$$

Due to (3.54) and (3.55) the last expression is bounded by  $\varepsilon/2$  for  $\mathbf{n} \geq \mathbf{n}_0$ , where  $\mathbf{n}_0$  depends on  $\varepsilon$  and  $\tau$ . Thus for  $\mathbf{n} \geq \mathbf{n}_0$  we have  $P_1(\mathbf{n}, \varepsilon, \tau) \leq P'_1(\mathbf{n}, \varepsilon, \tau)$ , where

$$P'_1(\mathbf{n}, \varepsilon, \tau) = P \left( \max_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \frac{|\Delta_{k_1}^{(1)} S'_{n,\tau}(\mathbf{k})|}{(\Delta b_1(k_1))^\alpha} > \varepsilon/2 \right). \quad (3.65)$$

Since

$$\text{Var} X_{n,\mathbf{k},\tau} \leq \mathbf{E} X_{n,\mathbf{k},\tau}^2 \leq \mathbf{E} X_{n,\mathbf{k}}^2 = \sigma_{n,\mathbf{k}}^2,$$

using Markov, Doob and Rosenthal inequalities for  $q > 1/(1/2 - \alpha)$  we get

$$\begin{aligned}
P'_1(\mathbf{n}, \varepsilon, \tau) &\leq \sum_{k=1}^{k_n^1} (\varepsilon/2)^{-q} (\Delta b_1(k))^{-q\alpha} \mathbf{E} |\Delta_k^{(1)} S'_{\mathbf{n}, \tau}(\mathbf{k}_n)|^q \\
&\leq c \sum_{k=1}^{k_n^1} (\varepsilon/2)^{-q} (\Delta b_1(k))^{-q\alpha} \left( (\Delta b_1(k))^{q/2} + \sum_{\mathbf{k}_{2:d}=1}^{\mathbf{k}_{n,2:d}} \mathbf{E} |X_{\mathbf{n}, \mathbf{k}, \tau}|^q \right) \\
&\leq c (\varepsilon/2)^{-q} \left( \sum_{k=1}^{k_n^1} (\Delta b_1(k))^{q(1/2-\alpha)} + \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \sigma_{\mathbf{n}, \mathbf{k}}^{-2q\alpha} \mathbf{E} |X_{\mathbf{n}, \mathbf{k}, \tau}|^q \right).
\end{aligned}$$

Note that this estimate holds for each  $\tau > 0$ . Combining all the estimates we get

$$\forall \tau > 0, \quad \limsup_{\mathbf{m}(\mathbf{n}) \rightarrow \infty} P(\mathbf{n}, \varepsilon) \leq c \limsup_{\mathbf{m}(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \sigma_{\mathbf{n}, \mathbf{k}}^{-2q\alpha} \mathbf{E} |X_{\mathbf{n}, \mathbf{k}, \tau}|^q.$$

with the constant  $c$  depending only on  $q$ . By letting  $\tau \rightarrow 0$  due to (3.57), (3.49) follows.

*Proof of (3.50) and (3.51)* We again prove only (3.50) since (3.51) is treated similarly. Introduce definitions  $\psi_{\mathbf{n}, \tau}(r, r^-)$  and  $\psi'_{\mathbf{n}, \tau}(r, r^-)$  by exchanging variables  $X_{\mathbf{n}, \mathbf{k}}$  with variables  $X_{\mathbf{n}, \mathbf{k}, \tau}$  and  $X'_{\mathbf{n}, \mathbf{k}, \tau} := X_{\mathbf{n}, \mathbf{k}, \tau} - \mathbf{E} X_{\mathbf{n}, \mathbf{k}, \tau}$  respectively. Define

$$P(J, \mathbf{n}, \varepsilon) = P\left(\sup_{j \geq J} 2^{\alpha j} \max_{r \in D_j} \psi_{\mathbf{n}}(r, r^-) > \varepsilon\right).$$

Similar to the proof of (3.49) we need only to deal with asymptotics of  $P_1(J, \mathbf{n}, \varepsilon, \tau)$ , where

$$P_1(J, \mathbf{n}, \varepsilon, \tau) = P\left(\sup_{j \geq J} 2^{\alpha j} \max_{r \in D_j} \psi_{\mathbf{n}, \tau}(r, r^-) > \varepsilon\right).$$

Again we need to estimate the expectation term. We have

$$\begin{aligned}
& \sup_{j \geq J} 2^{\alpha j} \max_{r \in D_j} \max_{\mathbf{1}_{2:d} \leq \mathbf{k}_{2:d} \leq \mathbf{k}_{n,2:d}} \left| \sum_{i=u_1(r^-)+2}^{u_1(r)} \Delta_i^{(1)} \mathbf{E} S_{n,\tau}((i, \mathbf{k}_{2:d})) \right| \\
& \leq \sup_{j \geq J} 2^{\alpha j} \max_{r \in D_j} \left( \sum_{i=u_1(r^-)+2}^{u_1(r)} \Delta b_1(i) \right)^{1/2} \left( \sum_{i=u_1(r^-)+2}^{u_1(r)} \sum_{\mathbf{k}_{2:d}=1}^{\mathbf{k}_{n,2:d}} P(|X_{n,\mathbf{k}}| > \sigma_{n,\mathbf{k}}^{2\alpha}) \right)^{1/2} \\
& \leq 2^{J(\alpha-1/2)} \left( \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} P(|X_{n,\mathbf{k}}| > \sigma_{n,\mathbf{k}}^{2\alpha}) \right)^{1/2}.
\end{aligned}$$

The last expression is bounded by  $\varepsilon/2$  for  $\mathbf{n} \geq \mathbf{n}_0$ , due to (3.55) where  $\mathbf{n}_0$  depends on  $\varepsilon$  and  $\tau$ , but not on  $J$ . Thus  $P_1(J, \mathbf{n}, \varepsilon, \tau) \leq P'_1(J, \mathbf{n}, \varepsilon, \tau)$ , where

$$P'_1(J, \mathbf{n}, \varepsilon, \tau) := P\left(\sup_{j \geq J} 2^{\alpha j} \max_{r \in D_j} \psi'_{n,\tau}(r, r^-) > \varepsilon/2\right). \quad (3.66)$$

Applying the same arguments as in proving (3.60) we get

$$\begin{aligned}
P'_1(J, \mathbf{n}, \varepsilon, \tau) & \leq \frac{c}{\varepsilon^q} \sum_{j \geq J} 2^{(q\alpha+1-q/2)j} \\
& \quad + \frac{c}{\varepsilon^q} \sum_{j \geq J} \sum_{r \in D_j} 2^{q\alpha j} \sum_{k_1=u_1(r^-)+2}^{u_1(r)} \sum_{\mathbf{1} \leq \mathbf{k}_{2:d} \leq \mathbf{k}_{n,2:d}} \mathbf{E} |X_{n,\mathbf{k},\tau}|^q.
\end{aligned}$$

Now using estimate (3.63) we get

$$P'_1(J, \mathbf{n}, \varepsilon, \tau) \leq C_1 2^{(q\alpha+1-q/2)J} + C_2 \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} \sigma_{n,\mathbf{k}}^{-2q\alpha} \mathbf{E} |X_{n,\mathbf{k},\tau}|^q,$$

where constants  $C_1$  and  $C_2$  depend on  $q$ ,  $\alpha$  and  $\varepsilon$ . Note that this inequality holds for each  $\tau > 0$ . Combining all the estimates we get

$$\forall \tau > 0, \quad \lim_{J \rightarrow \infty} \limsup_{m(\mathbf{n}) \rightarrow \infty} P(J, \mathbf{n}, \varepsilon) \leq C_2 \limsup_{m(\mathbf{n}) \rightarrow \infty} \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} \sigma_{n,\mathbf{k}}^{-2q\alpha} \mathbf{E} |X_{n,\mathbf{k},\tau}|^q.$$

By letting  $\tau \rightarrow 0$  due to (3.57), (3.50) follows.

### 3.2.3 Corollaries

Recalling the results from previous subsections we have the following functional central limit theorem for the summation process  $\Xi_{\mathbf{n}}$ .

**Theorem 29** Suppose there exists a function  $\mu : [0, 1]^d \rightarrow \mathbb{R}$  such that

$$\forall \mathbf{t} \in [0, 1]^d, \quad \lim_{m(\mathbf{n}) \rightarrow \infty} \mathbf{E} \Xi_{\mathbf{n}}^2(\mathbf{t}) = \mu(\mathbf{t}). \quad (3.67)$$

For  $0 < \alpha < 1/2$ , set  $p(\alpha) := 1/(1/2 - \alpha)$ . If

$$\max_{1 \leq l \leq d} \max_{1 \leq k_l \leq k_n^l} \Delta b_l(k_l) \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty \quad (3.68)$$

and for some  $q > p(\alpha)$

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} \sigma_{\mathbf{n}, \mathbf{k}}^{-2q\alpha} \mathbf{E} |X_{\mathbf{n}, \mathbf{k}}|^q = 0, \quad (3.69)$$

then

$$\Xi_{\mathbf{n}} \xrightarrow[m(\mathbf{n}) \rightarrow \infty]{H_{\alpha}^o([0, 1]^d)} G, \quad (3.70)$$

where  $G$  is a centered Gaussian process satisfying  $\mathbf{E} G(\mathbf{t})G(\mathbf{s}) = \mu(\mathbf{t} \wedge \mathbf{s})$  for  $\mathbf{t}, \mathbf{s} \in [0, 1]^d$ .

*Proof.* We have

$$\sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} \mathbf{E} X_{\mathbf{n}, \mathbf{k}}^2 \mathbf{1}\{|X_{\mathbf{n}, \mathbf{k}}| \geq \varepsilon\} \leq \frac{1}{\varepsilon^{q-2}} \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} \mathbf{E} |X_{\mathbf{n}, \mathbf{k}}|^q.$$

Since  $\sigma_{\mathbf{n}, \mathbf{k}}^2 \leq 1$ , condition (3.69) ensures that  $\sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} \mathbf{E} |X_{\mathbf{n}, \mathbf{k}}|^q$  converges to zero, thus conditions of theorem 26 are satisfied and we have convergence of finite-dimensional distributions. Furthermore the conditions for theorem 27 are satisfied also, so the process  $\Xi_{\mathbf{n}}$  is asymptotically tight in the space  $H_{\alpha}^o([0, 1]^d)$ . The theorem then follows.  $\square$

Our motivation for introducing special construction for the collections of random variables with non-uniform variance was to get one limiting process in functional central limit theorem for all possible variance structures of the collection. It is clear from theorem 29 that this goal was not achieved. Nevertheless we think that this is an improvement compared to using non-modified construction. The convergence of process  $\xi_{\mathbf{n}}$  in case of non-uniform variance was investigated by Goldie and Greenwood [14], [15]. Their focus was on non-independent variables and although the domain of summation process was wider, for  $[0, 1]^d$  their process coincides with  $\xi_{\mathbf{n}}$ . They proved the convergence to Brownian sheet in case  $\mathbf{n} = (n, \dots, n)$  in the space of continuous functions, but naturally their result requires that  $\mathbf{E} \xi_{\mathbf{n}}(\mathbf{t}) \rightarrow \pi(\mathbf{t})$  for all  $\mathbf{t} \in [0, 1]^d$ , which is the special case of our requirement (3.67). Fur-

thermore to achieve the convergence they place quite strict conditions on variances of random variables by requiring that

$$\lim_{c \rightarrow \infty} \sup_{1 \leq \mathbf{k} \leq \mathbf{k}_n} \mathbf{E} n^d X_{\mathbf{n}, \mathbf{k}}^2 \mathbf{1}(|n^{d/2} X_{\mathbf{n}, \mathbf{k}}| > c) = 0.$$

This limit follows from Lindeberg condition if we take  $\mathbf{E} X_{\mathbf{n}, \mathbf{k}}^2 = n^{-d}$ . Thus the variances in Goldie-Greenwood case have the additional restriction, which is unnecessary when using our proposed construction. Furthermore for the special structure of variances the convergence  $\mathbf{E} \Xi_n(\mathbf{t}) \rightarrow \boldsymbol{\pi}(\mathbf{t})$  is always satisfied.

**Corollary 30** *Let  $\sigma_{\mathbf{n}, \mathbf{k}}^2 = \pi(\mathbf{a}_{\mathbf{n}, \mathbf{k}})$ , where  $\{\mathbf{a}_{\mathbf{n}, \mathbf{k}} = (a_{\mathbf{n}, \mathbf{k}_1}^1, \dots, a_{\mathbf{n}, \mathbf{k}_d}^d)\}$  is a triangular array of real vectors satisfying the following conditions for each  $i = 1, \dots, d$  and for all  $\mathbf{k} \leq \mathbf{k}_n$ .*

$$i) \sum_{k=1}^{k_n^i} a_{\mathbf{n}, k}^i = 1 \text{ with } a_{\mathbf{n}, k_i}^i > 0.$$

ii)

$$\max_{1 \leq k \leq k_n^i} a_{\mathbf{n}, k}^i \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty.$$

*Then condition (3.69) is sufficient for weak convergence of summation process  $\Xi_n$  in the space  $H_\alpha^o([0, 1]^d)$  and the limiting process is then Brownian sheet  $W$ .*

*Proof.* The result follows from theorem 29 if we check that

$$\mathbf{E} \Xi_n^2(\mathbf{t}) \rightarrow \boldsymbol{\pi}(\mathbf{t}), \quad (3.71)$$

since evidently the condition (3.37) is satisfied. We have

$$b_i(k_i) = \sum_{k=1}^{k_i} a_{\mathbf{n}, k}^i,$$

thus

$$\mathbf{1}\{\mathbf{B}(\mathbf{k}) \in [0, \mathbf{t}]\} = \prod_{i=1}^d \mathbf{1}\{b_i(k_i) \in [0, t_i]\},$$

so for jump process  $Z_n$  we have

$$\mathbf{E} Z_n^2(\mathbf{t}) = \sum_{\mathbf{k} \leq \mathbf{k}_n} \mathbf{1}\{\mathbf{B}(\mathbf{t}) \in [0, \mathbf{t}]\} \sigma_{\mathbf{n}, \mathbf{k}}^2 = \prod_{i=1}^d \sum_{k_i=1}^{k_n^i} \mathbf{1}\{b_i(k_i) \in [0, t_i]\} a_{\mathbf{n}, k_i}^i.$$

But

$$\sum_{k_i=1}^{k_n^i} \mathbf{1}\{b_i(k_i) \in [0, t_i]\} a_{n,k_i}^i = \sum_{k_i=1}^{u_i(t_i)} a_{n,k_i}^i \rightarrow t_i,$$

thus (3.71) holds due to lemma 24 and the result follows.  $\square$

The weak convergence of summation process for random variables with such variance structure was investigated by Bickel and Wichura [5] for case  $d = 2$ . They investigated weak convergence in the space of càdlàg functions. Naturally the convergence to Brownian sheet was proved.

Since in i.i.d. case we have  $\mathbf{E} X_{n,k}^2 = \pi(\mathbf{k}_n)$  this corollary then shows that theorem 29 is a generalization of invariance principle 20 in case of real valued random variables. The moment condition (3.69) in i.i.d. case then becomes

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{i \leq \mathbf{n}} \pi(\mathbf{n})^{q\alpha} \mathbf{E} |X_i / \pi(\mathbf{n})^{1/2}|^q = \lim_{m(\mathbf{n}) \rightarrow \infty} \pi(\mathbf{n})^{q(\alpha-1/2)} \mathbf{E} |X_1|^q = 0$$

for some  $q > 1/2 - \alpha$  and  $0 < \alpha < 1/2$  which holds whenever  $\mathbf{E} |X_1|^q < \infty$ . Compared to requirement  $\sup_{t>0} t^{1/2-\alpha} P(|X_1| > t)$  we see that our moment condition is not optimal, but not very far from optimality. We can further weaken it by introducing truncated variables

$$X_{n,k,\tau} := X_{n,k} \mathbf{1}\{|X_{n,k}| \leq \tau \sigma_{n,k}^{2\alpha}\}.$$

Then following theorem holds.

**Theorem 31** *Suppose there exists a function  $\mu : [0, 1]^d \rightarrow \mathbb{R}$  such that*

$$\forall \mathbf{t} \in [0, 1]^d, \quad \lim_{m(\mathbf{n}) \rightarrow \infty} \mathbf{E} \Xi_{\mathbf{n}}^2(\mathbf{t}) = \mu(\mathbf{t}). \quad (3.72)$$

*If*

$$\max_{1 \leq l \leq d} \max_{1 \leq k_l \leq k_n^l} \Delta b_l(k_l) \rightarrow 0, \text{ as } m(\mathbf{n}) \rightarrow \infty \quad (3.73)$$

*and following conditions hold:*

(a). *For every  $\varepsilon > 0$ ,*

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{1 \leq \mathbf{k} \leq \mathbf{k}_n} P(|X_{n,\mathbf{k}}| \geq \varepsilon \sigma_{n,\mathbf{k}}^{2\alpha}) = 0; \quad (3.74)$$

(b). For every  $\varepsilon > 0$ ,

$$\lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} \mathbf{E} X_{n,\mathbf{k}}^2 \mathbf{1}\{|X_{n,\mathbf{k}}| \geq \varepsilon\} = 0; \quad (3.75)$$

(c). For some  $q > 1/(1/2 - \alpha)$ ,

$$\lim_{\tau \rightarrow 0} \lim_{m(\mathbf{n}) \rightarrow \infty} \sum_{\mathbf{1} \leq \mathbf{k} \leq \mathbf{k}_n} \sigma_{n,\mathbf{k}}^{-2q\alpha} \mathbf{E} |X_{n,\mathbf{k},\tau}|^q = 0; \quad (3.76)$$

then

$$\Xi_{\mathbf{n}} \xrightarrow[m(\mathbf{n}) \rightarrow \infty]{H_{\alpha}^2([0,1]^d)} G,$$

where  $G$  is a centered Gaussian process satisfying  $\mathbf{E} G(\mathbf{t})G(\mathbf{s}) = \mu(\mathbf{t} \wedge \mathbf{s})$  for  $\mathbf{t}, \mathbf{s} \in [0, 1]^d$ .

*Proof.* The proof is identical to that of theorem 29. Just notice that the theorem conditions ensure the conditions of the theorems 28 and 19.  $\square$

# Chapter 4

## Applications

To apply our theoretical results we have to look at the examples where we can naturally assign multi-dimensional index to random observations. One of such examples is so called panel or longitudinal data, where a sample of individuals is observed over some period of time. In this case each observation has two indexes, one denoting the number of individual and another the time period at which the individual was observed. For such type of data all classical statistical problems can be discussed in a view of adjustments necessary for accomodation of the additional index. We restrict ourselves to regression and change point problems with the goal of developing the test for detecting the occurence of the change of the regression coefficient in a given sample.

We briefly recount the general setting. The classical panel data regression model which we investigate can be presented as

$$y_{it} = \alpha_i + \mathbf{x}_{it}'\boldsymbol{\beta} + u_{it}, \quad (4.1)$$

where  $i = 1, \dots, N$ ,  $t = 1, \dots, T$ ,  $y_{it}$  is an observation of dependent variable for individual  $i$  at time period  $t$ ,  $\mathbf{x}_{it}' = [x_{1it}, \dots, x_{Kit}]$  is the  $K \times 1$  vector of observations on the independent variables and  $u_{it}$  are zero mean disturbances. The classical panel regression problem is to estimate  $\boldsymbol{\beta}$  in a view of various assumptions on intercepts  $\alpha_i$ ,  $\mathbf{x}_{it}$  and  $u_{it}$ , see for example Baltagi [2], Hsiao [16]. After estimating  $\boldsymbol{\beta}$  the usual statistical procedure is to test the goodness-of-fit and the validity of the model assumptions. One of the possible violations of the validity is that relationship (4.1) holds only for certain subsample of data, i.e. the true model is

$$y_{it} = \begin{cases} \alpha_i + \mathbf{x}_{it}'\boldsymbol{\beta}_0 + u_{it}, & \text{for } (i, t) \in I, \\ \alpha_i + \mathbf{x}_{it}'\boldsymbol{\beta}_1 + u_{it}, & \text{for } (i, t) \in I^c, \end{cases} \quad (4.2)$$

with  $I \subset \{1, \dots, N\} \times \{1, \dots, T\}$ . Such violation is called change point problem. It can also appear for larger class of models, usually in parametric problems, for more general treatment see Csörgö and Horvath [7]. First test for detecting the change point in the regression setting was developed by Brown, Durbin and Evans [6], for testing the model

$$y_t = \mathbf{x}_t' \boldsymbol{\beta} + u_t, \quad (4.3)$$

against the alternative

$$y_t = \begin{cases} \mathbf{x}_t' \boldsymbol{\beta}_0 + u_t, & t = 1, \dots, t_0, \\ \mathbf{x}_t' \boldsymbol{\beta}_1 + u_t, & t = t_0 + 1, \dots, T, \end{cases} \quad (4.4)$$

where  $u_t$  are independent standard normal variables. They examined the cumulative sums of so called recursive regression residuals:

$$w_r = \frac{y_r - \mathbf{x}_r' \boldsymbol{\beta}_{r-1}}{\sqrt{1 + \mathbf{x}_r' (\sum_{k=1}^{r-1} \mathbf{x}_k \mathbf{x}_k')^{-1} \mathbf{x}_r}},$$

where  $\boldsymbol{\beta}_r$  is the least squares estimate of the model (4.3) calculated using first  $r$  observations. Suitably normalized jump sum process based on these residuals converges to Brownian motion. For the alternative model (4.4) they show the  $w_r$  no longer have zero mean, thus cumulative sum converges to infinity. The normality restriction was lifted by Sen [38], who proved similar result for the case of i.i.d. regression errors with finite variance. Ploberger and Kraämer[23] proved similar result for usual regression residuals. The limiting process in this case is the Brownian bridge.

All these three results use the same test statistic, the maximum of the cumulative sum. Since this is also a maximum norm of the jump sum process, and maximum norm is the continuous functional, due to FCLT and continuous mapping theorem, the statistic converges to maximum of limiting process (Brownian motion or Brownian bridge) under null hypothesis of no change.

Other types of alternative models are also considered. For epidemic alternative:

$$y_t = \begin{cases} \mathbf{x}_t' \boldsymbol{\beta}_0 + u_t, & t = 1, \dots, t_0, t_1 + 1, \dots, T \\ \mathbf{x}_t' \boldsymbol{\beta}_1 + u_t, & t = t_0 + 1, \dots, t_1, \end{cases} \quad (4.5)$$

Račkauskas [26] proves that it is better to use the statistic

$$\max_{1 \leq l \leq n} \frac{1}{l^\alpha} \max_{0 \leq k \leq n-l} |S(k+l) - S(k) - \frac{l}{n} S_n|,$$

where  $S(k)$  is the cumulative sum of the regression residuals. This statistic allows testing against shorter epidemics, than the usual maximum test.

In their paper Kao, Trapani and Urga [17] write “Despite the potential usefulness in economics, the econometric theory of the testing and estimation of structural changes in panels is still underdeveloped”. Current results focus on testing the change point in presence of unit roots.

In light of these results we first develop the test against epidemic rectangles using techniques from Csörgő and Horváth [7] and then apply these tests for panel regression to generalize the results of Ploberger and Krämer.

## 4.1 Tests for epidemic alternatives

### 4.1.1 Epidemic rectangles

The question arises of how to generalize epidemic alternatives for multi-indexed case. In case of panel data where we have interpretation of indexes as individuals and times several simple scenarios are immediately apparent.

- In some time interval the change occurs for all individuals.
- At the start of observation, the change occurs for certain individuals.
- At the end of observation, the change occurs for certain individuals.

For the moment assume that we are only testing the change of mean. Let  $\{X_{ij}, 1 \leq i \leq n, 1 \leq j \leq m\}$  be a sample of panel data. The null hypothesis of no change then is

$$(H_0) : X_{ij} \text{ have all the same mean } \mu_0.$$

The scenarios we want to test against fall into general setting:

$(H_A) : \text{There are integers } 1 < a^* \leq b^* < n, 1 < c^* \leq d^* < m \text{ and a constant } \mu_1 \neq \mu_0 \text{ such that}$

$$\mathbf{E} X_{ij} = \mu_0 + \mu_1 \mathbf{1} \left( (i, j) \in [a^*, b^*] \times [c^*, d^*] \cap \mathbb{N}^2 \right).$$

Classical log-likelihood statistic from Csörgő and Horváth [7] for testing change of a mean in a certain set  $I$  (if it is known in advance) is

$$R = \frac{1}{\sqrt{n}} \left( \sum_{i \in I} X_i - \frac{|I|}{n} \sum_{i=1}^n X_i \right), \quad (4.6)$$

where  $|I|$  is the cardinality of set  $I \subset \{1, \dots, n\}$ . This statistic is suitable for testing epidemics of size  $n^\gamma$ , where  $\gamma > 1/2$ . In order to test shorter epidemics you have to weight this statistic with some function of  $|I|$ .

For two-dimensional setting assume now, that integers  $a^*, b^*, c^*, d^*$  are known. Define the set

$$D^* = \left[ \frac{a^*}{n}, \frac{b^*}{n} \right] \times \left[ \frac{c^*}{m}, \frac{d^*}{m} \right]$$

and introduce the analog of statistic (4.6)

$$R = \sum_{i=1}^n \sum_{j=1}^m X_{ij} \mathbf{1} \left( \left( \frac{i}{n}, \frac{j}{m} \right) \in D^* \right) - \frac{k^* l^*}{nm} \sum_{i=1}^n \sum_{j=1}^m X_{ij}$$

where  $k^* = b^* - a^*$ ,  $l^* = d^* - c^*$ . Under hypothesis  $H_0$ , if  $X_{ij}$  are i.i.d. with finite variance  $\sigma^2$

$$(nm)^{-1/2} R \rightarrow N(0, \sigma^2 |D^*| (1 - |D^*|))$$

when  $n \wedge m \rightarrow \infty$ . Under alternative hypothesis if  $\{X_{ij}, (i/n, j/m) \in D^*\}$  and  $\{X_{ij}, (i/n, j/m) \in [0, 1]^2 \setminus D^*\}$  are separately i.i.d. but with different means, we have

$$(nm)^{-1/2} R = (nm)^{1/2} \frac{k^* l^*}{nm} \left( 1 - \frac{k^* l^*}{nm} \right) (\mu_1 - \mu_0) + O_P(1).$$

and we see that the statistic will converge to infinity as long as  $k^* \geq C_1 n^\gamma$  and  $l^* \geq C_2 m^\delta$  with  $\gamma, \delta > 1/2$  and some positive constants  $C_1, C_2$ .

In order to test shorter epidemics we have to weight the statistic  $R$ . One of the possible weights is  $\text{diam}(D)^\alpha$ , since clearly  $\text{diam}(D) \rightarrow 0$ , as  $k^*/n \rightarrow 0$  and  $l^*/m \rightarrow 0$  and vice versa.

Let us rewrite the  $Q = R / \text{diam}(D)^\alpha$  in terms of the summation process. Denote  $s_i = i/n$ ,  $t_j = j/m$ . Then

$$\begin{aligned}
Q &= \text{diam}(D)^{-\alpha} \left( \sum_{i=a^*+1}^{b^*} \sum_{j=c^*+1}^{d^*} X_{ij} - \frac{k^* l^*}{nm} \sum_{i=1}^n \sum_{j=1}^m X_{ij} \right) \\
&= \frac{\Delta_{k^*}^1 \Delta_{l^*}^2 S_{b^*, d^*} - (s_{b^*} - s_{a^*})(t_{d^*} - t_{c^*}) S_{n, m}}{\max\{s_{b^*} - s_{a^*}, t_{d^*} - t_{c^*}\}^\alpha}
\end{aligned}$$

When  $a^*, b^*, c^*, d^*$  are unknown it is reasonable to replace  $Q$  with maximum over all possible their combinations:

$$DUI(n, m, \alpha) = \max_{\substack{1 \leq a < b \leq n \\ 1 \leq c < d \leq m}} \frac{|\Delta_{b-a}^1 \Delta_{d-c}^2 S_{b, d} - (s_b - s_a)(t_d - t_c) S_{n, m}|}{\max\{s_b - s_a, t_d - t_c\}^\alpha} \quad (4.7)$$

For  $\mathbf{n} = (n, m) \in \mathbb{N}^2$ , consider the functionals  $g_{n, m}$  defined on  $H_\alpha^o([0, 1]^2)$  by

$$g_{n, m}(x) := \max_{1 \leq i < j \leq n} I(x, \mathbf{i}/\mathbf{n}, \mathbf{j}/\mathbf{n}), \quad (4.8)$$

where

$$I(x, \mathbf{s}, \mathbf{t}) = \frac{|\Delta_{t_1-s_1}^1 \Delta_{t_2-s_2}^2 x(\mathbf{t}) - (t_1 - s_1)(t_2 - s_2)x(\mathbf{1})|}{|\mathbf{t} - \mathbf{s}|^\alpha}.$$

It is clear that

$$DUI(n, m, \alpha) = g_{n, m}(\xi_{n, m}).$$

The following theorem holds.

**Theorem 32** *Functionals  $\{g_{n, m}, (n, m) \in \mathbb{N}^2\}$  and  $g$  are continuous in the Hölder space  $H_\alpha^o([0, 1]^2)$ . Furthermore  $\{g_{n, m}, (n, m) \in \mathbb{N}^2\}$  are equicontinuous and for each  $x \in H_\alpha^o([0, 1]^2)$*

$$\lim_{n \wedge m \rightarrow \infty} g_{n, m}(x) = g(x) \quad (4.9)$$

where

$$g(x) := \sup_{\mathbf{0} < \mathbf{s} < \mathbf{t} < \mathbf{1}} I(x, \mathbf{s}, \mathbf{t}). \quad (4.10)$$

*Proof.* To show continuity of functionals  $g_{n, m}$  and  $g$  and equicontinuity of family  $\{g_{n, m}, (n, m) \in \mathbb{N}^2\}$  we use lemma 9. Clearly the functional  $q = I(\cdot, \mathbf{s}, \mathbf{t})$  satisfies conditions (a) and (b) of lemma 9. Let us check condi-

tion (c). For all  $\mathbf{t}, \mathbf{s} \in [0, 1]^2$

$$\frac{(t_1 - s_1)(t_2 - s_2)}{|\mathbf{t} - \mathbf{s}|^\alpha} \leq 1.$$

Thus if  $t_1 - s_1 \leq t_2 - s_2$ , then

$$\begin{aligned} I(x, \mathbf{s}, \mathbf{t}) &\leq \frac{|x(t_1, t_2) - x(t_1, s_2)|}{|t_2 - s_2|^\alpha} + \frac{|x(s_1, t_2) - x(s_1, s_2)|}{|t_2 - s_2|^\alpha} + x(1, 1) \\ &\leq 2\|x\|_\alpha. \end{aligned} \quad (4.11)$$

Similarly if  $t_1 - s_1 > t_2 - s_2$

$$\begin{aligned} I(x, \mathbf{s}, \mathbf{t}) &\leq \frac{|x(t_1, t_2) - x(s_1, t_2)|}{|t_1 - s_1|^\alpha} + \frac{|x(t_1, s_2) - x(s_1, s_2)|}{|t_1 - s_1|^\alpha} + x(1, 1) \\ &\leq 2\|x\|_\alpha. \end{aligned} \quad (4.12)$$

So functional  $I(\cdot, \mathbf{s}, \mathbf{t})$  satisfies condition (c) with  $C = 2$ . Thus the continuity and equicontinuity follows immediately from (1.7).

For (4.9) it is sufficient to show that the function  $(\mathbf{s}, \mathbf{t}) \rightarrow I(x, \mathbf{s}, \mathbf{t})$  can be extended by continuity to the compact set  $T = \{(\mathbf{s}, \mathbf{t}) \in [0, 1]^4; \mathbf{0} \leq \mathbf{s} \leq \mathbf{t} \leq \mathbf{1}\}$ . From (4.11) and (4.12) we get  $0 \leq I(x, \mathbf{s}, \mathbf{t}) \leq 2w_\alpha(x, |\mathbf{t} - \mathbf{s}|) + |\mathbf{t} - \mathbf{s}|^{1-\alpha}x(1, 1)$ , which allows continuous extension when  $\mathbf{t} = \mathbf{s}$  putting  $I(x, \mathbf{s}, \mathbf{s}) = 0$ .  $\square$

Functionals  $g_{n,m}$  and  $g$  satisfy the conditions of lemma 8 thus FCLT for  $X_{ij}$  gives the limiting distribution of statistic  $DUI(n, m, \alpha)$ . Due to results in previous sections, the FCLT in the space  $H_\alpha^0([0, 1]^2)$  holds for the summation processes based on i.i.d. random variables. Thus we have to strengthen the null hypothesis:

$(H'_0) : X_{ij}$  are independent identically distributed with mean denoted by  $\mu_0$ .

For better clarity for any real function  $x$  with two dimensional argument introduce definition

$$\Delta_{[\mathbf{s}, \mathbf{t}]}x := x(\mathbf{t}) - x(s_1, t_2) - x(t_1, s_2) + x(\mathbf{s}).$$

This sometimes is called the increment of  $x$  around the rectangle  $[\mathbf{s}, \mathbf{t}]$ . Consider the following random variable

$$DUI(\alpha) = \sup_{\mathbf{0} \leq \mathbf{s} < \mathbf{t} \leq \mathbf{1}} \frac{|\Delta_{[\mathbf{s}, \mathbf{t}]}W - (t_1 - s_1)(t_2 - s_2)W(\mathbf{1})|}{|\mathbf{t} - \mathbf{s}|^\alpha}. \quad (4.13)$$

Then following theorem holds.

**Theorem 33** Under  $H'_0$  assume that  $0 < \alpha < 1/2$  and

$$\sup_{t>0} t^p P(|X_1| > t) > \infty$$

for  $p = 1/(1/2 - \alpha)$ . Then

$$\sigma^{-1}(nm)^{-1/2} DUI(n, m, \alpha) \xrightarrow{D} DUI(\alpha), \text{ as } n \wedge m \rightarrow \infty,$$

where  $\sigma^2 = \mathbf{E} X_1^2$ .

*Proof.* Note first, that under  $H'_0$  the value of statistic  $DUI(n, m, \alpha)$  does not change if  $X_i$  are exchanged with  $X_i - \mu_0$ . Assume then that  $\mu_0 = 0$ . Theorem 21 together with theorem 32 and lemma 8 gives us the result.  $\square$

The consistency of the test is given by following theorem.

**Theorem 34** Assume under  $(H_A)$  that the  $X_{ij}$  are independent and  $\sigma_0^2 = \sup_n \text{var}(X_n)$  is finite. If

$$\lim_{m(n) \rightarrow \infty} (nm)^{1/2} \frac{h_{n,m}}{d_{n,m}^\alpha} |\mu_1 - \mu_0| \rightarrow \infty, \quad (4.14)$$

where

$$h_{n,m} = \frac{k^* l^*}{nm} \left( 1 - \frac{k^* l^*}{nm} \right) \text{ and } d_{n,m} = \max \left\{ \frac{k^*}{n}, \frac{l^*}{m} \right\}, \quad (4.15)$$

then

$$(nm)^{-1/2} DUI(n, m, \alpha) \rightarrow \infty. \quad (4.16)$$

For the case  $d = 1$  our result replicates the result of Račkauskas and Suquet [32]. In this case the test will be able to detect epidemics of order  $n^{\frac{1-2\alpha}{2-2\alpha}}$ . Now for two dimensional case assume that  $k^* = n^\gamma$ ,  $l^* = m^\delta$  and that  $\mu_1 - \mu_0$  does not depend on  $(n, m)$ . Then the condition (4.14) becomes

$$\frac{n^{\gamma-1/2} m^{\delta-1/2}}{[n^{\gamma-1} \vee m^{\delta-1}]^\alpha} \rightarrow \infty.$$

If  $n^{\gamma-1} > m^{\delta-1}$  (4.14) reduces to

$$n^{\gamma(1-\alpha)+\alpha-1/2} m^{\delta-1/2} \rightarrow \infty,$$

thus we can detect very short epidemics of  $k^*$ , but we can never get better rate for epidemic length of  $l^*$ . Notice that for  $\delta > 1/2$  condition  $n^{\gamma-1} > m^{\delta-1}$  will be satisfied if  $m > n^2$ . So we see that to detect very short epidemics for one index we must have more data for the other index. In case  $n = m$ ,  $\gamma = \delta$  we have

$$n^{\gamma(1-\alpha)+\alpha-1/2+\gamma-1/2} = n^{\gamma(2-\alpha)+\alpha-1} \rightarrow \infty,$$

and the best rate is  $\gamma > 1/3$ . We get that in two-dimensional case the rates of epidemic are influenced not only by  $\alpha$ , but also by the relationship between  $n$  and  $m$ .

The choice of  $\alpha$  is important in the convergence of null hypothesis. In one dimensional case for the convergence we have the condition  $\lim_{t \rightarrow \infty} t^p P(|X_1| > t) = 0$ , where  $p = 1/(1/2 - \alpha)$ . Since  $\frac{1-2\alpha}{2-2\alpha} \rightarrow 0$ , when  $\alpha \rightarrow 1/2$ , we get better rates with higher moment conditions.

For case  $m = n^2$  the moment condition for convergence is  $\lim_{t \rightarrow \infty} t^{\frac{3}{3/2-2\alpha}} P(|X_1| > t) = 0$ . Thus  $\mathbf{E} X_1^6$  is sufficient.

For case  $m = n$  from theorem 22 it follows that  $4^{th}$  moment is sufficient for convergence for any  $\alpha$ , but the rate cannot be lower than  $1/3$ . In one dimensional case detecting an epidemic of length  $n^{1/3}$  comes with the choice  $\alpha = 1/4$ , which means that we need  $4^{th}$  moment.

### Proof of the theorem 34

Define set  $I_{n,m} = [a^*, b^*] \times [c^*, d^*] \cap \mathbb{N}^2$  and random variables

$$X_{ij} := \begin{cases} X_{ij} - \mu_0, & (i, j) \in I_{n,m}^c \\ X_{ij} - \mu_1, & (i, j) \in I_{n,m} \end{cases}$$

We have

$$\begin{aligned} \Delta_{k^*}^1 \Delta_{l^*}^2 S_{b^*, d^*} - (s_{b^*} - s_{a^*})(t_{d^*} - t_{c^*}) S_{n,m} &= S(I_{n,m}) - \frac{k^* l^*}{nm} (S(I_{n,m}) + S(I_{n,m}^c)) \\ &= k^* l^* \left( 1 - \frac{k^* l^*}{nm} \right) (\mu_1 - \mu_0) + R_{n,m}, \end{aligned} \quad (4.17)$$

where

$$R_{n,m} := -\frac{k^* l^*}{nm} \sum_{i \in I_{n,m}^c} X'_i + \left( 1 - \frac{k^* l^*}{nm} \right) \sum_{i \in I_{n,m}} X'_i.$$

Now

$$\begin{aligned} \text{var}((nm)^{-1/2}R_{n,m}) &\leq \frac{1}{nm} \left( \frac{k^*l^*}{nm} \right)^2 (nm - k^*l^*)\sigma_0^2 \\ &\quad + \frac{1}{nm} \left( 1 - \frac{k^*l^*}{nm} \right)^2 k^*l^*\sigma_0^2 = \sigma_0^2 h_{n,m}. \end{aligned}$$

This estimate together with (4.17) leads to the lower bound

$$(nm)^{-1/2}DUI(n, m, \alpha) \geq (nm)^{1/2} \frac{h_{n,m}}{d_{n,m}^\alpha} |\mu_1 - \mu_0| + O_P \left( \frac{h_{n,m}^{1/2}}{d_{n,m}^\alpha} \right).$$

Now  $h_{n,m} \leq d_{n,m}^2$ , thus  $\lim_{d_{n,m} \rightarrow 0} h_{n,m}^{1/2}/d_{n,m}^\alpha = 0$ , so the theorem follows due to condition (4.14).

#### 4.1.2 Some special cases

In previous section we constructed statistic for detecting the change in subrectangle of unit square. Our motivation for such statistic came from three simple scenarios:

- S1. At the start of observation, the change occurs for certain individuals
- S2. At the end of observation, the change occurs for certain individuals.
- S3. In some time interval the change occurs for all individuals

Using results from the previous section we can adapt the general statistic  $DUI(n, m, \alpha)$  for each of these scenarios. Recall that the alternative hypothesis was of the change in an epidemic rectangle

$$D^* = \left[ \frac{a^*}{n}, \frac{b^*}{n} \right] \times \left[ \frac{c^*}{m}, \frac{d^*}{m} \right].$$

Then the respective epidemic rectangles for the scenarios are

$$\begin{aligned} D_1^* &= \left[ 0, \frac{b^*}{n} \right] \times \left[ 0, \frac{d^*}{m} \right], \\ D_2^* &= \left[ \frac{a^*}{n}, 1 \right] \times \left[ \frac{c^*}{m}, 1 \right], \\ D_3^* &= [0, 1] \times \left[ \frac{c^*}{m}, \frac{d^*}{m} \right]. \end{aligned}$$

Denote  $s_i = i/n$ ,  $t_j = j/m$ . Then the statistic adapted for specific first scenario is

$$DUI_1(n, m, \alpha) = \max_{\substack{1 \leq b \leq n \\ 1 \leq d \leq m}} \frac{|S_{b,d} - s_b t_d S_{n,m}|}{\max\{s_b, t_d\}^\alpha},$$

since  $S_{0,d} = S_{0,0} = S_{b,0} = 0$ , and  $s_a = t_c = 0$ . For the second scenario we have

$$DUI_2(n, m, \alpha) = \max_{\substack{1 \leq b \leq n \\ 1 \leq d \leq m}} \frac{|\Delta_{n-a}^1 \Delta_{m-c}^2 S_{n,m} - (1 - s_a)(1 - t_c) S_{n,m}|}{\max\{(1 - s_a), (1 - t_c)\}^\alpha}.$$

For the third scenario the statistic is defined as

$$DUI_3(n, m, \alpha) = \max_{1 \leq c < d \leq m} \frac{|S_{n,d} - S_{n,c} - (t_d - t_c) S_{n,m}|}{(t_d - t_c)^\alpha},$$

where we changed the denominator, since  $S_{0,c} = S_{0,d} = 0$  and in the nominator only the difference of the second argument matters. All the three statistics are the functionals of summation process  $\xi_{n,m}$  similar to functional  $g_n$  defined by (4.10). With minimal adaptation similar proposition to the theorem 32 holds. Define following functionals of Brownian sheet

$$\begin{aligned} DUI_1(\alpha) &= \sup_{0 < t < 1} \frac{|W(\mathbf{t}) - t_1 t_2 W(1, 1)|}{|\mathbf{t}|^\alpha} \\ DUI_2(\alpha) &= \sup_{0 < t < 1} \frac{|\Delta_{[t,1]} W - (1 - t_1)(1 - t_2) W(\mathbf{1})|}{|\mathbf{1} - \mathbf{t}|^\alpha}, \\ DUI_3(\alpha) &= \sup_{0 < s < t < 1} \frac{|W(1, t) - W(1, s) - (t - s) W(1, 1)|}{|t - s|^\alpha}. \end{aligned}$$

Then following theorem holds.

**Theorem 35** *For i.i.d. sample of double-indexed data and under null hypothesis of no change for scenarios S1, S2 and S3 assume that  $0 < \alpha < 1/2$  and*

$$\sup_{t>0} t^p P(|X_1| > t) > \infty,$$

for  $p = 1/(1/2 - \alpha)$ . Then for  $i = 1, 2, 3$ .

$$\sigma^{-1}(nm)^{-1/2} DUI_i(n, m, \alpha) \xrightarrow{D} DUI_i(\alpha), \text{ as } n \wedge m \rightarrow \infty,$$

where  $\sigma^2 = \mathbf{E} X_1^2$ .

Under the alternative hypothesis of the change of the mean in the rectangles  $D_i^*$ ,  $i = 1, 2, 3$ , the lengths of epidemics are

$$\begin{aligned} k^* &= b^*, \quad l^* = d^*, \text{ for the rectangle } D_1^*, \\ k^* &= 1 - a^*, \quad l^* = 1 - c^*, \text{ for the rectangle } D_2^*, \\ k^* &= n, \quad l^* = d^* - c^*, \text{ for the rectangle } D_3^*. \end{aligned}$$

For the rectangles  $D_1^*$  and  $D_2^*$  the consistency is then the direct corollary of the theorem (34).

**Corollary 36** *Given the independent family  $X_{ij}$  with  $\sigma_0^2 = \sup_n \text{var}(X_n)$  finite under alternative hypothesis of the change of the mean in rectangles  $D_1^*$  and  $D_2^*$  we have*

$$(nm)^{-1/2} DUI_i(n, m, \alpha) \rightarrow \infty \quad (4.18)$$

for  $i = 1, 2$  if

$$\lim_{m(n) \rightarrow \infty} (nm)^{1/2} \frac{h_{n,m}}{d_{n,m}^\alpha} |\mu_1 - \mu_0| \rightarrow \infty, \quad (4.19)$$

where

$$h_{n,m} = \frac{k^* l^*}{nm} \left( 1 - \frac{k^* l^*}{nm} \right) \quad \text{and} \quad d_{n,m} = \max \left\{ \frac{k^*}{n}, \frac{l^*}{m} \right\}. \quad (4.20)$$

For the rectangle  $D_3^*$  the conditions for the consistency are slightly different, since the denominator in the test statistic is different.

**Corollary 37** *Given the independent family  $X_{ij}$  with  $\sigma_0^2 = \sup_n \text{var}(X_n)$  finite under alternative hypothesis of the change of the mean in rectangle  $D_3^*$  we have*

$$(nm)^{-1/2} DUI_3(n, m, \alpha) \rightarrow \infty, \quad (4.21)$$

if

$$\lim_{m(n) \rightarrow \infty} (nm)^{1/2} \left( \frac{l^*}{m} \right)^{1-\alpha} \left( 1 - \frac{l^*}{m} \right) |\mu_1 - \mu_0| \rightarrow \infty. \quad (4.22)$$

This corollary enables us to improve the detection of short epidemics for the scenario **S3**. If we let  $l^* = m^\delta$  for some  $\delta > 0$ , condition (4.22) becomes

$$n^{1/2} m^{\delta(1-\alpha)+\alpha-1/2} (1 - m^{\delta-1}) \rightarrow \infty,$$

for  $\delta > \frac{1/2-\alpha}{1-\alpha}$ . Since  $0 < \alpha < 1/2$  we can make  $\delta$  arbitrarily small. Thus for certain rectangles it is possible to get similar results as in one dimensional case.

## 4.2 Functional central limit theorems for panel data regressions

### 4.2.1 Models and the assumptions

Suppose we have a sample of panel data  $\{(y_{ij}, \mathbf{x}'_{ij}), i = 1, \dots, n; j = 1, \dots, m\}$  where  $\mathbf{x}'_{ij} = (x_{ij1}, \dots, x_{ijK})$ . We investigate FCLT for following panel regression models

$$y_{ij} = \mathbf{x}'_{ij}\boldsymbol{\beta} + u_{ij}, \quad (4.23)$$

$$y_{ij} = \mathbf{x}'_{ij}\boldsymbol{\beta} + \mu_i + u_{ij}, \quad (4.24)$$

where  $u_{ij}$  are the disturbances, mean-zero random variables with finite variance independent of  $y_{ij}$  and  $\mathbf{x}_{ij}$ .

The goal of panel regression is to estimate coefficient vector  $\boldsymbol{\beta}$ . The model (4.23) is the classical linear regression model for observations with two dimensional indexes. The coefficients  $\boldsymbol{\beta}$  are then usually estimated using least squares:

$$\hat{\boldsymbol{\beta}} = \left( \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}'_{ij} \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} y_{ij}. \quad (4.25)$$

In classical panel data literature (Baltagi [2], Hsiao [16]) this estimate is called pooled or ordinary least squares estimate, and it is assumed that  $x_{ij1} = 1$  for all  $i, j$ , i.e. there is only one constant term. For the model (4.24) the constant term is allowed to vary through  $i$  and is considered as a nuisance parameter. The coefficient vector  $\boldsymbol{\beta}$  in this case is estimated by solving least squares problem for the model

$$y_{ij} - \bar{y}_i = (\mathbf{x}_{ij} - \bar{\mathbf{x}}_i)\boldsymbol{\beta} + u_{ij} - \bar{u}_i, \quad (4.26)$$

where

$$\bar{y}_{i.} = \frac{1}{m} \sum_{j=1}^m y_{ij}, \quad \bar{\mathbf{x}}_{i.} = \frac{1}{m} \sum_{j=1}^m \mathbf{x}_{ij}, \quad \bar{u}_{i.} = \frac{1}{m} \sum_{j=1}^m u_{ij}.$$

The estimate of  $\boldsymbol{\beta}$  is then

$$\hat{\boldsymbol{\beta}}^{FE} = \left( \sum_{i=1}^n \sum_{j=1}^m (\mathbf{x}_{ij} - \bar{\mathbf{x}}_{i.})(\mathbf{x}_{ij} - \bar{\mathbf{x}}_{i.})' \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m (\mathbf{x}_{ij} - \bar{\mathbf{x}}_{i.})(y_{ij} - \bar{y}_{i.}). \quad (4.27)$$

We are interested in proving FCLT for regression residuals. For model (4.23) they are defined as

$$\hat{u}_{ij} = y_{ij} - \mathbf{x}_{ij}' \hat{\boldsymbol{\beta}} = u_{ij} - \mathbf{x}_{ij}' (\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}).$$

By substituting the expression for  $y_{ij}$  we immediately get

$$\hat{u}_{ij} = u_{ij} - \mathbf{x}_{ij}' (\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}). \quad (4.28)$$

Now

$$\hat{\boldsymbol{\beta}} - \boldsymbol{\beta} = \left( \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}_{ij}' \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} u_{ij},$$

and we see that FCLT for regression residuals depends on distributional properties of regression disturbances. For this section let us make the following assumption.

**Assumption F** *Let random variables  $u_{ij}$  have zero mean, variance  $\sigma^2$  and be independent of  $\mathbf{x}_{ij}$ . Assume that the summation process based on these random variables defined as*

$$\xi_{n,m}(t, s) = nm \sum_{i=1}^n \sum_{j=1}^m \left[ \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, s] \right] u_{ij}, \quad (4.29)$$

*satisfies the functional central limit theorem:*

$$\frac{1}{\sigma \sqrt{nm}} \xi_{n,m}(t, s) \xrightarrow{D} W(t, s), \text{ as } n \wedge m \rightarrow \infty,$$

*in the space  $H_\alpha^2([0, 1]^2)$  with  $0 < \alpha < 1/2$ .*

From (4.28) it is also evident that we have to make assumptions on  $\mathbf{x}_{ij}$  in the panel regression model (4.23).

**Assumption A** Let  $x_{ij1} = 1$  for all  $1 \leq (i, j) \leq (n, m)$ . Assume that

$$\lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}_{ij}' = R, \quad (4.30)$$

for some nonsingular  $(K \times K)$  matrix  $R$ . Furthermore assume that the model is reparameterized such that

$$R = \begin{bmatrix} 1 & 0 \\ 0 & R^* \end{bmatrix}, \quad (4.31)$$

which in turn implies that

$$\mathbf{c} \equiv \lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} = [1, 0, \dots, 0]'. \quad (4.32)$$

Assumption **F** implies that  $u_{ij}$  satisfy central limit theorem, which together with assumption **A** implies that

$$\frac{1}{\sqrt{nm}} \left( \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}_{ij}' \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} u_{ij} \xrightarrow{D} N(0, \sigma^2 R), \text{ as } n \wedge m \rightarrow \infty. \quad (4.33)$$

Assumption **A** also implies that

$$\frac{1}{nm} \sum_{i=1}^{[nt]} \sum_{j=1}^{[ms]} \mathbf{x}_{ij} \rightarrow t s \mathbf{c}, \text{ as } n \wedge m \rightarrow \infty,$$

for each fixed  $t$  and  $s$ . Using results from section 2.1 we get that summation process based on  $\mathbf{x}_{ij}$  also has the same limit

$$\begin{aligned} \mathbf{X}_{n,m}(t, s) &= \frac{1}{nm} \sum_{j=1}^m nm \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, s] \right| \mathbf{x}_{ij} \\ &\rightarrow t s \mathbf{c} \end{aligned} \quad (4.34)$$

for each fixed  $t$  and  $s$ . But assumption **A** also ensures that  $\{\mathbf{X}_{n,m}, (n, m) \in \mathbb{N}^2\}$  is equicontinuous in Hölder space  $H_\alpha^o([0, 1]^2, \mathbb{R}^K)$ . Thus we get that

$$\mathbf{X}_{n,m}(t, s) \rightarrow t s \mathbf{c}, \text{ as } n \wedge m \rightarrow \infty. \quad (4.35)$$

in  $H_\alpha^o([0, 1]^2, \mathbb{R}^K)$ . We can write

$$\sqrt{nm}(\hat{\beta} - \beta) = \begin{bmatrix} 1 & 0 \\ 0 & R^* \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{nm}} \sum_{i=1}^n \sum_{j=1}^m u_{ij} \\ \frac{1}{\sqrt{nm}} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij}^* u_{ij} \end{bmatrix} + o_P(1)$$

where  $\mathbf{x}_{ij}^* = [x_{ij2}, \dots, x_{ijK}]'$ . This representation together with (4.34) gives us the convergence

$$\begin{aligned} \frac{1}{\sqrt{nm}} \sum_{j=1}^m nm \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, s] \right| \mathbf{x}_{ij}' (\hat{\beta} - \beta) \\ \xrightarrow{D} tsW(1, 1), \end{aligned}$$

as  $n \wedge m \rightarrow \infty$  in the space  $H_\alpha^o([0, 1]^2)$ . We have proved the following theorem.

**Theorem 38** *For the panel regression model*

$$y_{ij} = \mathbf{x}_{ij}' \beta + u_{ij}, \quad (4.36)$$

define the summation process

$$\widehat{W}^{(n,m)}(t, s) = nm \sum_{i=1}^n \sum_{j=1}^m \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, s] \right| \widehat{u}_{ij}.$$

Given the assumptions **F** and **A** we have

$$\frac{1}{\sigma \sqrt{nm}} \widehat{W}^{(n,m)}(t, s) \xrightarrow{D} W(t, s) - tsW(1, 1), \text{ as } n \wedge m \rightarrow \infty,$$

in the space  $H_\alpha^o([0, 1]^2)$ , with  $0 < \alpha < 1/2$ .

Let us turn now to the model (4.24). Since the fixed effect estimate  $\hat{\beta}^{FE}$  comes from the adjusted regression (4.26) it is natural to define residuals as

$$\widehat{u}^{FE} = \widetilde{y}_{ij} - \widetilde{\mathbf{x}}_{ij}' \hat{\beta}^{FE},$$

where  $\widetilde{y}_{ij} = y_{ij} - \bar{y}_{i\cdot}$ , and  $\widetilde{\mathbf{x}}_{ij}$  is defined analogously. Substituting the model (4.24) we get that

$$\widehat{u}_{ij}^{FE} = u_{ij} - \bar{u}_{i\cdot} - \widetilde{\mathbf{x}}_{ij}' (\hat{\beta}^{FE} - \beta).$$

For model (4.24) we make slightly different assumptions about  $\mathbf{x}_{ij}$ .

**Assumption B** Assume that

$$\lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \tilde{\mathbf{x}}'_{ij} = \tilde{R} \quad (4.37)$$

for some nonsingular  $(K \times K)$  matrix  $\tilde{R}$ , and that

$$\lim_{n \wedge m \rightarrow \infty} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} = \mathbf{c} \quad (4.38)$$

for some  $\mathbf{c} \in \mathbb{R}^K$ .

Now

$$\begin{aligned} \hat{\beta}^{FE} - \beta &= \left( \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \tilde{\mathbf{x}}'_{ij} \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} (u_{ij} - \bar{u}_{i\cdot}) \\ &= \left( \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \tilde{\mathbf{x}}'_{ij} \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} u_{ij}, \end{aligned}$$

since

$$\sum_{j=1}^m \tilde{\mathbf{x}}_{ij} = 0.$$

Thus assumption (F) gives us

$$\sqrt{nm}(\hat{\beta}^{FE} - \beta) \xrightarrow{D} N(0, \sigma^2 \tilde{R}).$$

From condition (4.38) we get

$$\frac{1}{nm} \sum_{i=1}^{[nt]} \sum_{j=1}^{[ms]} \tilde{\mathbf{x}}_{ij} \rightarrow 0,$$

for each fixed  $t$  and  $s$ . Then similar to (4.35) for summation process  $\widetilde{\mathbf{X}}_{n,m}(t, s)$  based on  $\tilde{\mathbf{x}}_{ij}$  we get that

$$\widetilde{\mathbf{X}}_{n,m}(t, s) \rightarrow 0, \text{ as } n \wedge m \rightarrow \infty, \quad (4.39)$$

in  $H_\alpha^o([0, 1]^2, \mathbb{R}^K)$ . Now relationships

$$\begin{aligned} & \sum_{i=1}^n \sum_{j=1}^m nm \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, s] \right| \bar{u}_i. \\ &= \sum_{i=1}^n \sum_{l=1}^m n \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \cap [0, t] \right| u_{il} \sum_{j=1}^m \left| \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, s] \right| \\ &= s \sum_{i=1}^n \sum_{j=1}^m nm \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, 1] \right| u_{ij} \end{aligned}$$

complete the proof of the following theorem.

**Theorem 39** *For the panel regression model*

$$y_{ij} = \mathbf{x}_{ij}' \boldsymbol{\beta} + \mu_i + u_{ij}, \quad (4.40)$$

define the summation process

$$\widehat{W}_{nm}^{FE}(t, s) = nm \sum_{i=1}^n \sum_{j=1}^m \left| \left[ \frac{i-1}{n}, \frac{i}{n} \right) \times \left[ \frac{j-1}{m}, \frac{j}{m} \right) \cap [0, t] \times [0, s] \right| \widehat{u}_{ij}^{FE}.$$

Given the assumptions *F* and *B*, we have

$$\widehat{W}_{nm}^{FE}(t, s) \xrightarrow{D} W(t, s) - sW(t, 1), \text{ as } n \wedge m \rightarrow \infty,$$

in the space  $H_\alpha^o([0, 1]^2)$ , with  $0 < \alpha < 1/2$ .

## 4.2.2 Local alternatives

It is possible to get meaningful results if we alter the original regression models. Assume that coefficient  $\boldsymbol{\beta}$  in the pooled regression actually varies across  $i$  and  $j$ :

$$\boldsymbol{\beta}_{ij} = \boldsymbol{\beta} + \frac{1}{\sqrt{nm}} \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right) \quad (4.41)$$

where  $\mathbf{g}$  is  $K$ -vector valued function continuous on  $[0, 1]^2$ .

As  $n \wedge m \rightarrow \infty$  this alternative to the regression model

$$y_{ij}^{loc} = \mathbf{x}_{ij}' \boldsymbol{\beta}_{ij} + u_{ij},$$

converges to model (4.23). Define least squares estimate for this regression

as

$$\hat{\beta}^{loc} = \left( \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}_{ij}' \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} y_{ij}^{loc}.$$

The following theorem then holds

**Theorem 40** Define summation process  $\widehat{W}^{(n,m)}$  based on residuals

$$u_{ij}^{loc} = y_{ij}^{loc} - \mathbf{x}_{ij}' \hat{\beta}^{loc}.$$

Given the assumptions **F** and **A** we have

$$\begin{aligned} \widehat{W}^{(n,m)}(t, s) &\xrightarrow{D} W(t, s) - tsW(1, 1) \\ &\quad + \int_0^t \int_0^s \mathbf{c}' \mathbf{g}(u, v) \, du \, dv - ts \mathbf{c}' \int_0^1 \int_0^1 \mathbf{g}(u, v) \, du \, dv. \end{aligned}$$

*Proof.* Note that

$$\hat{\beta}^{loc} = \hat{\beta} + \frac{1}{\sqrt{nm}} \mathbf{d}_{n,m},$$

where

$$\mathbf{d}_{n,m} = \left( \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}_{ij}' \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}_{ij}' \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right).$$

Thus we can decompose the residuals  $u_{ij}^{loc}$  into following sums:

$$\begin{aligned} u_{ij}^{loc} &= u_{ij} + \mathbf{x}_{ij}' \beta_{ij} - \mathbf{x}_{ij}' \hat{\beta}^{loc} \\ &= u_{ij} - \mathbf{x}_{ij}' (\hat{\beta} - \beta) + \frac{1}{\sqrt{nm}} \mathbf{x}_{ij}' \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right) - \frac{1}{\sqrt{nm}} \mathbf{x}_{ij}' \mathbf{d}_{n,m} \\ &= \hat{u}_{ij} + \frac{1}{\sqrt{nm}} \mathbf{x}_{ij}' \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right) - \frac{1}{\sqrt{nm}} \mathbf{x}_{ij}' \mathbf{d}_{n,m}. \end{aligned}$$

Using assumption **A** and properties of  $\mathbf{g}$  for each fixed  $t$  and  $s$  we get

$$\frac{1}{nm} \sum_{i=1}^{[nt]} \sum_{j=1}^{[ms]} \mathbf{x}_{ij}' \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right) \rightarrow \int_0^t \int_0^s \mathbf{c}' \mathbf{g}(u, v) \, du \, dv, \quad (4.42)$$

as  $n \wedge m \rightarrow \infty$ . Using the same arguments as in (4.34) we get that summation process based on  $\mathbf{x}_{ij}' \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right)$  has the same limit in Hölder space  $H_\alpha^0([0, 1]^2)$ .

Similarly we get

$$\mathbf{d}_{n,m} \rightarrow \int_0^1 \int_0^1 \mathbf{g}(u, v) \, du \, dv, \text{ as } n \wedge m \rightarrow \infty. \quad (4.43)$$

Since  $\mathbf{d}_{n,m}$  does not depend on  $i$  and  $j$ , the convergence (4.34) and theorem 38 complete the proof.  $\square$

Consider the same local alternatives (4.41) for the fixed-effects panel regression. Then the alternative model is

$$y_{ij}^{loc} = \mu_i + \mathbf{x}_{ij}' \beta_{ij} + u_{ij}. \quad (4.44)$$

Define analogously fixed effect estimate

$$\beta_{loc}^{FE} = \left( \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \tilde{\mathbf{x}}_{ij}' \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \tilde{y}_{ij}^{loc}. \quad (4.45)$$

Then the following theorem holds.

**Theorem 41** Define summation process  $\widehat{W}^{(n,m)}$  based on residuals

$$\tilde{u}_{ij}^{loc} = \tilde{y}_{ij}^{loc} - \tilde{\mathbf{x}}_{ij}' \beta_{loc}^{FE}.$$

Given the assumptions *F* and *B* we have

$$\begin{aligned} \widehat{W}^{(n,m)}(t, s) &\xrightarrow{D} W(t, s) - sW(t, 1) \\ &\quad + \int_0^t \int_0^s \mathbf{c}' \mathbf{g}(u, v) \, du \, dv - s \int_0^t \int_0^1 \mathbf{c}' \mathbf{g}(u, v) \, du \, dv, \end{aligned}$$

as  $n \wedge m \rightarrow \infty$ , in the space  $H_\alpha^\circ([0, 1]^2)$ , with  $0 < \alpha < 1/2$ .

*Proof.* Introduce definitions

$$\mathbf{g}_{ij} = \mathbf{g}\left(\frac{i}{n}, \frac{j}{m}\right), \quad \bar{\mathbf{f}}_{i\cdot} = \frac{1}{m} \sum_{j=1}^m \mathbf{x}_{ij}' \mathbf{g}_{ij}.$$

Then for our alternative model (4.44) we get

$$\tilde{y}_{ij}^{loc} = \tilde{\mathbf{x}}_{ij}' \beta + \tilde{u}_{ij} + \frac{1}{\sqrt{nm}} \mathbf{x}_{ij}' \mathbf{g}_{ij} - \frac{1}{\sqrt{nm}} \bar{\mathbf{f}}_{i\cdot}.$$

Then from the definition (4.45) it follows that

$$\boldsymbol{\beta}_{loc}^{FE} = \hat{\boldsymbol{\beta}}^{FE} + \frac{1}{\sqrt{nm}} \mathbf{d}_{n,m},$$

where

$$\mathbf{d}_{n,m} = \left( \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \tilde{\mathbf{x}}_{ij}' \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} (\mathbf{x}_{ij}' \mathbf{g}_{ij} - \bar{\mathbf{f}}_{i\cdot}).$$

Substituting these expressions to the definition of the residuals we obtain

$$\begin{aligned} \hat{u}_{ij}^{loc} &= \tilde{\mathbf{x}}_{ij}' \boldsymbol{\beta} + \tilde{u}_{ij} + \frac{1}{\sqrt{nm}} (\mathbf{x}_{ij}' \mathbf{g}_{ij} - \bar{\mathbf{f}}_{i\cdot}) - \tilde{\mathbf{x}}_{ij}' \boldsymbol{\beta}_{loc}^{FE} \\ &= \tilde{u}_{ij} - \tilde{\mathbf{x}}_{ij}' (\hat{\boldsymbol{\beta}}^{FE} - \boldsymbol{\beta}) + \frac{1}{\sqrt{nm}} (\mathbf{x}_{ij}' \mathbf{g}_{ij} - \bar{\mathbf{f}}_{i\cdot}) - \frac{1}{\sqrt{nm}} \tilde{\mathbf{x}}_{ij}' \mathbf{d}_{n,m} \\ &= \hat{u}_{ij}^{FE} + \frac{1}{\sqrt{nm}} (\mathbf{x}_{ij}' \mathbf{g}_{ij} - \bar{\mathbf{f}}_{i\cdot}) - \frac{1}{\sqrt{nm}} \tilde{\mathbf{x}}_{ij}' \mathbf{d}_{n,m}. \end{aligned}$$

Now given assumption B, similar to (4.42) we get that

$$\frac{1}{nm} \sum_{i=1}^{[nt]} \sum_{j=1}^{[ms]} \mathbf{x}_{ij}' \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right) \rightarrow \int_0^t \int_0^s \mathbf{c}' \mathbf{g}(u, v) \, du \, dv,$$

as  $n \wedge m \rightarrow \infty$  for each fixed  $t$  and  $s$ . Similarly

$$\frac{1}{nm} \sum_{i=1}^{[nt]} \sum_{j=1}^{[ms]} \bar{\mathbf{f}}_{i\cdot} = \frac{1}{nm} \sum_{i=1}^{[nt]} \sum_{j=1}^m \mathbf{x}_{ij}' \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right) \rightarrow s \int_0^t \int_0^1 \mathbf{c}' \mathbf{g}(u, v) \, du \, dv,$$

as  $n \wedge m \rightarrow \infty$  for each fixed  $t$  and  $s$ . From

$$\sum_{j=1}^m \tilde{\mathbf{x}}_{ij} = 0$$

it follows that

$$\frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \bar{\mathbf{f}}_{i\cdot} = 0 \text{ and } \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \bar{\mathbf{x}}_{i\cdot}' = 0.$$

Since  $\mathbf{g}$  is continuous on  $[0, 1]^2$ , it is bounded:

$$\sup_{t,s \in [0,1]^2} |\mathbf{g}(t, s)| \leq C.$$

Then

$$\begin{aligned} \mathbf{d}_{n,m} &= \left( \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \tilde{\mathbf{x}}'_{ij} \right)^{-1} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} (\mathbf{x}_{ij} g_{ij} - \bar{\mathbf{f}}_i) \\ &= \left( \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \tilde{\mathbf{x}}'_{ij} \right)^{-1} \frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} (\tilde{\mathbf{x}}'_{ij} g_{ij} + \bar{\mathbf{x}}'_i g_{ij}) \leq C \end{aligned}$$

and similar to the proof of the theorem 40 the convergence (4.39) and theorem 38 complete the proof.  $\square$

## 4.3 Change point statistics for panel regressions

### 4.3.1 Tests and their behaviour under null hypothesis

Combining results from previous sections we can now suggest statistics for detecting the change point in panel regression models and give their limiting distributions. Under assumption F and respective assumptions A and B the residuals for our regression models

$$y_{ij} = \mathbf{x}'_{ij} \boldsymbol{\beta} + u_{ij}, \quad (4.46)$$

$$y_{ij} = \mathbf{x}'_{ij} \boldsymbol{\beta} + \mu_i + u_{ij}, \quad (4.47)$$

satisfy functional central limit theorem. Then it is natural to “plug” respective partial sums of these regression residuals to statistic  $DUI(n, m, \alpha)$  and its special variants  $DUI_i(n, m, \alpha)$ ,  $i = 1, 2, 3$ . For the model (4.46) define the partial sums

$$\hat{S}_{kl} = \sum_{i=1}^k \sum_{j=1}^l \hat{u}_{ij}$$

and the statistic

$$\widehat{DUI}(n, m, \alpha) = \max_{\substack{1 \leq a < b \leq n \\ 1 \leq c < d \leq m}} \frac{|\Delta_{b-a}^1 \Delta_{d-c}^2 \widehat{S}_{b,d} - (s_b - s_a)(t_d - t_c) \widehat{S}_{n,m}|}{\max\{s_b - s_a, t_d - t_c\}^\alpha} \quad (4.48)$$

with  $s_k = k/n$  and  $t_l = l/m$ . We have then the following corollary

**Corollary 42** *Under null hypothesis of no change in the regression coefficient  $\beta$  of the model 4.46 given the assumptions F and A*

$$\sigma^{-1}(nm)^{-1/2} \widehat{DUI}(n, m, \alpha) \xrightarrow{D} DUI(\alpha), \text{ as } n \wedge m \rightarrow \infty,$$

for the local alternatives model

$$\beta_{ij} = \beta + \frac{1}{\sqrt{nm}} \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right)$$

with  $\mathbf{g}$  continuous on  $[0, 1]^2$ , we have similar result with the limiting statistic

$$DUI^{loc}(\alpha) = \sup_{0 \leq s < t \leq 1} \frac{|\Delta_{[s,t]} W - \pi(t-s)W(1) + \int_{[s,t]} \mathbf{c}' \mathbf{g} - \pi(t-s) \int_{[0,1]^2} \mathbf{c}' \mathbf{g}|}{|t-s|^\alpha}. \quad (4.49)$$

Note that the limiting statistic remains the same as in the theorem 33. This corollary can be considered as the generalization of the results of Ploberger and Krämer [23] for the regression of double-indexed random variables.

For the regression model (4.47) define partial sums

$$\widehat{S}_{kl}^{FE} = \sum_{i=1}^k \sum_{j=1}^l \widehat{u}_{ij}^{FE}$$

and the statistic

$$\widehat{DUI}^{FE}(n, m, \alpha) = \max_{\substack{1 \leq a < b \leq n \\ 1 \leq c < d \leq m}} \frac{|\Delta_{b-a}^1 \Delta_{d-c}^2 \widehat{S}_{b,d} - (s_b - s_a)(t_d - t_c) \widehat{S}_{n,m}|}{\max\{s_b - s_a, t_d - t_c\}^\alpha}. \quad (4.50)$$

The limiting distribution then is

$$DUI^{FE}(\alpha) = \sup_{0 \leq s < t \leq 1} \frac{|\Delta_{[s,t]} W - (t_2 - s_2)[W(t_1, 1) - W(s_1, 1)]|}{|t-s|^\alpha}.$$

and the corollary

**Corollary 43** *Under null hypothesis of no change in the regression coefficient  $\beta$  of the model (4.24) given the assumptions  $F$  and  $B$*

$$\sigma^{-1}(nm)^{-1/2} \widehat{DUI}^{FE}(n, m, \alpha) \xrightarrow{D} DUI^{FE}(\alpha), \text{ as } n \wedge m \rightarrow \infty,$$

for the local alternatives model

$$\beta_{ij} = \beta + \frac{1}{\sqrt{nm}} \mathbf{g} \left( \frac{i}{n}, \frac{j}{m} \right)$$

with  $\mathbf{g}$  continuous on  $[0, 1]^2$ , we have similar result with the limiting statistic

$$DUI_{loc}^{FE}(\alpha) = \tag{4.51}$$

$$\sup_{0 \leq s < t \leq 1} \frac{|\Delta_{[s,t]} W - (t_2 - s_2) \Delta_{t_1-s_1}^1 W(t_1, 1) + \int_{[s,t]} \mathbf{c}' \mathbf{g} - (t_2 - s_2) \int_{s_1}^{t_1} \int_0^1 \mathbf{c}' \mathbf{g}|}{|t - s|^\alpha}. \tag{4.52}$$

### 4.3.2 Consistency of the epidemic alternatives

Consider that there is a change of the regression coefficient  $\beta$  in rectangle

$$D^* = \left[ \frac{a^*}{n}, \frac{b^*}{n} \right] \times \left[ \frac{c^*}{m}, \frac{d^*}{m} \right],$$

or that the true panel regression models are

$$y_{ij} = \mathbf{x}_{ij}' \beta_0 + \mathbf{x}_{ij}' \mathbf{d}_{ij} + u_{ij} \tag{4.53}$$

$$y_{ij} = \mathbf{x}_{ij}' \beta_0 + \mathbf{x}_{ij}' \mathbf{d}_{ij} + \mu_i + u_{ij}, \tag{4.54}$$

where

$$\mathbf{d}_{ij} = (\beta_1 - \beta_0) \mathbf{1} \left( \left[ \frac{i}{n}, \frac{j}{m} \right] \cap D^* \right).$$

Denote by

$$\mathbf{e}_{n,m} = \left( \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}_{ij}' \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}_{ij}' \mathbf{d}_{ij}$$

and introduce quantity

$$\begin{aligned} \Delta(n, m, D^*) &= \left(1 - \frac{k^* l^*}{nm}\right) \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}'_{ij} \mathbf{d}_{ij} \\ &\quad - \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}'_{ij} \mathbf{e}_{n,m} \mathbf{1} \left( \left[ \frac{i}{n}, \frac{j}{m} \right] \cap D^* \right) + \frac{k^* l^*}{nm} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}'_{ij} \mathbf{e}_{n,m}, \end{aligned}$$

where  $k^* = b^* - a^*$ ,  $l^* = d^* - c^*$  are the lengths of epidemics.

**Theorem 44** *Under alternative hypothesis of the change of the regression coefficient  $\beta$  of the model (4.46)*

$$(nm)^{-1/2} \widehat{DUI}(n, m, \alpha) \rightarrow \infty, \text{ as } n \wedge m \rightarrow \infty,$$

provided

$$(nm)^{-1/2} \max \left\{ \frac{k^*}{n}, \frac{l^*}{m} \right\}^{-\alpha} |\Delta(nm, D^*)| \rightarrow \infty, \text{ as } n \wedge m \rightarrow \infty.$$

*Proof.* The least squares estimate for the model (4.53) satisfies

$$\begin{aligned} \hat{\beta} &= \left( \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}'_{ij} \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}'_{ij} y_{ij} \\ &= \beta_0 + \left( \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} \mathbf{x}'_{ij} \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij} u_{ij} + \mathbf{e}_{n,m}. \end{aligned}$$

The regression residuals then satisfy

$$\hat{u}_{ij}^{alt} = \hat{u}_{ij} + \mathbf{x}'_{ij} \mathbf{d}_{ij} - \mathbf{x}'_{ij} \mathbf{e}_{n,m},$$

where  $\hat{u}_{ij}$  are the regression residuals of the model (4.46). Under alternative hypothesis then

$$\begin{aligned} \widehat{DUI}(n, m, \alpha) &\geq \max \left\{ \frac{k^*}{n}, \frac{l^*}{m} \right\}^{-\alpha} \left| \Delta_{k^*}^1 \Delta_{l^*}^2 \hat{S}_{b^*d}^{alt} - \frac{k^* l^*}{nm} \hat{S}_{n,m}^{alt} \right| \\ &\geq \max \left\{ \frac{k^*}{n}, \frac{l^*}{m} \right\}^{-\alpha} (|\Delta(nm, D^*)| - |T(n, m, D^*)|), \end{aligned}$$

where

$$T(n, m, D^*) = \Delta_{k^*}^1 \Delta_{l^*}^2 \hat{S}_{b^*d} - \frac{k^* l^*}{nm} \hat{S}_{n,m}$$

since

$$(nm)^{-1/2} \max \left\{ \frac{k^*}{n}, \frac{l^*}{m} \right\}^{-\alpha} T(n, m, D^*) = O_P(1),$$

due to corollary 42, the proof is complete.  $\square$

Denote by

$$\mathbf{x}_{ij}^* = \mathbf{x}_{ij} - \frac{1}{m} \sum_{j=c^*}^{d^*} \mathbf{x}_{ij},$$

and define

$$\begin{aligned} \Delta^{FE}(n, m, D^*) &= \left(1 - \frac{k^* l^*}{nm}\right) \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij}'^* \mathbf{d}_{ij} - \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij}'^* \mathbf{e}_{n,m}^{FE} \mathbf{1} \left( \left[ \frac{i}{n}, \frac{j}{m} \right] \cap D^* \right) \\ &\quad + \frac{k^* l^*}{nm} \sum_{i=1}^n \sum_{j=1}^m \mathbf{x}_{ij}'^* \mathbf{e}_{n,m}^{FE}, \end{aligned}$$

where

$$\mathbf{e}_{n,m}^{FE} = \left( \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \tilde{\mathbf{x}}_{ij}' \right)^{-1} \sum_{i=1}^n \sum_{j=1}^m \tilde{\mathbf{x}}_{ij} \mathbf{x}_{ij}'^* \mathbf{d}_{ij}.$$

**Theorem 45** *Under alternative hypothesis of the change of the regression coefficient  $\beta$  of the model (4.47)*

$$(nm)^{-1/2} \widehat{DUI}^{FE}(n, m, \alpha) \rightarrow \infty, \text{ as } n \wedge m \rightarrow \infty,$$

provided

$$(nm)^{-1/2} \max \left\{ \frac{k^*}{n}, \frac{l^*}{m} \right\}^{-\alpha} |\Delta^{FE}(n, m, D^*)| \rightarrow \infty, \text{ as } n \wedge m \rightarrow \infty.$$

*Proof.* Due to definition of  $\mathbf{d}_{ij}$  we have

$$\tilde{y}_{ij} = \tilde{\mathbf{x}}_{ij}' \beta_0 + \mathbf{x}_{ij}'^* \mathbf{d}_{ij} + \tilde{u}_{ij},$$

since  $\hat{\beta}^{FE}$  is the least squares estimate of

$$\tilde{y}_{ij} = \tilde{\mathbf{x}}_{ij}'\beta + \tilde{u}_{ij}.$$

The proof is identical to the proof of the theorem 44 with notation changed.  $\square$

### 4.3.3 Practical considerations and discussion

Note that all of the results in this section relied on the assumption, that regression disturbances satisfy FCLT in Hölder space  $H_\alpha^o([0, 1]^2)$ . From the chapter 3 we know that FCLT holds if disturbances are i.i.d. and satisfy the moment condition

$$\sup_{t>0} t^{1/(1/2-\alpha)} P(|u_{11}| > t) < \infty.$$

For the practical applications the i.i.d. condition sometimes can be too restrictive. On the other hand to lift this restriction we just need to prove the FCLT for wider class of double-indexed random variables. Statistics for testing the change of the regression coefficients remain the same.

Throughout this chapter we focused on constructing statistics for testing against epidemic alternatives. Then the FCLT in Hölder space is needed, since the statistics are the functionals which are continuous only in Hölder space. If in the statistic  $DUI(n, m, \alpha)$  we drop the denominator, then the statistics are continuous functionals in the spaces  $C([0, 1]^2)$  and  $D([0, 1]^2)$ . Then all the results from previous section apply for such statistics given the assumption of the FCLT in  $C([0, 1]^d)$  or  $D([0, 1]^2)$ . The FCLT in  $D([0, 1]^2)$  is proved for wider class of random variables, for strictly stationary multi-indexed random variables satisfying mixing condition by Deo [9] and for strictly stationary multi-indexed martingale differences by Basu and Dorea [4] to name a few. Thus we can apply these types of statistics for wider class of disturbances. In particular if we drop the denominator in the statistic  $DUI_1(n, m, \alpha)$ , our results are then a generalization of Ploberger and Krämer [23].

# Bibliography

- [1] K. Alexander and R. Pyke. A uniform central limit theorem for set-indexed partial-sum processes with finite variance. *Ann. Probab.*, 14:582–597, 1986.
- [2] B. H. Baltagi. *Econometric Analysis of Panel Data*. Wiley, New York, 2001.
- [3] R. F. Bass. Law of the iterated logarithm for set-indexed partial sum processes with finite variance. *Z. Wahrsch. verw. Gebiete*, 70:591–608, 1985.
- [4] A. K. Basu and C. C. Y. Dorea. On functional central limit theorem for stationary martingale random fields. *Acta Mathematica Academiae Scientiarum Hungaricae*, 33(3-4):307–316, 1979.
- [5] P. J. Bickel and M. J. Wichura. Convergence criteria for multiparameter stochastic processes and some applications. *Ann. Math. Statist.*, 42:1656–1670, 1971.
- [6] R. L. Brown, J. Durbin, and J. M. Evans. Techniques for testing the constancy of regression relationships over time. *Journal of the Royal Statistical Society. Series B (Methodological)*, 37(2):149–192, 1975.
- [7] M. Csörgő and L. Horváth. *Limit Theorems in Change-Point Analysis*. John Wiley & Sons, New York, 1997.
- [8] J. Dedecker. Exponential inequalities and functional central limit theorems for random fields. *ESAIM: Probability and Statistics*, 5:77–104, 2001.
- [9] C. M. Deo. A functional central limit theorem for stationary random fields. *The Annals of Probability*, 3(4):708–715, 1975.
- [10] M. Donsker. An invariance principle for certain probability limit theorems. *Mem. Amer. Math. Soc.*, 6:1–12, 1951.

- 
- [11] R. Dudley. Sample functions of the gaussian process. *Ann. Probab.*, 1:66–103, 1973.
  - [12] M. El Machkouri and L. Ouchti. Invariance principles for standard-normalized and self-normalized random fields. *ALEA Latin American Journal of Probability and Mathematical Statistics*, 2006. To appear.
  - [13] R. V. Erickson. Lipschitz smoothness and convergence with applications to the central limit theorem for summation processes. *Ann. Probab.*, 9:831–851, 1981.
  - [14] C. M. Goldie and P. E. Greenwood. Characterisations of set-indexed brownian motion and associated conditions for finite-dimensional convergence. *Ann. Probab.*, 11:802–816, 1986.
  - [15] C. M. Goldie and P. E. Greenwood. Variance of set-indexed sums of mixing random variables and weak convergence of set indexed processes. *Ann. Probab.*, 11:817–839, 1986.
  - [16] C. Hsiao. *Analysis of Panel Data*. Cambridge University Press, New York, 2003.
  - [17] C. Kao, L. Trapani, and G. Urga. Modelling and testing for structural changes in panel cointegration models with idiosyncratic stochastic trend. Working Paper 92, Center for Policy Research, 2007.
  - [18] D. Khoshnevisan. *Multiparameter processes. An introduction to random fields*. Springer Monographs in Mathematics. Springer, New York, 2002.
  - [19] J. Kuelbs. The invariance principle for a lattice of random variables. *Ann. Math. Statist.*, 39:382–389, 1968.
  - [20] J. Lamperti. On convergence of stochastic processes. *Trans. Amer. Math. Soc.*, 104:430–435, 1962.
  - [21] M. Ledoux. Sur une inégalité de H. P. Rosenthal et le théorème limite central dans les espaces de Banach. *Israel J. of Math.*, 50:290–318, 1985.
  - [22] M. Ledoux and M. Talagrand. *Probability in Banach Spaces*. Springer-Verlag, Berlin, Heidelberg, 1991.
  - [23] W. Ploberger and W. Krämer. The cusum test with ols residuals. *Econometrica*, 60(2):271–285, 1992.

- [24] Y. Prokhorov. Convergence of random processes and limit theorems in probability theory. *Theory of Probab. and Appl.*, 1:157–214, 1956.
- [25] R. Pyke. A uniform central limit theorem for partial-sum processes indexed by sets. *London Math. Soc. Lect. Notes Series*, 79:219–240, 1983.
- [26] A. Račkauskas. Hölderian properties of partial sums of regression residuals. *Metrika*, 63:191–205, 2006.
- [27] A. Račkauskas and C. Suquet. Hölder versions of Banach spaces valued random fields. *Georgian Mathematical Journal*, 8(2):347–362, 2001.
- [28] A. Račkauskas and C. Suquet. Necessary and sufficient condition for the Lamperti invariance principle. *Theory of Probability and Mathematical Statistics*, 68:115–124, 2003.
- [29] A. Račkauskas and C. Suquet. Principe d’invariance hölderien pour des tableaux triangulaires de variables aléatoires. *Lietuvos Matematikos Rinkiny*, 43(4):513–532, 2003.
- [30] A. Račkauskas and C. Suquet. Hölderian invariance principle for triangular arrays of random variables. *Lithuanian Mathematical Journal*, 43(4):423–438, 2003.
- [31] A. Račkauskas and C. Suquet. Necessary and sufficient condition for the Hölderian functional central limit theorem. *Journal of Theoretical Probability*, 17(1):221–243, 2004.
- [32] A. Račkauskas and C. Suquet. Hölder norm test statistics for epidemic change. *Journal of Statistical Planning and Inference*, 126(2):495–520, 2004.
- [33] A. Račkauskas and C. Suquet. Central limit theorems in Hölder topologies for Banach space valued random fields. *Theory of Probability and its Applications*, 49(1):109–125, 2004.
- [34] A. Račkauskas and C. Suquet. Testing epidemic changes of infinite dimensional parameters. *Statistical Inference for Stochastic Processes*, 9:111–134, 2006.
- [35] A. Račkauskas, C. Suquet, and V. Zemlys. A Hölderian functional central limit theorem for a multi-indexed summation process. *Stochastic Processes Appl.*, 117(8):1137–1164, 2007.

- [36] A. Račkauskas and V. Zemlys. Functional central limit theorem for a double-indexed summation process. *Liet. Matem. Rink.*, 45:401–412, 2005.
- [37] Z. Semadeni. *Schauder bases in Banach spaces of continuous functions*, volume 918 of *Lecture Notes Math.* Springer Verlag, Berlin, Heidelberg, New York, 1982.
- [38] P. K. Sen. Invariance principles for recursive residuals. *The Annals of Statistics*, 10(1):307–312, 1982.
- [39] C. Suquet. Tightness in Schauder decomposable Banach spaces. In *Amer. Math. Soc. Transl.*, volume 193, pages 201–224. AMS, 1999.
- [40] N. Vakhania, V. Tarieladze, and S. Chobanyan. *Probability distributions on Banach spaces*, volume 14 of *Mathematics and Its Applications (Soviet Series)*. Reidel, Dordrecht, 1987. Transl. from the Russian by Wojbor A. Woyczynski.
- [41] A. van der Vaart and J. Wellner. *Weak convergence and empirical processes*. Springer, New York, 1996.
- [42] M. Wichura. Inequalities with applications to the weak convergence of random processes with multidimensional time parameters. *Ann. Math. Statist.*, 40:681–687, 1969.
- [43] N. Ziegler. Functional central limit theorems for triangular arrays of function-indexed processes under uniformly integrable entropy conditions. *Journal of Multivariate Analysis*, 62:233–272, 1997.